World Journal of *Clinical Cases*

World J Clin Cases 2022 June 6; 10(16): 5124-5517





Published by Baishideng Publishing Group Inc

W J C C World Journal of Clinical Cases

Contents

Thrice Monthly Volume 10 Number 16 June 6, 2022

OPINION REVIEW

5124 Malignant insulinoma: Can we predict the long-term outcomes? Cigrovski Berkovic M, Ulamec M, Marinovic S, Balen I, Mrzljak A

MINIREVIEWS

- 5133 Practical points that gastrointestinal fellows should know in management of COVID-19 Sahin T, Simsek C, Balaban HY
- 5146 Nanotechnology in diagnosis and therapy of gastrointestinal cancer Liang M, Li LD, Li L, Li S
- 5156 Advances in the clinical application of oxycodone in the perioperative period Chen HY, Wang ZN, Zhang WY, Zhu T

ORIGINAL ARTICLE

Clinical and Translational Research

5165 Circulating miR-627-5p and miR-199a-5p are promising diagnostic biomarkers of colorectal neoplasia Zhao DY, Zhou L, Yin TF, Zhou YC, Zhou GYJ, Wang QQ, Yao SK

Retrospective Cohort Study

5185 Management and outcome of bronchial trauma due to blunt versus penetrating injuries Gao JM, Li H, Du DY, Yang J, Kong LW, Wang JB, He P, Wei GB

Retrospective Study

5196 Ovarian teratoma related anti-N-methyl-D-aspartate receptor encephalitis: A case series and review of the literature Li SJ, Yu MH, Cheng J, Bai WX, Di W

- Endoscopic surgery for intraventricular hemorrhage: A comparative study and single center surgical 5208 experience Wang FB, Yuan XW, Li JX, Zhang M, Xiang ZH
- 5217 Protective effects of female reproductive factors on gastric signet-ring cell carcinoma Li Y, Zhong YX, Xu Q, Tian YT
- 5230 Risk factors of mortality and severe disability in the patients with cerebrovascular diseases treated with perioperative mechanical ventilation

Zhang JZ, Chen H, Wang X, Xu K



. .	World Journal of Clinical Cases				
Conten	ts Thrice Monthly Volume 10 Number 16 June 6, 2022				
5241	 Awareness of initiative practice for health in the Chinese population: A questionnaire survey base network platform 				
	Zhang YQ, Zhou MY, Jiang MY, Zhang XY, Wang X, Wang BG				
5253	Effectiveness and safety of chemotherapy for patients with malignant gastrointestinal obstruction: A Japanese population-based cohort study				
	Fujisawa G, Niikura R, Kawahara T, Honda T, Hasatani K, Yoshida N, Nishida T, Sumiyoshi T, Kiyotoki S, Ikeya T, Arai M, Hayakawa Y, Kawai T, Fujishiro M				
	Observational Study				
5266	Long-term outcomes of high-risk percutaneous coronary interventions under extracorporeal membrane oxygenation support: An observational study				
	Huang YX, Xu ZM, Zhao L, Cao Y, Chen Y, Qiu YG, Liu YM, Zhang PY, He JC, Li TC				
5275	Health care worker occupational experiences during the COVID-19 outbreak: A cross-sectional study				
	Li XF, Zhou XL, Zhao SX, Li YM, Pan SQ				
	Prospective Study				
5287	Enhanced recovery after surgery strategy to shorten perioperative fasting in children undergoing non- gastrointestinal surgery: A prospective study				
	Ying Y, Xu HZ, Han ML				
5297	Orthodontic treatment combined with 3D printing guide plate implant restoration for edentulism and its influence on mastication and phonic function				
	Yan LB, Zhou YC, Wang Y, Li LX				
	Randomized Controlled Trial				
5306	Effectiveness of psychosocial intervention for internalizing behavior problems among children of parents with alcohol dependence: Randomized controlled trial				
	Omkarappa DB, Rentala S, Nattala P				
	CASE REPORT				
5317	Crouzon syndrome in a fraternal twin: A case report and review of the literature				
	Li XJ, Su JM, Ye XW				
5324	Laparoscopic duodenoieiunostomy for malignant stenosis as a part of multimodal therapy: A case report				
	Murakami T, Matsui Y				
5331	Chordoma of petrosal mastoid region: A case report				
	Hua JJ, Ying ML, Chen ZW, Huang C, Zheng CS, Wang YJ				
5337	Pneumatosis intestinalis after systemic chemotherapy for colorectal cancer: A case report				
	Liu H, Hsieh CT, Sun JM				
5343	Mammary-type myofibroblastoma with infarction and atypical mitosis-a potential diagnostic pitfall: A case report				
	Zeng YF, Dai YZ, Chen M				



0	World Journal of Clinical Cases
Conten	ts Thrice Monthly Volume 10 Number 16 June 6, 2022
5352	Comprehensive treatment for primary right renal diffuse large B-cell lymphoma with a renal vein tumor thrombus: A case report
	He J, Mu Y, Che BW, Liu M, Zhang WJ, Xu SH, Tang KF
5359	Ectopic peritoneal paragonimiasis mimicking tuberculous peritonitis: A care report
	Choi JW, Lee CM, Kim SJ, Hah SI, Kwak JY, Cho HC, Ha CY, Jung WT, Lee OJ
5365	Neonatal hemorrhage stroke and severe coagulopathy in a late preterm infant after receiving umbilical cord milking: A case report
	Lu Y, Zhang ZQ
5373	Heel pain caused by os subcalcis: A case report
	Saijilafu, Li SY, Yu X, Li ZQ, Yang G, Lv JH, Chen GX, Xu RJ
5380	Pulmonary lymphomatoid granulomatosis in a 4-year-old girl: A case report
	Yao JW, Qiu L, Liang P, Liu HM, Chen LN
5387	Idiopathic membranous nephropathy in children: A case report
	Cui KH, Zhang H, Tao YH
5394	Successful treatment of aortic dissection with pulmonary embolism: A case report
	Chen XG, Shi SY, Ye YY, Wang H, Yao WF, Hu L
5400	Renal papillary necrosis with urinary tract obstruction: A case report
	Pan HH, Luo YJ, Zhu QG, Ye LF
5406	Glomangiomatosis - immunohistochemical study: A case report
	Wu RC, Gao YH, Sun WW, Zhang XY, Zhang SP
5414	Successful living donor liver transplantation with a graft-to-recipient weight ratio of 0.41 without portal flow modulation: A case report
	Kim SH
5420	Treatment of gastric hepatoid adenocarcinoma with pembrolizumab and bevacizumab combination chemotherapy: A case report
	Liu M, Luo C, Xie ZZ, Li X
5428	Ipsilateral synchronous papillary and clear renal cell carcinoma: A case report and review of literature
	Yin J, Zheng M
5435	Laparoscopic radical resection for situs inversus totalis with colonic splenic flexure carcinoma: A case report
	Zheng ZL, Zhang SR, Sun H, Tang MC, Shang JK
5441	PIGN mutation multiple congenital anomalies-hypotonia-seizures syndrome 1: A case report
	Hou F, Shan S, Jin H



0	World Journal of Clinical Cases
Conten	Thrice Monthly Volume 10 Number 16 June 6, 2022
5446	Pediatric acute myeloid leukemia patients with i(17)(q10) mimicking acute promyelocytic leukemia: Two case reports
	Yan HX, Zhang WH, Wen JQ, Liu YH, Zhang BJ, Ji AD
5456	Fatal left atrial air embolism as a complication of percutaneous transthoracic lung biopsy: A case report
	Li YW, Chen C, Xu Y, Weng QP, Qian SX
5463	Diagnostic value of bone marrow cell morphology in visceral leishmaniasis-associated hemophagocytic syndrome: Two case reports
	Shi SL, Zhao H, Zhou BJ, Ma MB, Li XJ, Xu J, Jiang HC
5470	Rare case of hepatocellular carcinoma metastasis to urinary bladder: A case report
	Kim Y, Kim YS, Yoo JJ, Kim SG, Chin S, Moon A
5479	Osteotomy combined with the trephine technique for invisible implant fracture: A case report
	Chen LW, Wang M, Xia HB, Chen D
5487	Clinical diagnosis, treatment, and medical identification of specific pulmonary infection in naval pilots: Four case reports
	Zeng J, Zhao GL, Yi JC, Liu DD, Jiang YQ, Lu X, Liu YB, Xue F, Dong J
5495	Congenital tuberculosis with tuberculous meningitis and situs inversus totalis: A case report
	Lin H, Teng S, Wang Z, Liu QY
5502	Mixed large and small cell neuroendocrine carcinoma of the stomach: A case report and review of literature
	Li ZF, Lu HZ, Chen YT, Bai XF, Wang TB, Fei H, Zhao DB
	LETTER TO THE EDITOR
5510	Pleural involvement in cryptococcal infection
	Georgakopoulou VE, Damaskos C, Sklapani P, Trakas N, Gkoufa A

5515 Electroconvulsive therapy plays an irreplaceable role in treatment of major depressive disorder Ma ML, He LP



Contents

Thrice Monthly Volume 10 Number 16 June 6, 2022

ABOUT COVER

Editorial Board Member of World Journal of Clinical Cases, Shivanshu Misra, MBBS, MCh, MS, Assistant Professor, Surgeon, Department of Minimal Access and Bariatric Surgery, Shivani Hospital and IVF, Kanpur 208005, Uttar Pradesh, India. shivanshu_medico@rediffmail.com

AIMS AND SCOPE

The primary aim of World Journal of Clinical Cases (WJCC, World J Clin Cases) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for WJCC as 1.337; IF without journal self cites: 1.301; 5-year IF: 1.742; Journal Citation Indicator: 0.33; Ranking: 119 among 169 journals in medicine, general and internal; and Quartile category: Q3. The WJCC's CiteScore for 2020 is 0.8 and Scopus CiteScore rank 2020: General Medicine is 493/793.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Xu Guo; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL World Journal of Clinical Cases	INSTRUCTIONS TO AUTHORS https://www.wignet.com/bpg/gerinfo/204	
ISSN	GUIDELINES FOR ETHICS DOCUMENTS	
ISSN 2307-8960 (online)	https://www.wjgnet.com/bpg/GerInfo/287	
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH	
April 16, 2013	https://www.wjgnet.com/bpg/gerinfo/240	
FREQUENCY	PUBLICATION ETHICS	
Thrice Monthly	https://www.wjgnet.com/bpg/GerInfo/288	
EDITORS-IN-CHIEF	PUBLICATION MISCONDUCT	
Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku	https://www.wjgnet.com/bpg/gerinfo/208	
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE	
https://www.wjgnet.com/2307-8960/editorialboard.htm	https://www.wjgnet.com/bpg/gerinfo/242	
PUBLICATION DATE	STEPS FOR SUBMITTING MANUSCRIPTS	
June 6, 2022	https://www.wjgnet.com/bpg/GerInfo/239	
COPYRIGHT	ONLINE SUBMISSION	
© 2022 Baishideng Publishing Group Inc	https://www.f6publishing.com	

© 2022 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: bpgoffice@wjgnet.com https://www.wjgnet.com



W J C C World Journal of Clinical Cases

Submit a Manuscript: https://www.f6publishing.com

World J Clin Cases 2022 June 6; 10(16): 5420-5427

DOI: 10.12998/wjcc.v10.i16.5420

ISSN 2307-8960 (online)

CASE REPORT

Treatment of gastric hepatoid adenocarcinoma with pembrolizumab and bevacizumab combination chemotherapy: A case report

Mei Liu, Cheng Luo, Zong-Zhou Xie, Xun Li

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0 Grade B (Very good): B, B Grade C (Good): C Grade D (Fair): 0 Grade E (Poor): 0

P-Reviewer: Abdelbasset WK, Saudi Arabia; Khaled I, Egypt; Solanki SL; India

Received: September 19, 2021 Peer-review started: September 19, 2021 First decision: December 10, 2021 Revised: December 19, 2021 Accepted: April 22, 2022 Article in press: April 22, 2022 Published online: June 6, 2022



Mei Liu, Zong-Zhou Xie, Xun Li, Department of Medical Oncology, Haikou People's Hospital, Haikou 570100, Hainan Province, China

Cheng Luo, Department of Gynecology, Hainan Hospital of Chinese People's Liberation Army General Hospital, Sanya 572013, Hainan Province, China

Corresponding author: Xun Li, MD, Chief Physician, Department of Medical Oncology, Haikou People's Hospital, No. 43 Renmin Avenue, Haikou 570100, Hainan, China. 1984661032@qq.com

Abstract

BACKGROUND

Gastric hepatoid adenocarcinoma (GHA) is a rare and aggressive cancer that is characterized by foci with features of both hepatocellular differentiation and adenomatous differentiation. However, there is currently no standard treatment for this disease, which has a poor prognosis.

CASE SUMMARY

A 72-year-old male with a body mass index of 20.9 was diagnosed with GHA with perigastric lymph node and liver metastasis. He underwent first-line chemotherapy but that failed. Pembrolizumab and bevacizumab with chemotherapy were used in the second-line treatment. The progression-free survival and overall survival were 14 mo and 16 mo, respectively, after treatment. In addition, the main adverse reaction was tolerable. The patient did not die of tumor progression.

CONCLUSION

The combination of pembrolizumab and bevacizumab with chemotherapy is an effective and safe regimen for GHA and may be recommended as a new choice for GHA treatment. Further studies should evaluate this treatment in a larger cohort or a randomized controlled trial.

Key Words: Gastric hepatoid adenocarcinoma; Serum; Immunotherapy; Targeted therapy; Chemotherapy; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

WJCC | https://www.wjgnet.com

Core Tip: We report a rare case of gastric hepatoid adenocarcinoma in a patient who underwent general chemotherapy that failed. However, pembrolizumab and bevacizumab combination chemotherapy was successful. The overall survival was 16 mo, and the main adverse reaction was tolerable. The patient died of intestinal infection rather than tumor progression.

Citation: Liu M, Luo C, Xie ZZ, Li X. Treatment of gastric hepatoid adenocarcinoma with pembrolizumab and bevacizumab combination chemotherapy: A case report. *World J Clin Cases* 2022; 10(16): 5420-5427 **URL:** https://www.wjgnet.com/2307-8960/full/v10/i16/5420.htm **DOI:** https://dx.doi.org/10.12998/wjcc.v10.i16.5420

INTRODUCTION

Gastric hepatoid adenocarcinoma (GHA) is a primary gastric cancer with the characteristics of adenocarcinoma and hepatocellular carcinoma-like differentiation. GHA is a rare and special subtype of gastric cancer with a clinical incidence rate that is less than or equal to 1%[1-3]. GHA is more common in men. Most patients have high serum AFP levels and are prone to lymph node and liver metastasis. The median overall survival time of GHA is 6-17 mo[4-6]. Some studies have confirmed that the prognosis of patients with GHA is worse than that of patients with common gastric cancer. However, there is currently no standard treatment for this disease.

We report a case of GHA who experienced general chemotherapy failure. Pembrolizumab and bevacizumab combination chemotherapy was successful. The overall survival (OS) was 16 mo, and the main adverse reaction was tolerable.

CASE PRESENTATION

Chief complaints

A 72-year-old male was admitted to our hospital in December 2019 with a 3-mo history of epigastric pain.

History of present illness

Three months before admission, the patient began to experience epigastric pain. He had no abdominal distention, diarrhea, nausea, vomiting, *etc.* Because the symptoms persisted and tended to worsen, the patient visited our hospital for further evaluation.

History of past illness

He had no history of other diseases.

Personal and family history

No family history to note.

Physical examination

Upon physical examination, the abdomen was flat and soft. There was tenderness in the upper abdomen without rebound pain or muscle tension. There was no mass in the abdomen, and there was no swelling of the liver or spleen.

Laboratory examinations

Routine blood examination, blood coagulation function, urinalysis, stool analysis, liver chemistry tests, urea, creatinine, uric acid and electrocardiogram results were all within normal limits. His serum AFP was 339.6 μ g/L on December 5, 2019.

Imaging examinations

An abdominal enhanced computed tomography (CT) scan revealed the following: (1) The gastric fundus and cardia were occupied malignant tumors and multiple lymph node metastases were found around the stomach; and (2) The right anterior lobe of the liver had a low density and was considered metastatic (Figure 1). PET/CT examination showed gastric cancer with perigastric lymph node and liver metastasis. No obvious abnormality was found in the rest of the abdomen.

Zaishidena® WJCC | https://www.wjgnet.com



DOI: 10.12998/wjcc.v10.i16.5420 Copyright ©The Author(s) 2022.



Pathologic findings

A gastroscopy biopsy confirmed a histopathological diagnosis of gastric hepatoid adenocarcinoma. Immunohistochemistry showed HER-2 (-), Ki-67 (+ > 75%), MSH6 (+ > 75%), MSH2 (+ > 75%), PMS2 (+ > 75%), MLH1 (+ > 75%), AFP (+), hepatocytes (+), GPC-3 (+), and SALL4 (+) (Figure 2). Gene detection was performed (2019-12-27, Baishibo, sample type: plasma and paraffin section, SP142), and programmed cell death-ligand 1 (PD-L1) protein expression in paraffin sections was positive. The percentage of positive tumor cells was 0%. The comprehensive positive score was + (5%). In a paraffin section, microsatellites were stable. The TMB values were 8.94 mutations/MB (high) in plasma and 3.81 mutations/MB (moderate) in paraffin sections, with quantiles of 82.36% and 41.9%, respectively. The TP53 Levels were 57.79% in paraffin sections and 11.98% in plasma.

FINAL DIAGNOSIS

The patient was diagnosed with GHA, with a classification of clinical stage IV.

TREATMENT

According to the National Comprehensive Cancer Network guidelines, the patient received first-line treatment of 2 cycles of chemotherapy with oxaliplatin + teggio on December 17, 2019. The doses of oxaliplatin and teggio were 130 mg/m² (Day 1) and 60 mg bid (Days 1-14), respectively. Then, reexamination of abdominal enhanced CT showed the following: (1) the gastric fundus and cardia were occupied, the abdominal cavity and retroperitoneal lymph node showed signs of metastasis, and the previously identified lesions were larger than before; and (2) the range of lesions in the right lobe of the liver was larger than before (Figure 3).

The following evaluation of curative effect is based on the Response Evaluation Criteria in Solid Tumors (RECIST) Version 1.1 standard. We selected liver metastases and the two perigastric malignant lymph nodes (indicated by the arrow in Figure 3) as target lesions, in which the maximum diameter of liver metastases and the minimum diameter of perigastric malignant lymph nodes were measured. The sum of the three was the measurable lesion length. See Table 1 for detailed data. Disease progression (PD) was evaluated. Due to failure of first-line chemotherapy, the effective probability of second-line chemotherapy by itself was not high. GHA has similar components to hepatocellular carcinoma. In recent years, drug treatments for liver cancer have been tried as treatments for GHA. After communicating with the patient and his family and obtaining their consent, we decided to try secondline chemotherapy with pembrolizumab and bevacizumab in February 2020. The doses of epirubicin, albumin binding paclitaxel, pembrolizumab and bevacizumab were 90 mg/m² (Day 1), 260 mg/m² (Day 1), 2 mg/kg (Day 2) and 7.5 mg/kg (Day 0). This was repeated every 3 wk. However, the patient developed 4 degrees of myelosuppression and agranulocytosis with fever and 1 degree of gastrointestinal reaction after the first cycle of the above treatment. He returned to normal soon after symptomatic treatment. Therefore, epirubicin was reduced to 80 mg/m², while other drug doses remained unchanged in cycles 2-4. The above side effects did not reoccur. The patient achieved remission after second-line treatment (Figure 3E vs 3C; Figure 3F vs 3D). After general surgery consultation, surgery was recommended. Therefore, we halted bevacizumab treatment in the 5th cycle (June 2020) and recommended that the patient undergo general surgery for surgical treatment after 3



Table 1 Data on the patient's target lesions								
Date	The minimum diameter of perigastric malignant lymph nodes 1 (cm)	The minimum diameter of perigastric malignant lymph nodes 2 (cm)	The maximum diameter of liver metastases (cm)	The sum (cm)				
December 13, 2019	1.49	2.15	1.82	5.46				
February 7, 2020	2.25	2.82	3.19	8.26				
April 10, 2020	1.54	1.02	1.71	4.27				



DOI: 10.12998/wjcc.v10.i16.5420 Copyright ©The Author(s) 2022.

Figure 2 Histological analysis of the patient's tumour tissue. A: Hematoxylin and eosin staining: tumor cells are arranged in a trabecular patten, with a glandular and hepatoid component (magnification, ×100); B: Hematoxylin and eosin staining: most of them are hepatocyte like differentiation area, and a few are adenocarcinoma area (magnification, ×200); C: Immunohistochemical staining: cells are positively stained for SALL4 (magnification, ×100).

> wk. However, due to his advanced age, he did not follow the doctor's advice undergo surgery. In August 2020, he received pembrolizumab by itself for the last time. However, he received no further treatment and recuperated at home for personal reasons. However, in January 2021, he developed diarrhea with fever and could not eat normally. His body mass index dropped to 18. Thus, he returned to the hospital. PET/CT examination showed that the tumor was still stable (Figure 4).

> Laboratory examinations showed that his inflammatory indices were high, leukocytes and neutrophils were slightly low, and other measures, such as thyroid function and the presence hepatitis B virus, were normal. However, the patient refused to allow examination of the stool routine, culture, etc., thus preventing the collection of etiological evidence. The possibility of intestinal infection was considered in the diagnosis. The patient was given parenteral nutrition, anti-infection therapy, an indwelling gastric tube for enteral nutrition and other supportive treatment. After the symptoms improved, the patient was discharged and returned to his hometown.

OUTCOME AND FOLLOW-UP

The patient's family reported that the patient had diarrhea with fever again in March 2021. He was treated in a local hospital but died in April 2021. The cause of death was intestinal infection. The last telephone follow-up was June 18, 2021. The patient, whose progression-free survival (PFS) and OS were



Zeishidene® WJCC | https://www.wjgnet.com



DOI: 10.12998/wjcc.v10.i16.5420 Copyright ©The Author(s) 2022.

Figure 3 Imaging examinations before and after first-line and second-line treatments. A and C: Curative effect of perigastric lymph nodes: PD; B and D: Curative effect of liver metastases: PD; C and E: Curative effect of perigastric lymph nodes: PR; D and F: Curative effect of liver metastases: PR.

> 14 mo and 16 mo, achieved remission after second-line treatment. The main adverse reaction of the treatment was tolerable. He died of intestinal infection rather than tumor progression.

DISCUSSION

The incidence rate of GHA is low. In addition, there is no standard treatment. Most of the treatment methods follow the principles of general gastric cancer. Surgery and chemotherapy are the main treatments. Molecular targeting therapy and immunotherapy are also being explored. D2 radical resection is the first choice for patients with early-stage GHA. For patients with isolated liver metastasis, palliative gastrectomy plus simultaneous resection of liver metastasis can be considered. Palliative gastrectomy plus local treatment of liver metastasis, such as hepatic artery chemoembolization or radiofrequency ablation, can also be considered. Chemotherapy is the main treatment for patients with advanced GHA that cannot be removed by surgery. Related studies have indicated that the first-line standard chemotherapy plan for GHA includes 5-FU and platinum-based chemotherapy, combined with simultaneous Taxol, irinotecan, methotrexate, mitomycin-C and other chemotherapy [3,7-8]. With the development of molecular detection technology, molecular targeted therapy has also been a topic of major interest in recent years. Trastuzumab combined with chemotherapy in the first-line treatment of HER2-positive common gastric cancer achieved positive results in a phase 3 Large-scale clinical study (ToGA study). Ranuciumab, which is a VEGFR2 antibody, has been approved as the first antiangiogenic drug for the treatment of advanced common gastric cancer^[8]. In this case, the patient's HER-2 status was negative; thus, his cancer was not suitable for anti-Her-2 treatment. In addition, because of drug accessibility, ramucirumab-targeted therapy was not carried out. SOX chemotherapy was carried out as the first-line treatment, but it failed. The selection of the second-line treatment was based on the following four considerations: (1) The patient was diagnosed with GHA, had eating difficulties (medication was inconvenient) and had similar components of hepatocellular carcinoma. However, chemotherapy alone was ineffective. According to the literature, multiple targeted antiangiogenic drugs, such as apatinib and sorafenib[9-10], can be used to treat clinical hepatoid adenocarcinoma; (2) IMbrave150, a phase 3 clinical trial of liver cancer[11], showed that the OS and PFS of atezolizumab + bevacizumab in unresectable HCC patients without previous systemic therapy were statistically and clinically significant and improved compared with those of sorafenib, with controllable safety; (3) PD-L1 protein expression was positive in this patient, with a TMB of 8.94 mutations/MB (high) in plasma and 3.81 mutations/MB in paraffin sections; and (4) Because of the accessibility of atezolizumab and the patient's financial means, we tried a new kind of "T + A"-like combined chemotherapy: pembrolizumab + bevacizumab combined chemotherapy. Ultimately, the patient achieved an extended PFS. The main adverse reaction was hematological toxicity, which was tolerated. Unfortunately, the patient died of complications: intestinal infection, not tumor progression. His OS might have been longer.





Figure 4 PET- computed tomography examinations at different time after second-line treatment. Persistent remission of perigastric lymph nodes.

Immunotherapy treatment of GHA is relatively understudied. However, a recent phase III randomized clinical trial, Keynote-062, showed that pembrolizumab was not inferior to chemotherapy for untreated advanced gastric/gastroesophageal junction cancer. In addition, fewer adverse events were observed. pembrolizumab or pembrolizumab plus chemotherapy was not superior to chemotherapy for OS and PFS endpoints[12]. However, the results of the Checkmate 649 study showed that for patients with advanced gastric cancer/gastroesophageal junction cancer/esophageal adenocarcinoma who had not been treated in the past, nivolumab was the first PD-1 inhibitor that was clinically shown to be superior to chemotherapy alone in terms of OS and PFS, with controllable safety [13]. The first-line chemotherapy treatment failed in this patient. However, the second-line attempt of pembrolizumab + bevacizumab combined chemotherapy led to tumor remission, and the side effects were tolerable. Consistent with the results of the Checkmate 649 study, pembrolizumab + bevacizumab combined chemotherapy may be effective. However, in the Keynote-062 clinical trial, pembrolizumab + chemotherapy was not superior to chemotherapy alone for OS and PFS. Possible explanations are as follows: (1) The lack of a synergistic effect of anti-angiogenesis targeted drugs such as bevacizumab; and (2) first-line chemotherapy may increase antigen exposure, and as a result, the benefits of second-line immunotherapy. However, these possible explanations are only speculation; there is no current evidence.

Previous studies have shown that surgery, chemotherapy and targeted therapy can be used in patients with GHA. However, the application of immunotherapy in such patients has not been reported in the literature. In this case of GHA, we tried a new regimen of pembrolizumab and bevacizumab with chemotherapy, and the patient benefited. However, the sample size of this was very small. Further studies should evaluate this treatment in a larger cohort or a randomized controlled trial.

Baisbideng® WJCC | https://www.wjgnet.com

CONCLUSION

The combination of pembrolizumab and bevacizumab with chemotherapy is an effective and safe regimen for treating this GHA patient. However, the sample size of this study was very small. Further evaluation of this treatment in a larger cohort or a randomized controlled trial is needed.

FOOTNOTES

Author contributions: Liu M contributed to the design, analysis, and drafted the manuscript; Li X contributed to the analysis, and critically revised the manuscript; Luo C contributed to the analysis; Xie ZZ collected medical history information, and edited charts; and all authors read and approved the final manuscript.

Informed consent statement: Informed written consent was obtained from the patient for publication of this report and any accompanying images.

Conflict-of-interest statement: The authors declare that they have no conflict of interest.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is noncommercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country/Territory of origin: China

ORCID number: Mei Liu 0000-0002-8863-6577; Cheng Luo 0000-0001-9275-8772; Zong-Zhou Xie 0000-0002-1533-7189; Xun Li 0000-0002-1663-170X.

S-Editor: Ma YJ L-Editor: A P-Editor: Ma Y]

REFERENCES

- Zhao M, Sun L, Lai JZ, Shi H, Mei K, He X, Jin X, Lai J, Cao D. Expression of RNA-binding protein LIN28 in classic 1 gastric hepatoid carcinomas, gastric fetal type gastrointestinal adenocarcinomas, and hepatocellular carcinomas: An immunohistochemical study with comparison to SALL4, alpha-fetoprotein, glypican-3, and Hep Par1. Pathol Res Pract 2018; 214: 1707-1712 [PMID: 30196987 DOI: 10.1016/j.prp.2018.07.037]
- 2 Lin CY, Yeh HC, Hsu CM, Lin WR, Chiu CT. Clinicopathologial features of gastric hepatoid adenocarcinoma. Biomed J 2015; 38: 65-69 [PMID: 25163499 DOI: 10.4103/2319-4170.126860]
- Zhang ZR, Wu J, Li HW, Wang T. Hepatoid adenocarcinoma of the stomach: Thirteen case reports and review of 3 literature. World J Clin Cases 2020; 8: 1164-1171 [PMID: 32258088 DOI: 10.12998/wjcc.v8.i6.1164]
- Wang Y, Sun L, Li Z, Gao J, Ge S, Zhang C, Yuan J, Wang X, Li J, Lu Z, Gong J, Lu M, Zhou J, Peng Z, Shen L, Zhang X. Hepatoid adenocarcinoma of the stomach: a unique subgroup with distinct clinicopathological and molecular features. Gastric Cancer 2019; 22: 1183-1192 [PMID: 30989433 DOI: 10.1007/s10120-019-00965-5]
- Chen EB, Wei YC, Liu HN, Tang C, Liu ML, Peng K, Liu T. Hepatoid Adenocarcinoma of Stomach: Emphasis on the Clinical Relationship with Alpha-Fetoprotein-Positive Gastric Cancer. Biomed Res Int 2019; 2019: 6710428 [PMID: 31915699 DOI: 10.1155/2019/6710428]
- 6 Yang J, Wang R, Zhang W, Zhuang W, Wang M, Tang C. Clinicopathological and prognostic characteristics of hepatoid adenocarcinoma of the stomach. Gastroenterol Res Pract 2014; 2014: 140587 [PMID: 24669215 DOI: 10.1155/2014/140587
- Zou M, Li Y, Dai Y, Sun L, Huang T, Yuan X, Qiu H. AFP-producing hepatoid adenocarcinoma (HAC) of peritoneum and 7 omentum: a case report and literature review. Onco Targets Ther 2019; 12: 7649-7654 [PMID: 31571915 DOI: 10.2147/OTT.S216501
- Soreide JA. Therapeutic Approaches to Gastric Hepatoid Adenocarcinoma: Current Perspectives. Ther Clin Risk Manag 8 2019; 15: 1469-1477 [PMID: 31920320 DOI: 10.2147/TCRM.S204303]
- Gavrancic T, Park YH. A novel approach using sorafenib in alpha fetoprotein-producing hepatoid adenocarcinoma of the lung. J Natl Compr Canc Netw 2015; 13: 387-91; quiz 391 [PMID: 25870375 DOI: 10.6004/jnccn.2015.0054]
- 10 Petrelli F, Ghilardi M, Colombo S, Stringhi E, Barbara C, Cabiddu M, Elia S, Corti D, Barni S. A rare case of metastatic pancreatic hepatoid carcinoma treated with sorafenib. J Gastrointest Cancer 2012; 43: 97-102 [PMID: 21365478 DOI: 10.1007/s12029-011-9264-2
- Ducreux MP, Cheng AL, Qin S, Zhu AX, Ikeda M, Kim TY, Xu DZ, Verret W, Liu J, Finn RS, Galle PR. Atezolizumab + 11



bevacizumab vs sorafenib in locally advanced or metastatic hepatocellular carcinoma: The randomised phase III study IMbrave150. Annals of Oncology 2018; 29 (suppl 8) [DOI: 10.1093/annonc/mdy282.165]

- 12 Shitara K, Van Cutsem E, Bang YJ, Fuchs C, Wyrwicz L, Lee KW, Kudaba I, Garrido M, Chung HC, Lee J, Castro HR, Mansoor W, Braghiroli MI, Karaseva N, Caglevic C, Villanueva L, Goekkurt E, Satake H, Enzinger P, Alsina M, Benson A, Chao J, Ko AH, Wainberg ZA, Kher U, Shah S, Kang SP, Tabernero J. Efficacy and Safety of Pembrolizumab or Pembrolizumab Plus Chemotherapy vs Chemotherapy Alone for Patients With First-line, Advanced Gastric Cancer: The KEYNOTE-062 Phase 3 Randomized Clinical Trial. JAMA Oncol 2020; 6: 1571-1580 [PMID: 32880601 DOI: 10.1001/jamaoncol.2020.3370]
- 13 Moehler M, Shitara K, Garrido M, Salman P, Shen L, Wyrwicz L, Yamaguchi K, Skoczylas T, Bragagnoli AC, Liu T, Schenker M, Yanez P, Tehfe M, Poulart V, Cullen D, Lei M, Kondo K, Li M, Ajani JA, Janjigian YY. Nivolumab (nivo) plus chemotherapy (chemo) vs chemo as first-line (1L) treatment for advanced gastric cancer/gastroesophageal junction cancer (GC/GEJC)/esophageal adenocarcinoma (EAC): first results of the CheckMate 649 study. Annals Oncol 2020; 31 (suppl 4): S1142-S1215 [DOI: 10.1016/j.annonc.2020.08.2296]





Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: bpgoffice@wjgnet.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

