

World Journal of *Gastrointestinal Oncology*

World J Gastrointest Oncol 2022 April 15; 14(4): 748-946



REVIEW

- 748 Regulatory RNAs, microRNA, long-non coding RNA and circular RNA roles in colorectal cancer stem cells
Chao HM, Wang TW, Chern E, Hsu SH
- 765 Role of three-dimensional printing and artificial intelligence in the management of hepatocellular carcinoma: Challenges and opportunities
Christou CD, Tsoulfas G
- 794 Role of sirtuins in esophageal cancer: Current status and future prospects
Otsuka R, Hayano K, Matsubara H

MINIREVIEWS

- 808 Vasoactive intestinal peptide secreting tumour: An overview
Una Cidon E
- 820 Management of single pulmonary metastases from colorectal cancer: State of the art
Chiappetta M, Salvatore L, Congedo MT, Bensi M, De Luca V, Petracca Ciavarella L, Camarda F, Evangelista J, Valentini V, Tortora G, Margaritora S, Lococo F
- 833 Current guidelines in the surgical management of hereditary colorectal cancers
Kudchadkar S, Ahmed S, Mukherjee T, Sagar J

ORIGINAL ARTICLE**Basic Study**

- 842 Berberine retarded the growth of gastric cancer xenograft tumors by targeting hepatocyte nuclear factor 4 α
Li LL, Peng Z, Hu Q, Xu LJ, Zou X, Huang DM, Yi P
- 858 Bi-specific T1 positive-contrast-enhanced magnetic resonance imaging molecular probe for hepatocellular carcinoma in an orthotopic mouse model
Ma XH, Chen K, Wang S, Liu SY, Li DF, Mi YT, Wu ZY, Qu CF, Zhao XM
- 872 Xihuang pills induce apoptosis in hepatocellular carcinoma by suppressing phosphoinositide 3-kinase/protein kinase-B/mechanistic target of rapamycin pathway
Teng YJ, Deng Z, Ouyang ZG, Zhou Q, Mei S, Fan XX, Wu YR, Long HP, Fang LY, Yin DL, Zhang BY, Guo YM, Zhu WH, Huang Z, Zheng P, Ning DM, Tian XF

Retrospective Study

- 887 Effect of hepatic artery resection and reconstruction on the prognosis of patients with advanced hilar cholangiocarcinoma
Li YM, Bie ZX, Guo RQ, Li B, Wang CE, Yan F

- 897** Prognostic significance of serum inflammation indices for different tumor infiltrative pattern types of gastric cancer

Wang YF, Yin X, Fang TY, Wang YM, Zhang L, Zhang XH, Zhang DX, Zhang Y, Wang XB, Wang H, Xue YW

- 920** Regorafenib combined with programmed cell death-1 inhibitor against refractory colorectal cancer and the platelet-to-lymphocyte ratio's prediction on effectiveness

Xu YJ, Zhang P, Hu JL, Liang H, Zhu YY, Cui Y, Niu P, Xu M, Liu MY

Clinical Trials Study

- 935** Genome-wide methylation profiling of early colorectal cancer using an Illumina Infinium Methylation EPIC BeadChip

Wu YL, Jiang T, Huang W, Wu XY, Zhang PJ, Tian YP

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Oncology*, Hanlin L Wang, MD, PhD, Professor, Department of Pathology and Laboratory Medicine, University of California Los Angeles, David Geffen School of Medicine and Ronald Reagan UCLA Medical Center, Los Angeles, CA 90095, United States.
hanlinwang@mednet.ucla.edu

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Oncology* (*WJGO*, *World J Gastrointest Oncol*) is to provide scholars and readers from various fields of gastrointestinal oncology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGO mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal oncology and covering a wide range of topics including liver cell adenoma, gastric neoplasms, appendiceal neoplasms, biliary tract neoplasms, hepatocellular carcinoma, pancreatic carcinoma, cecal neoplasms, colonic neoplasms, colorectal neoplasms, duodenal neoplasms, esophageal neoplasms, gallbladder neoplasms, etc.

INDEXING/ABSTRACTING

The *WJGO* is now indexed in Science Citation Index Expanded (also known as SciSearch®), PubMed, PubMed Central, and Scopus. The 2021 edition of Journal Citation Reports® cites the 2020 impact factor (IF) for *WJGO* as 3.393; IF without journal self cites: 3.333; 5-year IF: 3.519; Journal Citation Indicator: 0.5; Ranking: 163 among 242 journals in oncology; Quartile category: Q3; Ranking: 60 among 92 journals in gastroenterology and hepatology; and Quartile category: Q3. The *WJGO*'s CiteScore for 2020 is 3.3 and Scopus CiteScore rank 2020: Gastroenterology is 70/136.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xiang Li*; Editorial Office Director: *Ya-Juan Ma*.

NAME OF JOURNAL

World Journal of Gastrointestinal Oncology

ISSN

ISSN 1948-5204 (online)

LAUNCH DATE

February 15, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Monjur Ahmed, Florin Burada

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-5204/editorialboard.htm>

PUBLICATION DATE

April 15, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Regorafenib combined with programmed cell death-1 inhibitor against refractory colorectal cancer and the platelet-to-lymphocyte ratio's prediction on effectiveness

Yu-Jie Xu, Peng Zhang, Jin-Long Hu, Hong Liang, Yan-Yan Zhu, Yao Cui, Po Niu, Min Xu, Ming-Yue Liu

Specialty type: Oncology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Cho YS, South Korea; Shomura M, Japan

Received: October 9, 2021

Peer-review started: October 9, 2021

First decision: December 12, 2021

Revised: January 4, 2022

Accepted: March 25, 2022

Article in press: March 25, 2022

Published online: April 15, 2022



Yu-Jie Xu, Jin-Long Hu, Yan-Yan Zhu, Yao Cui, Po Niu, Ming-Yue Liu, Department of Oncology, Henan Provincial People's Hospital, People's Hospital of Zhengzhou University, Zhengzhou 450003, Henan Province, China

Peng Zhang, Hong Liang, Department of Gastrointestinal Surgery, Henan Provincial People's Hospital, People's Hospital of Zhengzhou University, Zhengzhou 450003, Henan Province, China

Min Xu, Department of Hepatology, The Third People's Hospital of Zhengzhou, Zhengzhou 450003, Henan Province, China

Corresponding author: Ming-Yue Liu, Doctor, MD, PhD, Chief Doctor, Department of Oncology, Henan Provincial People's Hospital, People's Hospital of Zhengzhou University, No. 7 Weiwu Road, Zhengzhou 450003, Henan Province, China. 1816160187@e.gzhu.edu.cn

Abstract

BACKGROUND

The effectiveness of regorafenib plus programmed cell death-1 (PD-1) inhibitor in treating microsatellite stable (MSS) metastatic colorectal cancer (mCRC) remains controversial.

AIM

To investigate the benefits of regorafenib combined with PD-1 inhibitor in treating MSS mCRC and explore indicators predicting response.

METHODS

This retrospective study included a total of 30 patients with microsatellite stable metastatic colorectal cancer treated with regorafenib combined with programmed cell death-1 inhibitor at Henan Provincial People's Hospital between December 2018 and December 2020. During a 4-wk treatment cycle, regorafenib was performed for 3 continuous weeks. PD-1 inhibitor was intravenously injected starting on the first day of the oral intake of regorafenib. We reviewed tumor response, progression-free survival (PFS), overall survival, and treatment-related adverse events (TRAEs) and evaluated association between platelet-to-lymphocyte ratio (PLR) and outcomes in this retrospective study.

RESULTS

Stable disease and progressive disease were found in 18 (60.0%) and 12 (40.0%) patients, respectively. The disease control rate was 60.0%. The median follow-up time was 12.0 mo, and median PFS was 3.4 mo [95% confidence interval (CI): 2.2-4.6 mo]. Of the 12 patients with progressive disease, 10 (83.3%) had liver metastasis before starting the combined treatment. Among the 18 patients with SD, 10 (55.6%) did not have liver metastases. One patient without liver metastases at baseline was found with a substantially prolonged PFS of 11.2 mo. The liver metastasis, the choice of programmed cell death-1 inhibitor other than nivolumab or pembrolizumab and previous exposure to regorafenib was't associated with treatment outcome. The median PFS in the low-PLR group was 4.2 mo (95%CI: 3.5-4.9 mo), compared with 2.8 mo (95%CI: 1.4-4.2 mo) in the high-PLR group ($P = 0.005$). The major TRAEs included hand-foot syndrome (33.3%), hypertension (23.3%), malaise (20.0%), and gastrointestinal reaction (16.7%). The incidence of grade 3 TRAEs was 13.3% (4/30), which comprised abnormal capillary proliferation ($n = 1$), transaminase elevation ($n = 1$), and hand-foot syndrome ($n = 2$). No grade 4 or higher toxicity was observed.

CONCLUSION

Regorafenib combined with PD-1 inhibitor could lead to a longer PFS in some patients with MSS mCRC. The PLR might be a prediction of the patient response to this therapy.

Key Words: Colorectal neoplasms; Microsatellite stable; Programmed cell death-1 inhibitor; Platelet-to-lymphocyte ratio; Regorafenib; Progression-free survival

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The use of regorafenib combined with programmed cell death-1 inhibitor in the treatment of refractory microsatellite stable colorectal cancer has contradictory results in some small-scale studies. The purpose of this paper is to analyze the real-world data of our center in the past 2 years so as to provide more treatment experience and reference for treatment selection. The progression-free survival and overall survival of patients with refractory microsatellite stable colorectal cancer treated with regorafenib combined with programmed cell death-1 inhibitor were analyzed retrospectively, and the safety and adverse reactions under different doses were reviewed. The platelet-to-lymphocyte ratio was found as a potential screening index for patients with prolonged progression-free survival.

Citation: Xu YJ, Zhang P, Hu JL, Liang H, Zhu YY, Cui Y, Niu P, Xu M, Liu MY. Regorafenib combined with programmed cell death-1 inhibitor against refractory colorectal cancer and the platelet-to-lymphocyte ratio's prediction on effectiveness. *World J Gastrointest Oncol* 2022; 14(4): 920-934

URL: <https://www.wjgnet.com/1948-5204/full/v14/i4/920.htm>

DOI: <https://dx.doi.org/10.4251/wjgo.v14.i4.920>

INTRODUCTION

Colorectal cancer (CRC) is one of the most common malignant tumors worldwide. More than 1.8 million new patients with CRC were reported in 2018, of which 881000 died of CRC[1,2]. In China, CRC is the fifth leading cause of cancer-related death. The number of CRC-related deaths was about 191000 in 2015 [3]. A variety of patients are diagnosed with advanced CRC, accompanied by distant metastases in addition to the primary tumor[4]. Despite multidisciplinary management based on surgery, systemic therapy, and radiotherapy[5-7], the prognosis of patients with advanced CRC is still poor, with 5-year survival rates of 71% and 14% for regional and distant disease, respectively[8].

The Guidelines of the Chinese Society of Clinical Oncology (CSCO) recommend chemotherapy with or without targeted therapy (such as cetuximab and bevacizumab) for the first- and second-line therapies of advanced CRC[9]. Although the CSCO guidelines also recommended later-line therapy for CRC after progression, the treatment efficacy is generally limited[9]. Regorafenib is an oral, small-molecular multi-target kinase inhibitor that can exert anti-tumor effects through inhibiting several key processes, such as tumor cell proliferation, metastasis, angiogenesis, and immune escape[10,11]. The international CORRECT trial promoted regorafenib as the standard drug for treating metastatic CRC (mCRC)[12].

The emergence of immunotherapy in recent years has brought long-term survival benefits for many patients. Still, to date, only patients with microsatellite instability-high mCRC could benefit from immunotherapy using a single drug[13]. Considering that the efficacy of single targeted drug was suboptimal, combination therapy could bring new hopes for patients. The preliminary findings from the REGONIVO study reported the efficacy of regorafenib combined with nivolumab. The objective response rate (ORR) was 33% in 24 Asian patients with proficient mismatch repair (MMR)/microsatellite stable (MSS) refractory mCRC, and the median progression-free survival (PFS) was prolonged by more than 6 mo. Therefore, the findings possibly provided new chances for patients with repeated failures after third- or further-line therapies to prolong the survival time and improve the quality of life[14]. The combination of regorafenib with programmed cell death-1 (PD-1) for patients with refractory CRC after multi-line standard therapies is now used by many medical centers worldwide. A retrospective study in 18 patients with refractory MSS mCRC, including five Asian patients, performed by the National Cancer Institute in the United States recently failed to show the effectiveness of regorafenib combined with nivolumab or pembrolizumab but suggested that patients without liver metastases could benefit from the treatment[15]. A study conducted in 23 Chinese patients with MSS advanced CRC also did not demonstrate the effectiveness of this combination[16]. These two studies[15,16] were inconsistent with the REGONIVO study[14]. Hence, additional clinical studies are needed to investigate the benefits and adverse events of regorafenib combined with PD-1 inhibitors.

The effects of specific treatments among different patients and different studies vary greatly. Thus, identifying factors that could predict the prognosis of patients treated with regorafenib combined with PD-1 inhibitors is clinically relevant. Carcinoembryonic antigen (CEA) is commonly used in clinical practice to monitor treatment efficacy in patients with CRC[17,18]. Other indicators are also related to the prognosis of patients with cancer[19-23], such as neutrophil-to-lymphocyte ratio (NLR) and platelet-to-lymphocyte ratio (PLR).

Despite the emergence of new drugs and the changes in the combination of targeted therapy, the effectiveness of mCRC treatment is still suboptimal, especially the treatments after third-line therapy, with a 5-year survival rate of only 11%[24]. Therefore, this retrospective study investigated patients with MSS mCRC treated with regorafenib combined with a PD-1 inhibitor in the last 2 years. This study may suggest a novel therapeutic approach that could be tried in a clinical trial.

MATERIALS AND METHODS

Study design and patients

This retrospective study included patients with MSS mCRC treated with regorafenib combined with PD-1 inhibitor at Henan Provincial People's Hospital between December 2018 and December 2020. This study was approved by the ethics committee of People's Hospital of Zhengzhou University (Henan Province, China) and performed in accordance with the Declaration of Helsinki. The requirement for informed consent was waived by the committee because of the retrospective nature of the study.

The inclusion criteria were: (1) Histologically or cytologically proven with MSS mCRC; (2) Treated with more than two lines of standard chemotherapy regimens (including fluorouracil, oxaliplatin, and irinotecan, with or without biological agents such as bevacizumab and cetuximab); (3) Treated with regorafenib combined with PD-1 inhibitor (due to the accessibility of drugs and financial burden of patients, other low-cost PD-1 inhibitors approved in China, such as camrelizumab, sintilimab, toripalimab, and tislelizumab, could also be used in addition to nivolumab and pembrolizumab); (4) With evaluable lesions and with detailed clinical data and follow-up results; and (5) Downregulation of the expression of four MMR enzymes (MutL homolog 1/MutS homolog 2/MutS homolog 6/PMS1 homolog 2) assessed by immunohistochemistry, or the MMR/MSI status was evaluated by the 2B3D method.

Treatment regimen

During a 4 wk treatment cycle, oral drug administration was performed for 3 continuous weeks. Regorafenib was orally administered at 80, 120, or 160 mg once per day. Dose reduction or temporary discontinuation of regorafenib was performed for the patients with treatment-related toxicities. PD-1 inhibitor was intravenously injected starting on the 1st day of the oral intake of regorafenib, including 240 mg every 3 wk for toripalimab, 200 mg every 2 or 3 wk for camrelizumab, 240 mg every 2 wk for nivolumab, 200 mg every 3 wk for pembrolizumab and sintilimab, or 200 mg every 3 wk for tislelizumab.

Data collection

Collected data included age, sex, Eastern Cooperative Oncology Group (ECOG) performance status (PS), site of the primary tumor, site of the metastases, MSI/MMR, gene status, lines of treatment, and previous treatments. The blood routine examination and CEA results before treatment and after three and five cycles of combination therapy were longitudinally analyzed. CEA was detected by microarray chemiluminescence immunoassay (Sunlant Biological SLXP-001, Wuxi, Jiangsu Province, China). The

platelet and lymphocyte counts were measured using a Hessian-Meikang XN-9000 automatic modular blood and body fluid analyzer. The PLR was calculated as the absolute platelet count divided by the absolute lymphocyte count in full blood.

According to the Response Evaluation Criteria in Solid Tumors (RECIST) version 1.1, tumor responses were evaluated every two or three cycles of immunotherapy. If signs of rapid disease progression were noted, the evaluation was performed at earlier time points. ORR included complete response (CR) and partial response (PR). Disease control rate (DCR) was defined as the sum of the ORR and stable disease (SD) rate. PFS was defined as the time from the start of treatment to disease progression or death. Overall survival (OS) referred to the time from the start of treatment to death. Toxicity was evaluated according to the Common Toxicity Criteria for Adverse Events Version 5.0 (CTCAE 5.0) and the 2019 CSCO immune checkpoint inhibitor-related toxicity management guideline [25]. Follow-up was censored on April 15, 2021.

Statistical analysis

SPSS 26.0 (IBM, Armonk, NY, United States) was used for statistical analysis. Continuous data with a normal distribution (according to the Shapiro-Wilk test) were described using means \pm SD, while continuous data without a normal distribution were described as medians and interquartile range (Q1, Q3). The t-test was used for comparing continuous data with a normal distribution, and the rank-sum test was used for comparing continuous data without a normal distribution. Categorical data were presented as *n* (%) and tested using the χ^2 test. The receiver operating characteristic (ROC) curve was used to estimate the best cutoff value of NLR and PLR before treatment. The patients were divided into the low-NLR, high-NLR, low-PLR, and high-PLR groups according to the cut-off value of ROC curve. The Kaplan-Meier method was used for univariable survival analysis. The log-rank test was used to analyze the differences in PFS and OS among different influencing factors. Two-sided *P* values < 0.05 were considered statistically significant.

RESULTS

Characteristics of the patients

Between December 2018 and December 2020, a total of 30 patients with MSS mCRC were included. **Table 1** and **Supplementary Table 1** present the characteristics of the patients. The patients were treated with regorafenib combined with PD-1 inhibitor as the third- (46.7%) or further-line therapy (53.3%). Of these patients, 12 (40.0%) were \geq 60 years and 18 (60.0%) < 60 years. Fourteen (46.7%) and sixteen (53.3%) were males and females, respectively. Twenty-one (70.0%) and 9 (30.0%) patients had an ECOG PS of 0-1 and \geq 2, respectively. The primary lesion was on the left side (including the left colon, sigmoid colon, and rectum) in 24 patients (80.0%) and on the right side (including the ascending colon and liver curvature of the transverse colon) in 6 patients (20.0%). Eighteen (60.0%) patients had liver metastases, 11 (36.7%) had lung metastases, 6 (20.0%) had peritoneal metastases, and 18 patients (60.0%) had multiple metastases in \geq two sites (**Table 1**). All patients were proven with MSS (proficient MMR) type CRC using immunohistochemistry or polymerase chain reaction, and 1 (3.3%) patient had been treated with PD-1 inhibitor before. The genetic test showed that 7 (36.7%) and 12 (60.0%) patients had RAS mutation and wild-type RAS, respectively, while 11 patients did not have a genetic mutation test. No patient had BRAF mutation.

Characteristics of the treatments

Nineteen (63.3%) patients were treated with regorafenib for a median treatment duration of 4.3 mo [95% confidence interval (CI): 2.0-6.7 mo] before the combination therapy in this study. During the combination therapy, the initial dose of regorafenib was 80, 120, and 160 mg in 18, 8, and 4 patients, respectively. Of the 30 patients receiving combination therapy, 9 were treated with sintilimab, 8 with toripalimab, 5 with camrelizumab, 4 with tislelizumab, 3 with nivolumab, and 1 with pembrolizumab (**Table 1** and **Supplementary Table 1**).

Clinical response

According to RECIST 1.1, 18 (60.0%) of the 30 patients who received combination therapy achieved SD, and 12 (40.0%) had progressive disease (PD); no PR or CR was found. The DCR was 60.0% (**Table 2** and **Supplementary Table 1**). Of the 12 patients with PD, 10 (83.3%) had liver metastases at baseline. Among the 18 patients with SD, 10 (55.6%) had no liver metastases.

Survival

Up to April 15, 2021, the median follow-up time was 12.0 mo. The median PFS of the patients was 3.4 mo (95%CI: 2.2-4.6 mo) (**Figure 1A** and **B**). One patient without liver metastases at baseline was found with a substantially prolonged PFS of 11.2 mo. Still, of the patients with liver metastases at baseline, no prolonged PFS was observed in 2 patients even after the combination therapy using local hepatectomy

Table 1 Characteristics of the individual patients

No.	Age (yr)	Sex	ECOG PS	Primary tumor	Site of metastases	Liver metastases	RAS and BRAF mutation	Lines of treatment	Prior 5-Fu	Prior platinum	Prior irinotecan	Prior bevacizumab	Prior cetuximab	Prior regorafenib	Prior PD-1
1	56	Female	1	Sigmoid colon	Abdominal metastases	No	Wild-type	3	Yes	Yes	Yes	Yes	Yes	No	No
2	62	Female	3	Ascending colon	Multiple metastases	Yes	Not tested	4	Yes	Yes	Yes	Yes	No	Yes	No
3	51	Female	0	Rectum	Multiple metastases	Yes	Not tested	3	Yes	Yes	Yes	Yes	No	No	No
4	52	Female	1	Sigmoid colon	Multiple metastases	No	KRAS mutation	3	Yes	Yes	Yes	Yes	No	No	No
5	66	Female	1	Ascending colon	Multiple metastases	Yes	Not tested	4	Yes	Yes	Yes	Yes	No	Yes	No
6	57	Female	2	Sigmoid colon	Multiple metastases	No	Wild-type	3	Yes	Yes	Yes	Yes	Yes	No	No
7	48	Female	3	Liver curvature of the transverse colon	Multiple metastases	Yes	Not tested	3	Yes	Yes	Yes	Yes	No	No	No
8	61	Female	2	Rectum	Pelvic metastases	No	KRAS mutation	3	Yes	Yes	Yes	Yes	No	No	No
9	65	Female	1	Rectum	Ovary metastases	No	Wild-type	4	Yes	Yes	Yes	Yes	Yes	Yes	No
10	65	Female	0	Ascending colon	Lung metastases	No	Not tested	3	Yes	Yes	Yes	Yes	No	No	No
11	56	Female	2	Left half of the colon	Ovary metastases	No	Wild-type	3	Yes	Yes	Yes	Yes	Yes	No	No
12	27	Male	0	Sigmoid colon	Pelvic metastases	No	Wild-type	4	Yes	Yes	Yes	Yes	Yes	Yes	Yes
13	58	Male	1	Sigmoid colon	Liver metastases	Yes	Not tested	3	Yes	Yes	Yes	Yes	No	No	No
14	70	Female	1	Rectum	Lung metastases	No	Not tested	4	Yes	Yes	Yes	Yes	No	No	No
15	54	Male	1	Ascending colon	Multiple metastases	Yes	KRAS mutation	3	Yes	Yes	Yes	Yes	No	No	No
16	55	Male	1	Rectum	Multiple metastases	Yes	NRAS mutation	4	Yes	Yes	Yes	Yes	No	Yes	No
17	48	Female	0	Rectum	Multiple metastases	Yes	Wild-type	3	Yes	Yes	Yes	Yes	Yes	No	No
18	55	Male	1	Rectum	Multiple	Yes	Wild-type	3	Yes	Yes	Yes	Yes	Yes	No	No

					metastases											
19	64	Male	2	Rectum	Multiple metastases	Yes	KRAS mutation	3	Yes	Yes	Yes	Yes	No	No	No	
20	58	Female	0	Ascending colon	Multiple metastases	Yes	Wild-type	3	Yes	Yes	Yes	Yes	Yes	No	No	
21	44	Female	2	Sigmoid colon	Multiple metastases	Yes	Wild-type	3	Yes	Yes	Yes	Yes	Yes	No	No	
22	51	Male	1	Rectum	Multiple metastases	Yes	Not tested	4	Yes	Yes	Yes	Yes	No	Yes	No	
23	56	Male	1	Rectum	Lung metastases	No	Not tested	4	Yes	Yes	Yes	Yes	No	No	No	
24	61	Male	2	Rectum	Multiple metastases	Yes	Not tested	4	Yes	Yes	Yes	Yes	No	No	No	
25	73	Male	0	Rectum	Multiple metastases	Yes (radiofrequency)	Wild-type	3	Yes	Yes	Yes	Yes	Yes	No	No	
26	61	Male	1	Rectum	Multiple metastases	Yes	Wild-type	4	Yes	Yes	Yes	Yes	Yes	Yes	No	
27	58	Male	2	Sigmoid colon	Multiple metastases	No	KRAS mutation	3	Yes	Yes	Yes	Yes	No	No	No	
28	57	Female	1	Sigmoid colon	Liver metastases	Yes (hepatectomy)	KRAS mutation	3	Yes	Yes	Yes	Yes	No	No	No	
29	67	Male	2	Rectum	Liver metastases	Yes	Not tested	3	Yes	Yes	Yes	Yes	No	No	No	
30	61	Male	1	Sigmoid colon	Peritoneal metastasis	No	Wide-type	3	Yes	Yes	Yes	Yes	Yes	No	No	

ECOG PS: Eastern Cooperative Oncology Group performance status.

and radiofrequency ablation. The median PFS was 2.8 mo (95%CI: 0.8-4.7 mo) for the 18 patients with liver metastases and 3.5 mo (95%CI: 3.0-4.0 mo) for the 12 patients without liver metastases, with no significant difference ($P = 0.17$) (Figure 1C). The median PFS of the 19 patients treated with previous regorafenib treatment was 2.8 mo (95%CI: 1.0-4.6 mo), while the median PFS of the 11 patients naïve to regorafenib was 3.4 mo (95%CI: 2.8-4.0 mo); the difference was not statistically significant ($P = 0.55$) (Figure 1D). The median PFS was not significantly different between the patients with or without RAS mutation ($P = 0.37$). The 4 patients with imported PD-1 inhibitor (nivolumab or pembrolizumab) and 26 patients with domestic PD-1 inhibitor (sintilimab, toripalimab, camrelizumab, or tislelizumab) did not differ in median PFS ($P = 0.31$). At the time of April 15, 2021, the data for OS were still not mature. Detailed PFS and OS data can be found in Table 2.

Table 2 Treatments and outcomes

No.	Initial dose of regorafenib (mg)	Dose adjustment (mg)	PD-1 inhibitor used in combination	Number of treatment cycles	Best response	PFS (mo)	OS (mo)
1	120		Camrelizumab	16	SD	11.2	12.0
2	80		Tislelizumab	7	SD	5.0	7.0
3	160	160→80	Nivolumab	4	PD	3.4	Not reached
4	80		Camrelizumab	4	PD	3.4	4.0
5	80		Tislelizumab	3	SD	2.0	5.0
6	80		Toripalimab	5	SD	3.5	Not reached
7	80		Tislelizumab	7	SD	5.0	9.0
8	80		Pembrolizumab	5	PD	3.5	Not reached
9	120		Sintilimab	4	SD	2.8	Not reached
10	160	160→80	Toripalimab	8	SD	5.6	Not reached
11	80		Tislelizumab	2	SD	1.4	Not reached
12	160		Toripalimab	4	SD	2.8	Not reached
13	120		Nivolumab	6	SD	4.2	Not reached
14	80		Toripalimab	5	SD	3.5	Not reached
15	120		Sintilimab	6	SD	3.5	Not reached
16	80		Nivolumab	15	SD	10.5	Not reached
17	160	160→80	Sintilimab	4	PD	2.8	Not reached
18	80		Camrelizumab	5	PD	3.5	Not reached
19	80		Toripalimab	2	SD	1.4	Not reached
20	120		Camrelizumab	4	PD	3.4	7.0
21	80		Toripalimab	2	PD	1.4	Not reached
22	120		Camrelizumab	1	SD	0.7	12.0
23	80		Sintilimab	2	PD	1.4	10.0
24	80		Toripalimab	9	SD	6.3	11.0
25	120	120→80	Sintilimab	2	PD	1.4	Not reached
26	80		Sintilimab	3	SD	2.1	10.0
27	80		Toripalimab	6	SD	4.2	Not reached
28	120		Sintilimab	2	PD	1.4	Not reached
29	80		Sintilimab	1	PD	0.7	not reached
30	80		Sintilimab	2	PD	1.4	Not

OS: Overall survival; PD: Progressive disease; PFS: Progression-free survival; SD: Stable disease; PD-1: Programmed cell death-1.




Figure 1 Kaplan-Meier survival curves. A: Progression-free survival (PFS) of 30 patients; B: Overall survival of the whole cohort; C: PFS according to the patients with or without liver metastasis ($P > 0.05$); D: PFS according to previous exposure to regorafenib or not ($P > 0.05$). PFS: Progression-free survival; OS: Overall survival.

Safety profile

Of the 30 patients, treatment-related adverse events (TRAEs) occurred in 17 patients (56.7%). The major TRAEs included hand-foot syndrome (33.3%), hypertension (23.3%), malaise (20.0%), and gastrointestinal reaction (16.7%). The other TRAEs included transaminase elevation (13.3%), diarrhea (10.0%), abnormal capillary proliferation (6.7%), thrombocytopenia (6.7%), hypothyroidism (6.7%), proteinuria (6.7%), rash (6.7%), anemia (3.3%), myocardial enzyme elevation (3.3%), and oral mucositis (3.3%). The incidence of grade 3 TRAEs was 13.3% (4/30), which comprised abnormal capillary proliferation ($n = 1$), transaminase elevation ($n = 1$), and hand-foot syndrome ($n = 2$). No grade 4 or higher toxicity was observed. For patients treated with different initial doses of regorafenib, more grade 3 TRAEs were observed in the 160 mg group. Specifically, 3 of the 4 patients treated with an initial dose of regorafenib of 160 mg had grade 3 TRAEs, including grade 3 hand-foot syndrome in 2 patients and grade 3 abnormal capillary proliferation in 1 patient. For all 3 patients, the dose of regorafenib was reduced to 80 mg. In the 8 patients treated with an initial dose of regorafenib of 120 mg, 1 patient had grade 3 transaminase elevation. For this patient, regorafenib and PD-1 inhibitor were discontinued until the transaminase levels returned to normal. Subsequently, the regorafenib dose was reduced to 80 mg to

treat the patient combined with a PD-1 inhibitor. All 18 patients in the 80 mg group had good tolerance; they could tolerate the therapy after symptomatic treatment until PD. Table 3 lists the details of the TRAEs.

Relationship between PLR and treatment response

The NLR and PLR of the 30 patients were calculated, and the ROC curves were plotted. The area under the ROC curve (AUC) of NLR was 0.533, which could not effectively predict the treatment efficacy (Figure 2A). The AUC of PLR was 0.774, the maximum Youden index was 0.549, the corresponding cutoff value of PLR was 118, and the sensitivity and specificity were 85.7% and 69.2%, respectively (Figure 2B).

The median PLR was 151.1 (96.0-211.8). The PLR values differed significantly between the SD and PD groups ($P = 0.047$) (Figure 3A). Subsequently, the patients were divided into the PLR < 118 group (low-PLR group, $n = 11$) and PLR ≥ 118 group (high-PLR group, $n = 19$) according to the cutoff value. The median PFS in the low-PLR group was 4.2 mo (95%CI: 3.5-4.9 mo), compared with 2.8 mo (95%CI: 1.4-4.2 mo) in the high-PLR group ($P = 0.005$) (Figure 3B). The PLR may be an indicator to predict patient response and selected those who with longer PFS to regorafenib combined with PD-1 inhibitor. For the 12 patients with PD, the CEA increased after one and two cycles of treatment (Figure 4A). For the 18 patients in the SD group, the CEA in 1 patient increased transiently with a low amplitude after the first treatment cycle (higher than the normal range). In contrast, after the second treatment cycle, the CEA in all the patients was stable or decreased compared with before treatment (Figure 4B). The PLR in 3 patients decreased slightly after the first treatment cycle, while the PLR in 11 patients increased after two treatment cycles (Figure 4C). Still, the PLR level in 5 patients increased slightly after the first treatment cycle (higher than the cutoff value). The PLR in 1 patient still increased after the second treatment cycle (Figure 4D). The sensitivity of PLR was slightly lower than that of CEA for predicting treatment response.

DISCUSSION


This study aimed to investigate the benefits of regorafenib combined with a PD-1 inhibitor in treating MSS mCRC and explore indicators predicting treatment response and prognosis. Treatment using regorafenib combined with PD-1 inhibitor could lead to a longer PFS in some patients with MSS mCRC with failure to standard treatment. Our study also analyzed and compared the PLR of the patients in different treatment cycles with the corresponding CEA levels and explored the possible predictive value of PLR for predicting the response to treatment.

Recently, many studies on combination therapy with immunotherapy were performed to improve the immune responses and clinical efficacies on malignant tumors[26,27]. Still, studies on immunotherapy combined with vascular endothelial growth factor (VEGF) inhibitors have not shown significant improvements in PFS or OS[28,29]. Combination therapy using regorafenib and PD-1 inhibitors has already shown synergistic effects in mouse models[15,30]. In addition, the REGONIVO trial in Japan reported that the response rate was 36%, and the median PFS was 7.9 mo in 25 patients with mCRC (including one with microsatellite instability-high mCRC) treated with regorafenib plus nivolumab [14]. On the other hand, a retrospective study in the United States did not replicate the findings of the REGONIVO study[15]. Specifically, the clinical responses of the patients were relatively poor; the rate of disease progression was as high as 69%, and the rate of SD was only 31%, without PR or CR[15]. A retrospective study in Shandong, China analyzed the data of 23 patients; SD was found in 18 patients, and the DCR was 78.3%; PD was found in 5 patients, and the PD rate was 21.7%[16]. In the present study, the data of 30 patients were retrospectively analyzed, making it the largest study to date. Of these patients, 18 had SD during the treatment, and the DCR was 60.0%, which was substantially higher than the 31% reported by the American study[15]. The median PFS was 3.4 mo in this study, which was not as outstanding as in the REGONIVO study (7.9 mo) but was longer than the 2.0 mo reported by the American study and comparable with the 3.1 mo reported by the Shandong study. The PFS in 2 patients was longer than 10 mo and was 11.2 mo for the patient with the longest PFS.

The differences between the findings of this study and those of the American study could be due to the following reasons. First, 77.8% of the patients included in the American study had baseline liver metastases. Second, only 4 Asian patients were included in the American study, and patients of different ethnicities could respond differently to the treatment. Liver metastasis was considered an important factor influencing the study results. As an immune-tolerant organ, the liver is related to high percentages of immune suppressor cells[31]. The immune tolerance of the liver is used by primary hepatocellular carcinoma and liver metastases to inhibit the anti-tumor immune responses and decrease the efficacy of treatments using immune checkpoint inhibitors[32]. In addition, several studies demonstrated that liver metastases could also exert systemic immunosuppression effects in patients with cancer, which consequently inhibited the intra- and extrahepatic immune responses[33,34]. A promising method to overcome the inherent immune escape of liver tumors is the combination therapy of liver cancer using anti-VEGF drugs and immune checkpoint inhibitors, as the anti-VEGF drugs could

Table 3 Treatment-related adverse event

Adverse event	Number of patients (n = 30), n (%)		
	Any grade	Grade 1-2	Grade 3
Any event	17 (56.7)	13 (43.3)	4 (13.3)
Hand-foot syndrome	10 (33.3)	8 (26.7)	2 (6.7)
Hypertension	7 (23.3)	7 (23.3)	0
Malaise	6 (20.0)	6 (20.0)	0
Gastrointestinal reaction	5 (16.7)	5 (16.7)	0
Transaminase elevation	4 (13.3)	3 (10.0)	1 (3.3)
Diarrhea	3 (10.0)	3 (10.0)	0
Abnormal capillary proliferation	2 (6.7)	1 (3.3)	1 (3.3)
Hypothyroidism	2 (6.7)	2 (6.7)	0
Proteinuria	2 (6.7)	2 (6.7)	0
Rash	2 (6.7)	2 (6.7)	0
Thrombocytopenia	2 (6.7)	2 (6.7)	0
Anemia	1 (3.3)	1 (3.3)	0
Myocardial enzyme elevation	1 (3.3)	1 (3.3)	0
Oral mucositis	1 (3.3)	1 (3.3)	0
Leukopenia/neutropenia	0	0	0



DOI: 10.4251/wjgo.v14.i4.920 Copyright ©The Author(s) 2022.

Figure 2 Receiver operating characteristics curves. A: The receiver operating characteristics curves of neutrophil-to-lymphocyte ratio; B: The receiver operating characteristics curves of platelet-to-lymphocyte ratio. NLR: Platelet -to-lymphocyte ratio; PLR: Neutrophil-to-lymphocyte ratio; ROC: Receiver operating characteristics.

reverse the VEGF-mediated immune suppression, promote T-cell infiltration of tumor microenvironment, and consequently enhance the treatment effects of ICIs[35]. The multicenter study in Shandong [25] and the REGONIVO study[14] included 56.5% and 52.0% of patients with liver metastases, respectively, while this frequency was 77.8% in the American study. The present study included 60% of patients with liver metastases. The relatively encouraging findings of the REGONIVO study could be related to the good ECOG PS of the patients, while the failure of the American study could be directly related to the high percentage of patients with liver metastases. In the present study, the ECOG PS and the percentage of liver metastases were more evenly distributed. Moreover, the number of patients with each domestic PD-1 inhibitor was limited in our study. Due to the national conditions and patients’




Figure 3 Relationship between platelet-to-lymphocyte ratio and treatment response. A: Difference in platelet-to-lymphocyte ratio according to stable disease and progressive disease; B: Progression-free survival according to the platelet-to-lymphocyte ratio (*P* < 0.05). PLR: Platelet-to-lymphocyte ratio; PFS: Progression-free survival; SD: Stable disease; PD: Progressive disease.

financial burden, the number of patients who received nivolumab (*n* = 3) or pembrolizumab (*n* = 1) was also very low. Thus, we analyzed the difference in PFS between patients with imported PD-1 inhibitor (nivolumab or pembrolizumab) and those with domestic anti-PD-1 inhibitor (sintilimab, toripalimab, camrelizumab, or tislelizumab), which showed no statistical significance. This comforting result was also supported by previous clinical trials of domestic PD-1 inhibitors. Thus, the study could provide more objective evidence for evaluating the clinical efficacy of this treatment regimen.

In the present study, the doses of regorafenib included 160, 120, and 80 mg. The dose of regorafenib in all 4 patients treated with 160 mg was reduced to 80 mg due to grade 3 TRAE, and the regorafenib dose in some patients treated with 120 mg was also reduced to 80 mg due to abnormal transaminase levels. Regorafenib treatment could be continued at this dose for all patients, suggesting that 80 mg could be used as the best dose in the combination therapy for further investigation. Besides nivolumab and pembrolizumab used worldwide, four other PD-1 inhibitors common in China were included in this study because many patients cannot afford nivolumab and pembrolizumab.

Since the cost of the combination treatment is relatively high, discovering predictive markers for treatment efficacy is important to identify the patients who could best benefit from the treatment. The findings showed that the specificity of PLR was slightly lower than that of CEA (the PLR of one patient with SD still increased slightly after two treatment cycles), while the sensitivity was comparable (the trend of changes in most patients conformed to the treatment efficacy after the first treatment cycle). The PFS was significantly different between the PLR-low and -high groups, indicating that the PLR was negatively correlated to PFS. The findings suggested that the PLR could be used as a reference to predict the treatment efficacy and PFS of patients when selecting the combination therapy. An Italian study demonstrated that PLR was an independent factor influencing the outcomes of CRC[36]. Moreover, patients with high PLR also showed a high expression level of programmed cell death-ligand 1 in circulating tumor cells, suggesting that PLR may also be a predictive marker of change in tumor immune microenvironment[36]. This may explain why PLR can predict the effectiveness of PD-1 inhibitor combination therapy but NLR cannot.

This study has limitations. Although we retrospectively included all the patients, only 30 were included in this study. The treatment responses of this treatment regimen need to be verified through prospective studies with large sample sizes. No multivariable Cox regression model could be established due to the relatively small sample size of this study, and the predict value of PLR should be further validated.

CONCLUSION

In conclusion, treatment using regorafenib combined with PD-1 inhibitor could lead to a longer PFS in some patients with MSS mCRC with failure to standard treatment. The PLR should be examined further for its ability to predict response to regorafenib combined with a PD-1 inhibitor. These results could help the design of a prospective trial in patients with refractory MSS mCRC.




Figure 4 Dynamic changes in carcinoembryonic antigen ($\mu\text{g/mL}$) and platelet-to-lymphocyte ratio after one and two treatment cycles. A: Carcinoembryonic antigen ($\mu\text{g/mL}$) in time in patients with stable disease; B: Carcinoembryonic antigen ($\mu\text{g/mL}$) in time in patients with progressive disease; C: Platelet-to-lymphocyte in time in patients with stable disease; D: Platelet-to-lymphocyte in time in patients with progressive disease. PLR: Platelet-to-lymphocyte ratio; SD: Stable disease; PD: Progressive disease.

ARTICLE HIGHLIGHTS

Research background

The effectiveness of the combination therapy using regorafenib and programmed cell death-1 (PD-1) inhibitors in treating metastatic colorectal cancer (mCRC) in the REGONIVO trial in Japan and a retrospective study in the United States are inconsistent.

Research motivation

As the effectiveness of the combination therapy remains controversial, we evaluated the situation and data of the combination therapy including the efficacy and safety in our medical centre in order to provide more clinical evidence for this treatment.

Research objectives

The objectives of this study were to investigate the tumor response, progression-free survival, overall survival, and treatment-related adverse events of the treatment and explore a potential indicators predicting response and prognosis.

Research methods

We identified patients with microsatellite stable (MSS) mCRC treated with regorafenib combined with PD-1 inhibitor at Henan Provincial People's Hospital between December 2018 and December 2020. Collected data included age, sex, Eastern Cooperative Oncology Group (ECOG) performance status (PS), site of the primary tumor, site of the metastases, MSI/MMR, gene status, lines of treatment, and previous treatments. The blood routine examination and CEA results before treatment and after three and five cycles of combination therapy were longitudinally analyzed.

Research results

We included 30 patients with MSS mCRC treated with regorafenib combined with PD-1 inhibitor. The disease control rate was 60.0%. The median follow-up time was 12.0 mo, and median PFS was 3.4 mo [95% confidence interval (CI): 2.2-4.6 mo]. The median PFS in the low-PLR group was 4.2 mo (95% CI: 3.5-4.9 mo), compared with 2.8 mo (95% CI: 1.4-4.2 mo) in the high-PLR group ($P = 0.005$). Four (13.3%) patients experienced grade 3 TRAE.

Research conclusions

We find that some patients can benefit from the combination therapy even after multi-line therapy and adverse events are generally tolerable. The PLR might be a potential indicator to predict patient response to this combination therapy.

Research perspectives

This study provides experiences and could help to design a prospective trial for patients with MSS mCRC those who failure to standard therapy.

FOOTNOTES

Author contributions: Liu MY and Xu YJ designed the research; Xu YJ, Zhang P, and Hu JL performed the research; Liang H, Zhu YY, and Cui Y contributed new reagents/analytic tools; Xu YJ, Zhang P, Niu P, and Xu M analyzed the data; Liu MY, Xu YJ, and Zhang P wrote the paper.

Supported by the Henan Provincial Department of Science and Technology, No. 212102310047.

Institutional review board statement: This study was approved by the ethics committee of People's Hospital of Zhengzhou University (Henan Province, China) and performed in accordance with the Declaration of Helsinki.

Informed consent statement: The requirement for informed consent was waived by the committee because of the retrospective nature of the study.

Conflict-of-interest statement: The authors declare no conflict of interest.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Yu-Jie Xu 0000-0003-1833-6541; Peng Zhang 0000-0002-1359-8035; Jin-Long Hu 0000-0002-0032-0202; Hong Liang 0000-0003-4686-9791; Yan-Yan Zhu 0000-0003-2395-0351; Yao Cui 0000-0003-0273-9178; Po Niu 0000-0001-9369-6980; Min Xu 0000-0002-2949-0133; Ming-Yue Liu 0000-0001-6090-0295.

S-Editor: Wang LL

L-Editor: Filipodia

P-Editor: Li X

REFERENCES

- 1 Ferlay J, Colombet M, Soerjomataram I, Mathers C, Parkin DM, Piñeros M, Znaor A, Bray F. Estimating the global cancer incidence and mortality in 2018: GLOBOCAN sources and methods. *Int J Cancer* 2019; **144**: 1941-1953 [PMID: 30350310 DOI: 10.1002/ijc.31937]

- 2 **Bray F**, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018; **68**: 394-424 [PMID: 30207593 DOI: 10.3322/caac.21492]
- 3 **Chen W**, Zheng R, Baade PD, Zhang S, Zeng H, Bray F, Jemal A, Yu XQ, He J. Cancer statistics in China, 2015. *CA Cancer J Clin* 2016; **66**: 115-132 [PMID: 26808342 DOI: 10.3322/caac.21338]
- 4 **Fu Q**, Zhang Q, Lou Y, Yang J, Nie G, Chen Q, Chen Y, Zhang J, Wang J, Wei T, Qin H, Dang X, Bai X, Liang T. Correction: Primary tumor-derived exosomes facilitate metastasis by regulating adhesion of circulating tumor cells via SMAD3 in liver cancer. *Oncogene* 2019; **38**: 5740-5741 [PMID: 31068668 DOI: 10.1038/s41388-019-0830-6]
- 5 **Van Cutsem E**, Cervantes A, Adam R, Sobrero A, Van Krieken JH, Aderka D, Aranda Aguilar E, Bardelli A, Benson A, Bodoky G, Ciardiello F, D'Hoore A, Diaz-Rubio E, Douillard JY, Ducreux M, Falcone A, Grothey A, Gruenberger T, Haustermans K, Heinemann V, Hoff P, Köhne CH, Labianca R, Laurent-Puig P, Ma B, Maughan T, Muro K, Normanno N, Österlund P, Oyen WJ, Papamichael D, Pentheroudakis G, Pfeiffer P, Price TJ, Punt C, Ricke J, Roth A, Salazar R, Scheithauer W, Schmoll HJ, Tabernero J, Taieb J, Tejpar S, Wasan H, Yoshino T, Zaanan A, Arnold D. ESMO consensus guidelines for the management of patients with metastatic colorectal cancer. *Ann Oncol* 2016; **27**: 1386-1422 [PMID: 27380959 DOI: 10.1093/annonc/mdw235]
- 6 **NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines)**. Colon Cancer. Version 2.2021. Fort Washington: National Comprehensive Cancer Network, 2021
- 7 **NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines)**. Rectal Cancer. Version 1.2021. Fort Washington: National Comprehensive Cancer Network, 2021
- 8 **Siegel RL**, Miller KD, Jemal A. Cancer statistics, 2019. *CA Cancer J Clin* 2019; **69**: 7-34 [PMID: 30620402 DOI: 10.3322/caac.21551]
- 9 **Diagnosis And Treatment Guidelines For Colorectal Cancer Working Group CSOCOC**. Chinese Society of Clinical Oncology (CSCO) diagnosis and treatment guidelines for colorectal cancer 2018 (English version). *Chin J Cancer Res* 2019; **31**: 117-134 [PMID: 30996570 DOI: 10.21147/j.issn.1000-9604.2019.01.07]
- 10 **Dhillon S**. Regorafenib: A Review in Metastatic Colorectal Cancer. *Drugs* 2018; **78**: 1133-1144 [PMID: 29943375 DOI: 10.1007/s40265-018-0938-y]
- 11 **Grothey A**, Blay JY, Pavlakis N, Yoshino T, Bruix J. Evolving role of regorafenib for the treatment of advanced cancers. *Cancer Treat Rev* 2020; **86**: 101993 [PMID: 32199197 DOI: 10.1016/j.ctrv.2020.101993]
- 12 **Yamaguchi K**, Komatsu Y, Satoh T, Uetake H, Yoshino T, Nishida T, Yamazaki N, Takikawa H, Morimoto T, Chosa M, Sunaya T, Hamada Y, Muro K, Sugihara K. Large-Scale, Prospective Observational Study of Regorafenib in Japanese Patients with Metastatic Colorectal Cancer in a Real-World Clinical Setting. *Oncologist* 2019; **24**: e450-e457 [PMID: 30606885 DOI: 10.1634/theoncologist.2018-0377]
- 13 **Le DT**, Uram JN, Wang H, Bartlett BR, Kemberling H, Eyring AD, Skora AD, Luber BS, Azad NS, Laheru D, Biedrzycki B, Donehower RC, Zaheer A, Fisher GA, Crocenzi TS, Lee JJ, Duffy SM, Goldberg RM, de la Chapelle A, Koshiji M, Bhaijee F, Hrubner T, Hruban RH, Wood LD, Cuka N, Pardoll DM, Papadopoulos N, Kinzler KW, Zhou S, Cornish TC, Taube JM, Anders RA, Eshleman JR, Vogelstein B, Diaz LA Jr. PD-1 Blockade in Tumors with Mismatch-Repair Deficiency. *N Engl J Med* 2015; **372**: 2509-2520 [PMID: 26028255 DOI: 10.1056/NEJMoa1500596]
- 14 **Fukuoka S**, Hara H, Takahashi N, Kojima T, Kawazoe A, Asayama M, Yoshii T, Kotani D, Tamura H, Mikamoto Y, Hirano N, Wakabayashi M, Nomura S, Sato A, Kuwata T, Togashi Y, Nishikawa H, Shitara K. Regorafenib Plus Nivolumab in Patients With Advanced Gastric or Colorectal Cancer: An Open-Label, Dose-Escalation, and Dose-Expansion Phase Ib Trial (REGONIVO, EPOC1603). *J Clin Oncol* 2020; **38**: 2053-2061 [PMID: 32343640 DOI: 10.1200/JCO.19.03296]
- 15 **Wang C**, Chevalier D, Saluja J, Sandhu J, Lau C, Fakhri M. Regorafenib and Nivolumab or Pembrolizumab Combination and Circulating Tumor DNA Response Assessment in Refractory Microsatellite Stable Colorectal Cancer. *Oncologist* 2020; **25**: e1188-e1194 [PMID: 32406541 DOI: 10.1634/theoncologist.2020-0161]
- 16 **Li J**, Cong L, Liu J, Peng L, Wang J, Feng A, Yue J, Li L, Wang X. The Efficacy and Safety of Regorafenib in Combination With Anti-PD-1 Antibody in Refractory Microsatellite Stable Metastatic Colorectal Cancer: A Retrospective Study. *Front Oncol* 2020; **10**: 594125 [PMID: 33282742 DOI: 10.3389/fonc.2020.594125]
- 17 **Cho WK**, Choi DH, Park HC, Park W, Yu JI, Park YS, Park JO, Lim HY, Kang WK, Kim HC, Cho YB, Yun SH, Lee WY. Elevated CEA is associated with worse survival in recurrent rectal cancer. *Oncotarget* 2017; **8**: 105936-105941 [PMID: 29285304 DOI: 10.18632/oncotarget.22511]
- 18 **Auclin E**, Taieb J, Lepage C, Aparicio T, Faroux R, Mini E, Folprecht G, Salazar R, Benetkiewicz M, Banzi M, Louvet C, Van Laethem JL, Tabernero J, Hickish T, de Gramont A, André T, Vernerey D. Carcinoembryonic Antigen Levels and Survival in Stage III Colon Cancer: *Post hoc* Analysis of the MOSAIC and PETACC-8 Trials. *Cancer Epidemiol Biomarkers Prev* 2019; **28**: 1153-1161 [PMID: 31263053 DOI: 10.1158/1055-9965.EPI-18-0867]
- 19 **Tanoglu A**, Karagoz E. Neutrophil and platelet-to-lymphocyte ratio: new predictors of dropout and recurrence after liver transplantation for hepatocellular cancer? *Transpl Int* 2014; **27**: e80-e81 [PMID: 24612189 DOI: 10.1111/tri.12291]
- 20 **Huang S**, Li D, Zhuang L, Sun L, Wu J. A meta-analysis of the efficacy and safety of adjuvant sorafenib for hepatocellular carcinoma after resection. *World J Surg Oncol* 2021; **19**: 168 [PMID: 34112190 DOI: 10.1186/s12957-021-02280-9]
- 21 **Miao Y**, Yan Q, Li S, Li B, Feng Y. Neutrophil to lymphocyte ratio and platelet to lymphocyte ratio are predictive of chemotherapeutic response and prognosis in epithelial ovarian cancer patients treated with platinum-based chemotherapy. *Cancer Biomark* 2016; **17**: 33-40 [PMID: 27314290 DOI: 10.3233/CBM-160614]
- 22 **Zhang Y**, Lu JJ, Du YP, Feng CX, Wang LQ, Chen MB. Prognostic value of neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio in gastric cancer. *Medicine (Baltimore)* 2018; **97**: e0144 [PMID: 29561419 DOI: 10.1097/MD.00000000000010144]
- 23 **Kim JY**, Jung EJ, Kim JM, Lee HS, Kwag SJ, Park JH, Park T, Jeong SH, Jeong CY, Ju YT. Dynamic changes of neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio predicts breast cancer prognosis. *BMC Cancer* 2020; **20**: 1206 [PMID: 33287745 DOI: 10.1186/s12885-020-07700-9]
- 24 **Messersmith WA**. NCCN Guidelines Updates: Management of Metastatic Colorectal Cancer. *J Natl Compr Canc Netw*

- 2019; **17**: 599-601 [PMID: [31117039](#) DOI: [10.6004/jnccn.2019.5014](#)]
- 25 **Zhang L**, Wang J, Zhang B, Chu Q, Su C, Wu H, Chen X, Wang B, Yin Y, Zhu B, Sun J. Attitudes and Practices of Immune Checkpoint Inhibitors in Chinese Patients With Cancer: A National Cross-Sectional Survey. *Front Pharmacol* 2021; **12**: 583126 [PMID: [33841138](#) DOI: [10.3389/fphar.2021.583126](#)]
- 26 **Golshani G**, Zhang Y. Advances in immunotherapy for colorectal cancer: a review. *Therap Adv Gastroenterol* 2020; **13**: 1756284820917527 [PMID: [32536977](#) DOI: [10.1177/1756284820917527](#)]
- 27 **Johdi NA**, Sukor NF. Colorectal Cancer Immunotherapy: Options and Strategies. *Front Immunol* 2020; **11**: 1624 [PMID: [33042104](#) DOI: [10.3389/fimmu.2020.01624](#)]
- 28 **Mettu NB**, Twohy E, Ou FS, Halfdanarson TR, Lenz HJ, Breakstone R, Boland PM, Crysler O, Wu C, Grothey A, Nixon AB, Bolch E, Niedzwiecki D, Fruth B, Schweitzer B, Elsing A, Hurwitz H, Fakih MG, Bekaii-Saab T. BACCI: A phase II randomized, double-blind, multicenter, placebo-controlled study of capecitabine (C) bevacizumab (B) plus atezolizumab (A) or placebo (P) in refractory metastatic colorectal cancer (mCRC): An ACCRU network study. *Ann Oncol* 2019; **30** [DOI: [10.1093/annonc/mdz246.011](#)]
- 29 **Grothey A**, Taberero J, Arnold D, De Gramont A, Ducreux MP, O'Dwyer PJ, Van Cutsem E, Bosanac I, Srock S, Mancao C, Gilberg F, Winter J, Schmoll HJ. Fluoropyrimidine (FP) + bevacizumab (BEV) + atezolizumab vs FP/BEV in BRAFwt metastatic colorectal cancer (mCRC): Findings from Cohort 2 of MODUL – a multicentre, randomized trial of biomarker-driven maintenance treatment following first-line induction therapy. *Ann Oncol* 2018; **29**: 714-715 [DOI: [10.1093/annonc/mdy424.020](#)]
- 30 **Shigeta K**, Matsui A, Kikuchi H, Klein S, Mamessier E, Chen IX, Aoki S, Kitahara S, Inoue K, Shigeta A, Hato T, Ramjiawan RR, Staiculescu D, Zopf D, Fiebig L, Hobbs GS, Quaas A, Dima S, Popescu I, Huang P, Munn LL, Cobbold M, Goyal L, Zhu AX, Jain RK, Duda DG. Regorafenib combined with PD1 blockade increases CD8 T-cell infiltration by inducing CXCL10 expression in hepatocellular carcinoma. *J Immunother Cancer* 2020; **8** [PMID: [33234602](#) DOI: [10.1136/jitc-2020-001435](#)]
- 31 **Zheng M**, Tian Z. Liver-Mediated Adaptive Immune Tolerance. *Front Immunol* 2019; **10**: 2525 [PMID: [31787967](#) DOI: [10.3389/fimmu.2019.02525](#)]
- 32 **Yu J**, Green MD, Li S, Sun Y, Journey SN, Choi JE, Rizvi SM, Qin A, Waninger JJ, Lang X, Chopra Z, El Naqa I, Zhou J, Bian Y, Jiang L, Tezel A, Skvarce J, Achar RK, Sitto M, Rosen BS, Su F, Narayanan SP, Cao X, Wei S, Szeliga W, Vatan L, Mayo C, Morgan MA, Schonewolf CA, Cuneo K, Kryczek I, Ma VT, Lao CD, Lawrence TS, Ramnath N, Wen F, Chinnaiyan AM, Cieslik M, Alva A, Zou W. Liver metastasis restrains immunotherapy efficacy via macrophage-mediated T cell elimination. *Nat Med* 2021; **27**: 152-164 [PMID: [33398162](#) DOI: [10.1038/s41591-020-1131-x](#)]
- 33 **Lee JC**, Mehdizadeh S, Smith J, Young A, Mufazalov IA, Mowery CT, Daud A, Bluestone JA. Regulatory T cell control of systemic immunity and immunotherapy response in liver metastasis. *Sci Immunol* 2020; **5** [PMID: [33008914](#) DOI: [10.1126/sciimmunol.aba0759](#)]
- 34 **Tumeh PC**, Hellmann MD, Hamid O, Tsai KK, Loo KL, Gubens MA, Rosenblum M, Harview CL, Taube JM, Handley N, Khurana N, Nosrati A, Krummel MF, Tucker A, Sosa EV, Sanchez PJ, Banayan N, Osorio JC, Nguyen-Kim DL, Chang J, Shintaku IP, Boasberg PD, Taylor EJ, Munster PN, Algazi AP, Chmielowski B, Dummer R, Grogan TR, Elashoff D, Hwang J, Goldinger SM, Garon EB, Pierce RH, Daud A. Liver Metastasis and Treatment Outcome with Anti-PD-1 Monoclonal Antibody in Patients with Melanoma and NSCLC. *Cancer Immunol Res* 2017; **5**: 417-424 [PMID: [28411193](#) DOI: [10.1158/2326-6066.CIR-16-0325](#)]
- 35 **Voron T**, Colussi O, Marcheteau E, Pernot S, Nizard M, Pointet AL, Latreche S, Bergaya S, Benhamouda N, Tanchot C, Stockmann C, Combe P, Berger A, Zinzindohoue F, Yagita H, Tartour E, Taieb J, Terme M. VEGF-A modulates expression of inhibitory checkpoints on CD8+ T cells in tumors. *J Exp Med* 2015; **212**: 139-148 [PMID: [25601652](#) DOI: [10.1084/jem.20140559](#)]
- 36 **Raimondi L**, Raimondi FM, Di Benedetto L, Cimino G, Spinelli GP. PD-L1 Expression on Circulating Tumour Cells May Be Predictive of Response to Regorafenib in Patients Diagnosed with Chemorefractory Metastatic Colorectal Cancer. *Int J Mol Sci* 2020; **21** [PMID: [32962309](#) DOI: [10.3390/ijms21186907](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

