

World Journal of *Cardiology*

World J Cardiol 2022 May 26; 14(5): 271-328



MINIREVIEWS

- 271 Same day discharge after structural heart disease interventions in the era of the coronavirus-19 pandemic and beyond

Asbeutah AA, Junaid M, Hassan F, Avila Vega J, Efeovbokhan N, Khouzam RN, Ibebuogu UN

ORIGINAL ARTICLE**Basic Study**

- 282 Bioinformatics prediction of potential mechanisms and biomarkers underlying dilated cardiomyopathy

Liu Z, Song YN, Chen KY, Gao WL, Chen HJ, Liang GY

Retrospective Study

- 297 Pledget-assisted hemostasis to fix residual access-site bleedings after double pre-closure technique

Burzotta F, Aurigemma C, Kovacevic M, Romagnoli E, Cangemi S, Bianchini F, Nesta M, Bruno P, Trani C

- 307 Day-to-day blood pressure variability predicts poor outcomes following percutaneous coronary intervention: A retrospective study

Weisel CL, Dyke CM, Klug MG, Haldis TA, Basson MD

META-ANALYSIS

- 319 Comparative efficacy and safety of adenosine and regadenoson for assessment of fractional flow reserve: A systematic review and meta-analysis

Gill GS, Gadre A, Kanmanthareddy A

ABOUT COVER

Editorial Board Member of *World Journal of Cardiology*, Yu-Li Huang, MD, PhD, Chief Doctor, Professor, Research Scientist, Clinical Research Centre, Shunde Hospital, Southern Medical University, Foshan 528300, Guangdong Province, China. hyuli821@smu.edu.cn

AIMS AND SCOPE

The primary aim of *World Journal of Cardiology* (*WJC*, *World J Cardiol*) is to provide scholars and readers from various fields of cardiology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJC mainly publishes articles reporting research results and findings obtained in the field of cardiology and covering a wide range of topics including acute coronary syndromes, aneurysm, angina, arrhythmias, atherosclerosis, atrial fibrillation, cardiomyopathy, congenital heart disease, coronary artery disease, heart failure, hypertension, imaging, infection, myocardial infarction, pathology, peripheral vessels, public health, Raynaud's syndrome, stroke, thrombosis, and valvular disease.

INDEXING/ABSTRACTING

The *WJC* is now abstracted and indexed in Emerging Sources Citation Index (Web of Science), PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2021 edition of Journal Citation Reports® cites the 2020 Journal Citation Indicator (JCI) for *WJC* as 0.36. The *WJC*'s CiteScore for 2020 is 0.3, and Scopus CiteScore rank 2020: Cardiology and Cardiovascular Medicine is 289/317.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Hua-Ge Yin; Production Department Director: Xiang Li; Editorial Office Director: Ya-Juan Ma.

NAME OF JOURNAL

World Journal of Cardiology

ISSN

ISSN 1949-8462 (online)

LAUNCH DATE

December 31, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Ramdas G Pai, Dimitrios Tousoulis, Marco Matteo Ciccone, Pal Pacher

EDITORIAL BOARD MEMBERS

<https://www.wjnet.com/1949-8462/editorialboard.htm>

PUBLICATION DATE

May 26, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Basic Study

Bioinformatics prediction of potential mechanisms and biomarkers underlying dilated cardiomyopathy

Zhou Liu, Ying-Nan Song, Kai-Yuan Chen, Wei-Long Gao, Hong-Jin Chen, Gui-You Liang

Specialty type: Engineering, biomedical

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Emran TB, Bangladesh; Tanabe S, Japan

Received: December 14, 2021

Peer-review started: December 14, 2021

First decision: January 25, 2022

Revised: February 19, 2022

Accepted: April 26, 2022

Article in press: April 26, 2022

Published online: May 26, 2022



Zhou Liu, Gui-You Liang, School of Basic Medical Sciences, Guizhou Medical University, Guiyang 550025, Guizhou Province, China

Zhou Liu, Ying-Nan Song, Kai-Yuan Chen, Wei-Long Gao, Hong-Jin Chen, Gui-You Liang, Translational Medicine Research Center, Guizhou Medical University, Guiyang 550025, Guizhou Province, China

Ying-Nan Song, Hong-Jin Chen, Gui-You Liang, Department of Cardiovascular Surgery, the Affiliated Hospital of Guizhou Medical University, Guiyang 510000, Guizhou Province, China

Corresponding author: Gui-You Liang, MD, Professor, School of Basic Medical Sciences, Guizhou Medical University, Dangwu, Guian District, Guiyang 550025, Guizhou Province, China. guiyou515@163.com

Abstract

BACKGROUND

Heart failure is a health burden responsible for high morbidity and mortality worldwide, and dilated cardiomyopathy (DCM) is one of the most common causes of heart failure. DCM is a disease of the heart muscle and is characterized by enlargement and dilation of at least one ventricle alongside impaired contractility with left ventricular ejection fraction < 40%. It is also associated with abnormalities in cytoskeletal proteins, mitochondrial ATP transporter, microvasculature, and fibrosis. However, the pathogenesis and potential biomarkers of DCM remain to be investigated.

AIM

To investigate the candidate genes and pathways involved in DCM patients.

METHODS

Two expression datasets (GSE3585 and GSE5406) were downloaded from the Gene Expression Omnibus database. The differentially expressed genes (DEGs) between the DCM patients and healthy individuals were identified using the R package "linear models for microarray data." The pathways with common DEGs were analyzed *via* Gene Ontology (GO), Kyoto Encyclopedia of Genes and Genomes (KEGG), and gene set enrichment analyses. Moreover, a protein-protein interaction network (PPI) was constructed to identify the hub genes and modules. The MicroRNA Database was applied to predict the microRNAs (miRNAs) targeting the hub genes. Additionally, immune cell infiltration in DCM was

analyzed using CIBERSORT.

RESULTS

In total, 97 DEGs (47 upregulated and 50 downregulated) were identified. GO analysis showed that the DEGs were mainly enriched in “response to growth factor,” “extracellular matrix,” and “extracellular matrix structural constituent.” KEGG pathway analysis indicated that the DEGs were mainly enriched in “protein digestion and absorption” and “interleukin 17 (IL-17) signaling pathway.” The PPI network suggested that collagen type III alpha 1 chain (COL3A1) and COL1A2 contribute to the pathogenesis of DCM. Additionally, visualization of the interactions between miRNAs and the hub genes revealed that hsa-miR-5682 and hsa-miR-4500 interacted with both COL3A1 and COL1A2, and thus these miRNAs might play roles in DCM. Immune cell infiltration analysis revealed that DCM patients had more infiltrated plasma cells and fewer infiltrated B memory cells, T follicular helper cells, and resting dendritic cells.

CONCLUSION

COL1A2 and COL3A1 and their targeting miRNAs, hsa-miR-5682 and hsa-miR-4500, may play critical roles in the pathogenesis of DCM, which are closely related to the IL-17 signaling pathway and acute inflammatory response. These results may provide useful clues for the diagnosis and treatment of DCM.

Key Words: Dilated cardiomyopathy; Bioinformatics; Differentially expressed genes; Function enrichment analysis; Protein-protein interaction network; Immune cell infiltration

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: As the most common cause of heart failure, the diagnosis and therapy for dilated cardiomyopathy (DCM) are still unsatisfactory because of its indistinct pathogenesis and specific biomarkers. Thus, we comprehensively utilized the microarray data from the Gene Expression Omnibus database to uncover the biomarker and mechanisms underlying DCM development. Collagen type III alpha 1 chain and collagen type I alpha 2 chain, which are regulated by hsa-miR-5682 and hsa-miR-4500, may play critical roles in the pathogenesis of DCM through the acute inflammatory response and interleukin 17 signaling pathway. These biomarkers and mechanisms need to be further studied.

Citation: Liu Z, Song YN, Chen KY, Gao WL, Chen HJ, Liang GY. Bioinformatics prediction of potential mechanisms and biomarkers underlying dilated cardiomyopathy. *World J Cardiol* 2022; 14(5): 282-296

URL: <https://www.wjgnet.com/1949-8462/full/v14/i5/282.htm>

DOI: <https://dx.doi.org/10.4330/wjc.v14.i5.282>

INTRODUCTION

Dilated cardiomyopathy (DCM) is a progressive myocardial disease. It accounts for 30%–40% of heart failure cases and leads to high mortality worldwide[1]. DCM is characterized by biventricular dilatation, cardiac systolic dysfunction, and ventricular remodeling[2]. Recently, several studies have reported that mutations, myocarditis, hypertension, and ischemia are the induction factors of DCM[3,4]. Increasing evidence shows that various gene mutations and biomarkers are associated with DCM[5-7]. Mutations in cytoskeletal proteins, including dystrophin[8] and desmin[9], impair muscular force transmission and thereby contribute to the development of DCM. Mutations in lamin A/C, a nuclear membrane protein, usually cause DCM with atrioventricular block and atrial fibrillation[10]. Li *et al*[11] reported that mutation of aryl hydrocarbon receptor nuclear translocator-like protein 1 (known as BMAL1) plays a critical role in the development of DCM through the regulation of mitochondrial fission and mitophagy via mitochondrial protein B cell leukemia/lymphoma 2 interacting protein 3. Mutations in thin filament regulatory proteins including cardiac troponin T, cardiac troponin I, and α -tropomyosin can cause DCM with systolic dysfunction by reducing fractional shortening and systolic calcium level[12]. Moreover, some biomarkers associated with the development of DCM have been reported. For example, syndecan-1 and syndecan-4 may serve as useful biomarkers for predicting adverse cardiovascular events in DCM patients[13,14]. Carbonic anhydrase 2 and 3 are associated with heart failure and are potential risk biomarkers for DCM[15], and serum fibroblast growth factor 21 level is linked to the prognosis of DCM[16].

Several studies have sought DCM-related genes and mechanisms *via* bioinformatic methods and found some meaningful results. Huang *et al*[17] found that Fos proto-oncogene, AP-1 transcription factor subunit, tissue inhibitor of metalloprotease-1, and serpin family E member 1 may serve as therapeutic targets in DCM. Zhao *et al*[18] identified 89 differentially expressed genes (DEGs), mainly enriched in the extracellular matrix and biological adhesion signaling pathways, which may play significant roles in the development of DCM. However, the main cause(s) and pathogenic mechanism(s) underlying DCM are still unknown; thus, DCM is mostly diagnosed late, which causes a poor prognosis in turn. More studies are urgently needed to improve the diagnostic and therapeutic efficiency in DCM. The Gene Expression Omnibus (GEO) database includes many DCM-related microarray data, which have not been fully utilized. These data can be used to identify additional candidate biomarkers and pathways to further explore the cause(s) of DCM. To investigate the candidate genes and pathways involved in DCM patients, we analyzed the two gene expression data sets GSE3585 and GSE5406. Using the “linear models for microarray data” (limma) method, we identified 97 DEGs between healthy individuals and DCM patients. In addition, we identified the mechanisms commonly regulated by the DEGs *via* Gene Ontology (GO), Kyoto Encyclopedia of Genes and Genomes (KEGG) analyses, and gene set enrichment analyses (GSEA). Moreover, a protein-protein interaction network (PPI) network was applied to identify the hub genes that may contribute to the pathogenesis of DCM and predict the microRNAs (miRNAs) targeting the hub genes. Furthermore, we investigated the pattern of immune cell infiltration in DCM.

MATERIALS AND METHODS

Microarray data extraction from the GEO database

The mRNA expression profiles GSE3585 and GSE5406 in the GEO database (<http://www.ncbi.nlm.nih.gov/geo/>), which is a shared platform in which researchers deposit their microarray data related to various diseases, were downloaded. The GSE3585 dataset, generated by Barth *et al*[19], and the GSE5406 dataset, generated by Hannenhalli *et al*[20], consisted of 7 DCM patients and 5 healthy individuals, and 86 DCM patients and 16 healthy individuals, respectively. In total, 114 samples of the left ventricular myocardium, consisting of 93 DCM and 21 healthy samples (control group), were included in this study.

Data processing and DEGs identification

The two datasets GSE3585 and GSE5406 were loaded onto the GPL96 platform (Affymetrix Human Genome U133A Array [HG-U133A]). Additionally, the series matrix and platform annotation for the two databases were downloaded from the GEO database. The probe identity documents were transformed into gene symbols. Then, *via* the R package “Surrogate Variable Analysis,” the two databases were merged, and any batch effect was removed using the “Empirical Bayes” method[21]. The R package “limma” was applied to identify the DEGs between the DCM and healthy myocardium tissues[22]. The screening criteria were set as $P < 0.05$ and $|\log \text{ fold change (FC)}| > 0.589$ ($\text{FC} > 1.5$). Volcano and heat maps were generated using R software.

Gene expression enrichment analysis

The gene expression enrichment analysis in this study included GO analysis (<https://www.geneontology.org/>)[23], KEGG (<https://www.genome.jp/kegg/>)[24] pathway analysis, and GSEA (<https://www.gsea-msigdb.org/gsea/>) analysis[25]. The DEGs were inputted into Metascape (<https://metascape.org/>)[26]: The species was selected as *Homo sapiens*; The screening standard was set as $P < 0.05$; and The GO terms of biological process (BP), cellular component (CC), and molecular function (MF) were analyzed and KEGG pathway analysis was performed with the criteria of $P < 0.05$. GSEA interprets the biological function of the expression dataset. The expression dataset in the DCM cases *vs* healthy tissues was loaded into GSEA 4.0.3 software, set gene sets database as GO gene set (c5.all.v7.1.symbols.gmt), set number of permutations as 1000, set phenotype labels as control *vs* DCM, set collapse/remap to gene symbols as no collapse, set permutation type as phenotype, and the other parameters were set at default parameters. Then the GSEA software was used to obtain the enrichment results. The enriched terms were defined as significant with nominal $P < 0.05$.

PPI network construction and hub gene identification

The PPI network was constructed with the online website Search Tool for the Retrieval of Interacting Genes/Proteins (STRING, <https://stringdb.org/cgi/input.pl>)[27] to contribute to the understanding of the interactive relationship among DEGs. The DEGs were inputted into this website, species of *Homo sapiens* was selected, and the identification criterion was set as combined score > 0.4 (medium confidence). Then the profile of interacting node pairs was imported into Cytoscape (<https://cytoscape.org/>)[28] to visualize the PPI network. The top 10 hub genes were identified with the standard of connectivity degree by using the CytoHubba plugin. The plugin Molecular Complex




Figure 1 Identification of differentially expressed genes. A: Volcano map, in which the red node represents the upregulated differentially expressed genes (DEGs) and the blue node represents the downregulated DEGs; B: Heatmap of DEGs.

Detection (MCODE) was applied to identify the essential module within the PPI network in Cytoscape with the default parameters (degree cutoff, 2; node score cutoff, 0.2; kcore, 2; and maximum depth, 100).

Construction of the miRNA-mRNA interaction network

miRNAs, a class of small non-coding RNAs, regulate the expression of various genes by binding to their transcripts and play critical roles in DCM progression[29]. By using the MicroRNA Database (miRDB) [30] (<http://mirdb.org/>), we predicted miRNAs targeting any of the top 10 hub genes. Then, we sorted these miRNAs according to their prediction scores and selected the top 10 miRNAs. The mRNA-miRNA pairs were imported into Cytoscape to visualize the miRNA-mRNA network.

Immune cell infiltration analysis

The CIBERSORT (cibersort.stanford.edu) algorithm was applied to analyze the normalized gene expression data, and the proportions of 22 types of immune cells in each sample were analyzed[31]. The gene expression data were normalized *via* “limma” and transformed into the 22 types of immune cell expression data through the source of CIBERSORT[32] *via* R. Then the results were filtered out *via* Perl (<https://www.perl.org>) with $P < 0.05$, and the immune cell infiltration matrix was obtained. Next, the “vioplot” package was used to draw violin diagrams to visualize the difference in immune cell infiltration between the DCM and healthy groups in detail. The “ggplot2”[33] package was applied to perform principal component analysis (PCA) and draw a PCA clustering map. The “corrplot” package was used to analyze the correlation among immune cell infiltration and draw a correlation heatmap.


RESULTS

Identification of DEGs

After merging the two datasets, 97 DEGs, including 47 upregulated and 50 downregulated genes, were obtained in the DCM group compared with the control group. Figure 1 shows the volcano map and heatmap of the 97 DEGs. The details of the top 10 upregulated or downregulated genes are shown in Table 1.

GO and KEGG enrichment analyses

Next, the DEGs were used to perform enrichment analysis for BP, CC, MF, and KEGG pathways. By using the Metascape website, the BP of GO was found to be significantly enriched in “response to growth factor,” “blood vessel development,” “regulation of smooth muscle cell proliferation,” “muscle tissue development,” and “acute inflammatory response” (Figure 2A). The DEGs in CC were mainly enriched in “extracellular matrix,” “cytoplasmic vesicle lumen,” “collagen trimer,” and “sarcoplasm” (Figure 2B). Regarding MF, the DEGs were mainly enriched in “extracellular matrix structural constituent,” “growth factor binding,” “alpha-actinin binding,” and “calcium ion binding” (Figure 2C). KEGG pathway analysis revealed significant pathway enrichment of DEGs in “protein digestion and absorption,” “interleukin 17 (IL-17) signaling pathway,” “advanced glycation end products-receptor for advanced glycation end products signaling pathway in diabetic complications,” “complement,” and “coagulation cascades” (Figure 2D).



DOI: 10.4330/wjc.v14.i5.282 Copyright ©The Author(s) 2022.

Figure 2 Gene Ontology and Kyoto Encyclopedia of Genes and Genomes analysis of enrichment for differentially expressed genes. A: Biological process; B: Cellular component; C: Molecular function; D: Kyoto Encyclopedia of Genes and Genomes.

GSEA analysis

GSEA analysis results revealed that, compared with the control group, the DCM group was significantly enriched in GO terms “heart development,” “response to ischemia,” “vascular smooth muscle cell differentiation,” “response to transforming growth factor beta,” “stem cell proliferation,” and “regulation of mitochondrial fission” (Figure 3).

PPI network and identification of hub genes

To further explore the relationship among the DEGs at the protein level, the PPI network of the 97 DEGs was constructed using STRING with the criterion of combined score > 0.4 and visualized using Cytoscape. The PPI network consisted of 77 nodes and 145 edges (Figure 4A). The top 10 hub genes included collagen type III alpha 1 chain (COL3A1), COL1A2, signal transducer and activator of transcription 3 (STAT3), C-C motif chemokine ligand 2 (CCL2), fibromodulin (FMOD), asporin (ASPN), C-X-C motif chemokine ligand 12 (CXCL12), lumican (LUM), heat shock protein 90 alpha family class A member 1 (HSP90AA1), and osteoglycin (OGN) (Figure 4B). The detailed information of these hub genes is provided in Table 2. MCODE analysis identified five essential modules, and COL3A1, myosin heavy chain 6, activating transcription factor 3, B cell leukemia/lymphoma 6 transcription repressor,




Figure 3 Gene set enrichment analyses of enrichment of differentially expressed genes. A: Heart development; B: Response to ischemia; C: Vascular smooth muscle cell differentiation; D: Response to transforming growth factor beta; E: Stem cell proliferation; F: Regulation of mitochondrial fission.




Figure 4 Protein-protein interaction network. A: Protein-protein interaction (PPI) network of differentially expressed genes (DEGs), in which the red node represents upregulated DEGs and the green node represents the downregulated DEGs; B: Top 10 hub genes; C: Module 1; D: Module 2; E: Module 3; F: Module 4; G: Module 5. Yellow node represents the Seed gene of each module.

and pentraxin 3 were the seeds of clusters 1, 2, 3, 4, and 5, respectively (Figure 4C–G).

miRNA–mRNA interaction network


Increasing evidence shows that miRNAs play important roles in the development and progression of DCM. By using the miRDB database, we predicted miRNAs targeting any of the top 10 hub genes. Then we sorted these miRNAs according to their prediction scores and selected the top 10 miRNAs. Additionally, the top 100 miRNA–mRNA pairs were visualized using Cytoscape (Figure 5). Consequently, hsa-miR-5682, hsa-miR-4500, hsa-miR-32-3p, and hsa-miR-374a-3p were each found to target ≥ 2 hub genes.

Immune cell infiltration in DCM

Immune cells infiltrate into the myocardium upon myocardial injury[28]. Thus, a violin plot was constructed to investigate the difference in immune-cell infiltration between the DCM and control groups (Figure 6A). Compared with the control group, the DCM group had more infiltrated plasma cells and fewer infiltrated B memory cells, T follicular helper cells, and resting dendritic cells (DCs), whereas there was no significant difference in the remaining 18 types of immune cells. However, the PCA results showed that the control and DCM groups could not be well distinguished according to the infiltration patterns of the 22 types of immune cells (Figure 6B). We generated a correlation heatmap to assess the correlation among the 19 immune cells that were found infiltrated in the DCM or control group. As shown in Figure 6C, the number of infiltrated B memory cells was positively correlated with that of the infiltrated resting DCs, activated natural killer (NK) cells and T follicular helper cells, the number of infiltrated activated NK cells was negatively correlated with that of infiltrated resting NK cells, and the number of infiltrated resting memory CD4 T cells was negatively associated with that of infiltrated B memory cells and regulatory T cells.

DISCUSSION

DCM is one of the main reasons of sudden cardiac death and heart failure. It is a heterogeneous disease caused by various types of pathogenic factors including genetic, infectious, hormonal, and environmental factors[34]. The causes of DCM should be explored in depth to improve the diagnosis, treatment, and prognosis of DCM patients. Therefore, it is of great significance to elucidate the genetic mechanisms involved in DCM.



DOI: 10.4330/wjc.v14.i5.282 Copyright ©The Author(s) 2022.

Figure 6 Immune cell infiltration analyses of differentially expressed genes. A: Violin plot, in which red represents dilated cardiomyopathy group and blue represents the normal group; B: Principal component analysis; C: Correlation heatmap.

Table 1 Top 10 differentially expressed genes in the dilated cardiomyopathy group vs normal group

Expression	Gene symbol	Description	logFC	adj. P value
UP	<i>ASPN</i>	Asporin	2.31	3.05E-22
	<i>NPPA</i>	Natriuretic peptide A	2.08	1.26E-11
	<i>LUM</i>	Lumican	1.87	4.56E-18
	<i>MXRA5</i>	Matrix remodeling associated 5	1.43	5.47E-14
	<i>HBB</i>	Hemoglobin subunit beta	1.20	3.08E-05
	<i>COL1A2</i>	Collagen type I alpha 2 chain	1.09	1.37E-10
	<i>COL3A1</i>	Collagen type III alpha 1 chain	1.05	2.39E-07
	<i>OGN</i>	Osteoglycin	1.01	5.38E-09
	<i>FRZB</i>	Frizzled related protein	0.99	1.08E-13
	<i>EIF1AY</i>	Eukaryotic translation initiation factor 1A Y-linked	0.96	1.33E-02
	<i>MYOT</i>	Myotilin	-1.89	1.70E-13
	<i>ANKRD2</i>	Ankyrin repeat domain 2	-1.44	3.00E-08
	<i>PLA2G2A</i>	Phospholipase A2 group IIA	-1.35	4.43E-07
	<i>MYH6</i>	Myosin heavy chain 6	-1.35	1.80E-05
	<i>SERPINA3</i>	Serpin family A member 3	-1.28	1.75E-18
Down	<i>CYP4B1</i>	Cytochrome P450 family 4 subfamily B member 1	-1.20	1.06E-07
	<i>FCN3</i>	Ficolin 3	-1.19	2.48E-13
	<i>CNN1</i>	Calponin 1	-1.16	2.09E-10
	<i>PTX3</i>	Pentraxin 3	-1.13	1.27E-04
	<i>NRAP</i>	Nebulin related anchoring protein	-1.11	1.30E-03

Table 2 Top 10 hub genes in the protein-protein interaction network

Gene symbol	Description	Rank	Degree
<i>COL3A1</i>	Collagen type III alpha 1 chain	1	18
<i>COL1A2</i>	Collagen type I alpha 2 chain	2	12
<i>STAT3</i>	Signal transducer and activator of transcription 3	3	11
<i>CCL2</i>	C-C motif chemokine ligand 2	3	11
<i>FMOD</i>	Fibromodulin	5	10
<i>ASPN</i>	Asporin	6	9
<i>CXCL12</i>	C-X-C motif chemokine ligand 12	6	9
<i>LUM</i>	Lumican	8	8
<i>HSP90AA1</i>	Heat shock protein 90 alpha family class A member 1	8	8
<i>OGN</i>	Osteoglycin	8	8

KEGG pathway analysis showed that the DEGs were significantly enriched in the IL-17 signaling pathway. DCM is induced by viral myocarditis, accompanying autoimmune dysfunction, affecting the secretion of IL-17 cytokine by T helper 17 cells, and IL-17 itself promotes myocardial cell injury[51]. Wang *et al*[52] reported that elevated IL-17 levels are significantly associated with DCM incidence and progression. Additionally, the serum levels of other inflammatory factors such as IL-6, tumor necrosis factor- α , and IL-21 are significantly increased in DCM patients[53]. Thus, the IL-17 signaling pathway may be one of the major signaling pathways involved in the development of DCM.

Through construction of PPI and miRNA-mRNA interaction networks, we identified the hub genes and the miRNAs targeting them. The hub genes, COL1A2 and COL3A1, encode the pro- α 2/1 chains of type I and III collagens, respectively. Collagens I and III, the main collagens of cardiac extracellular

matrix, are classical biomarkers of cardiac fibrosis in DCM[54]. Mihailovici *et al*[55] reported that collagens I and III are upregulated in DCM patients compared with matched healthy controls. Additionally, Zhao *et al*[18] reported that COL1A2 may participate in DCM pathogenesis by regulating the cardiac remodeling characterized by collagen deposition in the extracellular matrix[56]. Consistent with our results, Liu *et al*[53] identified STAT3 as a hub gene in DCM *via* bioinformatic analysis. Other studies have also indicated a role of STAT3 in DCM. Podewski *et al*[57] showed that STAT3 protein level is significantly decreased in the cardiomyocytes of patients with end-stage DCM. Moreover, inhibition of the IL-6-mediated STAT3 signaling pathway can improve myocardial remodeling through reducing myocardial apoptosis in a mouse model of DCM[58]. Thus, the roles of COL1A2, COL3A1, and STAT3 in DCM should be further investigated. Moreover, the identified miRNAs hsa-miR-5682, hsa-miR-4500, hsa-miR-32-3p, and hsa-miR-374a-3p may participate in DCM pathogenesis through their interaction with $\geq$ two hub genes. Previous studies have also suggested that miRNAs play significant roles in DCM. It has been found that miR-21, miR-29a, and miR-133b are differentially expressed in DCM patients[59]. miR-133a expression is associated with fibrosis, myocyte necrosis, left ventricular function, and clinical outcome in patients with inflammatory DCM[60]. Moreover, Satoh *et al*[61] showed that a low let-7i level can serve as an independent predictor of cardiac death and heart failure (relative risk = 3.76).

Immune cells commonly infiltrate into the myocardium upon various types of cardiac damage[49, 62]. Overactivation of immune cells could be investigated in pathological examination about cardiac biopsy specimens in DCM patients. Noutsias *et al*[63] reported that the upregulation of genes associated with T cells exacerbates DCM progression. Therefore, our study also assessed the correlation between DEGs and immune cell infiltration. The results indicated infiltration of 19 types of immune cells in DCM pathogenesis. Notably, compared with the control group, the DCM group had more infiltrated plasma cells and fewer infiltrated B memory cells, T follicular helper cells, and resting DCs. However, Liu *et al* [53] have demonstrated that, compared with healthy controls, DCM patients have more infiltrated T follicular helper cells and fewer T follicular regulatory cells, and infiltration of T follicular regulatory cells is positively correlated with left ventricular ejection fraction.

Our study had some limitations. First, the gene expression data were acquired from a public database. Moreover, we did not experimentally verify the relevance of the identified DEGs with DCM and their enriched functions or hub genes. Likewise, we did not verify the predicted miRNA-mRNA interactions and their relevance.

CONCLUSION

In summary, we first identified that COL1A2 and COL3A1 may be both presumably regulated by hsa-miR-5682 and hsa-miR-4500, and play significant roles in the pathogenesis of DCM through acute inflammatory response and IL-17 signaling pathway. These results may provide useful biomarkers for the diagnosis and treatment of DCM, but further research is needed to clarify the roles of the predicted genes and pathways.

ARTICLE HIGHLIGHTS

Research background

Dilated cardiomyopathy (DCM), a disease of the heart muscle, is one of the most common causes of heart failure. However, the original cause and pathogenesis in development of DCM are still remain elusive.

Research motivation

The early diagnosis and prognosis of DCM patients are unsatisfactory because of DCM main cause and pathogenesis are still unclear. Increasing DCM datasets were provided online but little was been explored. Bioinformatics could further investigate the DCM mechanism and biomarkers for improving the diagnostic and therapeutic efficiency.

Research objectives

This study investigated the candidate genes and pathways involved in DCM patients.

Research methods

Expression datasets were downloaded from the Gene Expression Omnibus database. Gene Ontology, Kyoto Encyclopedia of Genes and Genomes, and gene set enrichment analyses investigated the key pathway in differentially expressed genes (DEGs) between the DCM patients and healthy individuals. Protein-protein interaction network identified the hub genes and modules in DCM. MicroRNA Database predicted the microRNAs which targeting the hub genes. CIBERSORT analyzed the immune-

ell infiltration in DCM.

Research results

Ninety-seven DEGs mainly enriched in “response to growth factor,” “extracellular matrix,” and “extracellular matrix structural constituent.” Moreover, the top two pathways were “protein digestion and absorption” and “interleukin 17 signaling pathway.” Collagen type III alpha 1 chain (COL3A1) and COL1A2, whose were regulated by hsa-miR-5682 and hsa-miR-4500, mainly contributed to the pathogenesis of DCM. Compared with the control group, DCM patients had more infiltrated plasma cells and fewer infiltrated B memory cells, T follicular helper cells, and resting dendritic cells.

Research conclusions

DCM progression closely related to IL-17 signaling pathway and acute inflammatory response. COL1A2 and COL3A1 and their targeting miRNAs, hsa-miR-5682 and hsa-miR-4500, are the potential biomarkers of DCM.

Research perspectives

This study may provide valuable pathways and biomarkers for the diagnosis or treatment of DCM. Further studies should investigate the functions of the predicted genes and pathways.

FOOTNOTES

Author contributions: Liu Z and Song YN designed this study; Chen KY collected the relevant data; Liu Z analyzed the data; Liu Z and Gao WL drafted the manuscript; Chen HJ and Liang GY reviewed and supervised this manuscript; All authors approved the final version of the article.

Supported by National Nature Science Foundation of China, No. 81960051, No. 8217021743, and No. 82160060; Project of High-Level Innovative Talents of Guizhou Province, No. [2016]4034; and Construction Funding from Characteristic Key Laboratory of Guizhou Province, No. [2021]313.

Conflict-of-interest statement: The authors have no conflicts of interest to declare.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Zhou Liu 0000-0002-6773-1073; Ying-Nan Song 0000-0003-4464-0512; Kai-Yuan Chen 0000-0002-0375-1091; Wei-Long Gao 0000-0002-0464-970X; Hong-Jin Chen 0000-0002-0513-9584; Gui-You Liang 0000-0002-4555-9102.

S-Editor: Ma YJ

L-Editor: Filipodia

P-Editor: Ma YJ

REFERENCES

- 1 **Papadakis M**, Sharma S, Cox S, Sheppard MN, Panoulas VF, Behr ER. The magnitude of sudden cardiac death in the young: a death certificate-based review in England and Wales. *Europace* 2009; **11**: 1353-1358 [PMID: 19700472 DOI: 10.1093/europace/eup229]
- 2 **Haas J**, Frese KS, Peil B, Kloos W, Keller A, Nietsch R, Feng Z, Müller S, Kayvanpour E, Vogel B, Sedaghat-Hamedani F, Lim WK, Zhao X, Fradkin D, Köhler D, Fischer S, Franke J, Marquart S, Barb I, Li DT, Amr A, Ehlermann P, Mereles D, Weis T, Hassel S, Kremer A, King V, Wirsz E, Isnard R, Komajda M, Serio A, Grasso M, Syrris P, Wicks E, Plagnol V, Lopes L, Gadgaard T, Eiskjær H, Jørgensen M, Garcia-Giustiniani D, Ortiz-Genga M, Crespo-Leiro MG, Deprez RH, Christiaans I, van Rijsingen IA, Wilde AA, Waldenström A, Bolognesi M, Bellazzi R, Mörmér S, Bermejo JL, Monserrat L, Villard E, Mogensen J, Pinto YM, Charron P, Elliott P, Arbustini E, Katus HA, Meder B. Atlas of the clinical genetics of human dilated cardiomyopathy. *Eur Heart J* 2015; **36**: 1123-135a [PMID: 25163546 DOI: 10.1093/eurheartj/ehu301]
- 3 **Bozkurt B**, Colvin M, Cook J, Cooper LT, Deswal A, Fonarow GC, Francis GS, Lenihan D, Lewis EF, McNamara DM, Pahl E, Vasan RS, Ramasubbu K, Rasmussen K, Towbin JA, Yancy C; American Heart Association Committee on Heart Failure and Transplantation of the Council on Clinical Cardiology; Council on Cardiovascular Disease in the Young; Council on Cardiovascular and Stroke Nursing; Council on Epidemiology and Prevention; and Council on Quality of Care

- and Outcomes Research. Current Diagnostic and Treatment Strategies for Specific Dilated Cardiomyopathies: A Scientific Statement From the American Heart Association. *Circulation* 2016; **134**: e579-e646 [PMID: [27832612](#) DOI: [10.1161/CIR.0000000000000455](#)]
- 4 **Vikhorev PG**, Vikhoreva NN. Cardiomyopathies and Related Changes in Contractility of Human Heart Muscle. *Int J Mol Sci* 2018; **19**: 2234 [PMID: [30065175](#) DOI: [10.3390/ijms19082234](#)]
 - 5 **Weintraub RG**, Semsarian C, Macdonald P. Dilated cardiomyopathy. *Lancet* 2017; **390**: 400-414 [PMID: [28190577](#) DOI: [10.1016/S0140-6736\(16\)31713-5](#)]
 - 6 **Willott RH**, Gomes AV, Chang AN, Parvatiyar MS, Pinto JR, Potter JD. Mutations in Troponin that cause HCM, DCM AND RCM: what can we learn about thin filament function? *J Mol Cell Cardiol* 2010; **48**: 882-892 [PMID: [19914256](#) DOI: [10.1016/j.yjmcc.2009.10.031](#)]
 - 7 **Pyun JH**, Kim HJ, Jeong MH, Ahn BY, Vuong TA, Lee DI, Choi S, Koo SH, Cho H, Kang JS. Cardiac specific PRMT1 ablation causes heart failure through CaMKII dysregulation. *Nat Commun* 2018; **9**: 5107 [PMID: [30504773](#) DOI: [10.1038/s41467-018-07606-y](#)]
 - 8 **Restrepo-Cordoba MA**, Wahbi K, Florian AR, Jiménez-Jáimez J, Politano L, Arad M, Climent-Paya V, Garcia-Alvarez A, Hansen RB, Larrañaga-Moreira JM, Kubanek M, Lopes LR, Ros A, Jurcut R, Rasmussen TB, Ruiz-Guerrero L, Pribe-Wolferts R, Palomino-Doza J, Bilinska Z, Rodríguez-Palomares JF, Van Loon RLE, Basurte Elorz MT, Quarta G, Robledo Iñarritu M, Verdonschot JAJ, Stojkovic T, Shomanova Z, Bermudez-Jimenez F, Palladino A, Freimark D, García-Álvarez MI, Jorda P, Dominguez F, Ochoa JP, Girolami F, Brugada R, Meder B, Barriales-Villa R, Mogensen J, Laforêt P, Yilmaz A, Elliott P, Garcia-Pavia P; European Genetic Cardiomyopathies Initiative Investigators (see online supplementary Appendix S1). Prevalence and clinical outcomes of dystrophin-associated dilated cardiomyopathy without severe skeletal myopathy. *Eur J Heart Fail* 2021; **23**: 1276-1286 [PMID: [34050592](#) DOI: [10.1002/ejhf.2250](#)]
 - 9 **Pawlak A**, Rejmak-Kozicka E, Gil KE, Ziemia A, Kaczmarek L, Gil RJ. Patterns of desmin expression in idiopathic dilated cardiomyopathy are related to the desmin mRNA and ubiquitin expression. *J Investig Med* 2019; **67**: 11-19 [PMID: [30097466](#) DOI: [10.1136/jim-2017-000707](#)]
 - 10 **Parks SB**, Kushner JD, Nauman D, Burgess D, Ludwigsen S, Peterson A, Li D, Jakobs P, Litt M, Porter CB, Rahko PS, Hershsberger RE. Lamin A/C mutation analysis in a cohort of 324 unrelated patients with idiopathic or familial dilated cardiomyopathy. *Am Heart J* 2008; **156**: 161-169 [PMID: [18585512](#) DOI: [10.1016/j.ahj.2008.01.026](#)]
 - 11 **Li E**, Li X, Huang J, Xu C, Liang Q, Ren K, Bai A, Lu C, Qian R, Sun N. BMAL1 regulates mitochondrial fission and mitophagy through mitochondrial protein BNIP3 and is critical in the development of dilated cardiomyopathy. *Protein Cell* 2020; **11**: 661-679 [PMID: [32277346](#) DOI: [10.1007/s13238-020-00713-x](#)]
 - 12 **Robinson P**, Sparrow AJ, Patel S, Malinowska M, Reilly SN, Zhang YH, Casadei B, Watkins H, Redwood C. Dilated cardiomyopathy mutations in thin-filament regulatory proteins reduce contractility, suppress systolic Ca²⁺, and activate NFAT and Akt signaling. *Am J Physiol Heart Circ Physiol* 2020; **319**: H306-H319 [PMID: [32618513](#) DOI: [10.1152/ajpheart.00272.2020](#)]
 - 13 **Liu W**, Wang Y, Zheng J, Song D, Zheng S, Ren L, Yao Y, Liu Y, Bai R, Dong J, Liu T. Syndecan-1 as an independent risk factor for the incidence of adverse cardiovascular events in patients having stage C and D heart failure with non-ischemic dilated cardiomyopathy. *Clin Chim Acta* 2019; **490**: 63-68 [PMID: [30578753](#) DOI: [10.1016/j.cca.2018.12.022](#)]
 - 14 **Bielecka-Dabrowa A**, Sakowicz A, Misztal M, von Haehling S, Ahmed A, Pietrucha T, Rysz J, Banach M. Differences in biochemical and genetic biomarkers in patients with heart failure of various etiologies. *Int J Cardiol* 2016; **221**: 1073-1080 [PMID: [27448535](#) DOI: [10.1016/j.ijcard.2016.07.150](#)]
 - 15 **Su H**, Hu K, Liu Z, Chen K, Xu J. Carbonic anhydrase 2 and 3 as risk biomarkers for dilated cardiomyopathy associated heart failure. *Ann Palliat Med* 2021; **10**: 12554-12565 [PMID: [35016406](#) DOI: [10.21037/apm-21-3561](#)]
 - 16 **Gu L**, Jiang W, Zheng R, Yao Y, Ma G. Fibroblast Growth Factor 21 Correlates with the Prognosis of Dilated Cardiomyopathy. *Cardiology* 2021; **146**: 27-33 [PMID: [33264784](#) DOI: [10.1159/000509239](#)]
 - 17 **Huang H**, Luo B, Wang B, Wu Q, Liang Y, He Y. Identification of Potential Gene Interactions in Heart Failure Caused by Idiopathic Dilated Cardiomyopathy. *Med Sci Monit* 2018; **24**: 7697-7709 [PMID: [30368515](#) DOI: [10.12659/MSM.912984](#)]
 - 18 **Zhao J**, Lv T, Quan J, Zhao W, Song J, Li Z, Lei H, Huang W, Ran L. Identification of target genes in cardiomyopathy with fibrosis and cardiac remodeling. *J Biomed Sci* 2018; **25**: 63 [PMID: [30115125](#) DOI: [10.1186/s12929-018-0459-8](#)]
 - 19 **Barth AS**, Kuner R, Buness A, Ruschhaupt M, Merk S, Zwermann L, Käab S, Kreuzer E, Steinbeck G, Mansmann U, Poustka A, Nabauer M, Stültmann H. Identification of a common gene expression signature in dilated cardiomyopathy across independent microarray studies. *J Am Coll Cardiol* 2006; **48**: 1610-1617 [PMID: [17045896](#) DOI: [10.1016/j.jacc.2006.07.026](#)]
 - 20 **Hannenhalli S**, Putt ME, Gilmore JM, Wang J, Parmacek MS, Epstein JA, Morrissey EE, Margulies KB, Cappola TP. Transcriptional genomics associates FOX transcription factors with human heart failure. *Circulation* 2006; **114**: 1269-1276 [PMID: [16952980](#) DOI: [10.1161/CIRCULATIONAHA.106.632430](#)]
 - 21 **Leek JT**, Johnson WE, Parker HS, Jaffe AE, Storey JD. The sva package for removing batch effects and other unwanted variation in high-throughput experiments. *Bioinformatics* 2012; **28**: 882-883 [PMID: [22257669](#) DOI: [10.1093/bioinformatics/bts034](#)]
 - 22 **Ritchie ME**, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* 2015; **43**: e47 [PMID: [25605792](#) DOI: [10.1093/nar/gkv007](#)]
 - 23 **Gene Ontology Consortium**. The Gene Ontology resource: enriching a GOLD mine. *Nucleic Acids Res* 2021; **49**: D325-D334 [PMID: [33290552](#) DOI: [10.1093/nar/gkaa1113](#)]
 - 24 **Kanehisa M**, Sato Y, Kawashima M. KEGG mapping tools for uncovering hidden features in biological data. *Protein Sci* 2022; **31**: 47-53 [PMID: [34423492](#) DOI: [10.1002/pro.4172](#)]
 - 25 **Subramanian A**, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* 2005; **102**: 15545-15550 [PMID: [16199517](#) DOI: [10.1073/pnas.0506580102](#)]
 - 26 **Zhou Y**, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, Benner C, Chanda SK. Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. *Nat Commun* 2019; **10**: 1523 [PMID: [30944313](#) DOI: [10.1038/s41467-019-0923-5](#)]

- 10.1038/s41467-019-09234-6]
- 27 **Szklarczyk D**, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, Simonovic M, Doncheva NT, Morris JH, Bork P, Jensen LJ, Mering CV. STRING v11: protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res* 2019; **47**: D607-D613 [PMID: [30476243](#) DOI: [10.1093/nar/gky1131](#)]
 - 28 **Shannon P**, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res* 2003; **13**: 2498-2504 [PMID: [14597658](#) DOI: [10.1101/gr.1239303](#)]
 - 29 **Isserlin R**, Merico D, Wang D, Vuckovic D, Bousette N, Gramolini AO, Bader GD, Emili A. Systems analysis reveals down-regulation of a network of pro-survival miRNAs drives the apoptotic response in dilated cardiomyopathy. *Mol Biosyst* 2015; **11**: 239-251 [PMID: [25361207](#) DOI: [10.1039/c4mb00265b](#)]
 - 30 **Chen Y**, Wang X. miRDB: an online database for prediction of functional microRNA targets. *Nucleic Acids Res* 2020; **48**: D127-D131 [PMID: [31504780](#) DOI: [10.1093/nar/gkz757](#)]
 - 31 **Chen B**, Khodadoust MS, Liu CL, Newman AM, Alizadeh AA. Profiling Tumor Infiltrating Immune Cells with CIBERSORT. *Methods Mol Biol* 2018; **1711**: 243-259 [PMID: [29344893](#) DOI: [10.1007/978-1-4939-7493-1_12](#)]
 - 32 **Newman AM**, Liu CL, Green MR, Gentles AJ, Feng W, Xu Y, Hoang CD, Diehn M, Alizadeh AA. Robust enumeration of cell subsets from tissue expression profiles. *Nat Methods* 2015; **12**: 453-457 [PMID: [25822800](#) DOI: [10.1038/nmeth.3337](#)]
 - 33 **Wickham H**. ggplot2. New York: Springer, Cham; 2016
 - 34 **Merlo M**, Cannatà A, Gobbo M, Stolfo D, Elliott PM, Sinagra G. Evolving concepts in dilated cardiomyopathy. *Eur J Heart Fail* 2018; **20**: 228-239 [PMID: [29271570](#) DOI: [10.1002/ehf.1103](#)]
 - 35 **Richardson P**, McKenna W, Bristow M, Maisch B, Mautner B, O'Connell J, Olsen E, Thiene G, Goodwin J, Gyarsas I, Martin I, Nordet P. Report of the 1995 World Health Organization/International Society and Federation of Cardiology Task Force on the Definition and Classification of cardiomyopathies. *Circulation* 1996; **93**: 841-842 [PMID: [8598070](#) DOI: [10.1161/01.cir.93.5.841](#)]
 - 36 **Maisch B**, Pankuweit S. Inflammatory dilated cardiomyopathy : Etiology and clinical management. *Herz* 2020; **45**: 221-229 [PMID: [32123933](#) DOI: [10.1007/s00059-020-04900-8](#)]
 - 37 **Barcena ML**, Pozdniakova S, Haritonow N, Breiter P, Kühl AA, Milting H, Baczkowski I, Ladilov Y, Regitz-Zagrosek V. Dilated cardiomyopathy impairs mitochondrial biogenesis and promotes inflammation in an age- and sex-dependent manner. *Aging (Albany NY)* 2020; **12**: 24117-24133 [PMID: [33303703](#) DOI: [10.18632/aging.202283](#)]
 - 38 **Brayson D**, Frustaci A, Verardo R, Chimenti C, Russo MA, Hayward R, Ahmad S, Vizcay-Barrena G, Protti A, Zammit PS, dos Remedios CG, Ehler E, Shah AM, Shanahan CM. Prelamin A mediates myocardial inflammation in dilated and HIV-associated cardiomyopathies. *JCI Insight* 2019; **4**: e126315 [PMID: [31622279](#) DOI: [10.1172/jci.insight.126315](#)]
 - 39 **Lynch TL 4th**, Ismahil MA, Jegga AG, Zilliox MJ, Troidl C, Prabhu SD, Sadayappan S. Cardiac inflammation in genetic dilated cardiomyopathy caused by MYBPC3 mutation. *J Mol Cell Cardiol* 2017; **102**: 83-93 [PMID: [27955979](#) DOI: [10.1016/j.yjmcc.2016.12.002](#)]
 - 40 **Xia Y**, Chen Z, Chen A, Fu M, Dong Z, Hu K, Yang X, Zou Y, Sun A, Qian J, Ge J. LCZ696 improves cardiac function via alleviating Drp1-mediated mitochondrial dysfunction in mice with doxorubicin-induced dilated cardiomyopathy. *J Mol Cell Cardiol* 2017; **108**: 138-148 [PMID: [28623750](#) DOI: [10.1016/j.yjmcc.2017.06.003](#)]
 - 41 **Liu GS**, Morales A, Vafiadaki E, Lam CK, Cai WF, Haghighi K, Adly G, Hershberger RE, Kranias EG. A novel human R25C-phospholamban mutation is associated with super-inhibition of calcium cycling and ventricular arrhythmia. *Cardiovasc Res* 2015; **107**: 164-174 [PMID: [25852082](#) DOI: [10.1093/cvr/cvv127](#)]
 - 42 **Knezevic T**, Myers VD, Gordon J, Tilley DG, Sharp TE 3rd, Wang J, Khalili K, Cheung JY, Feldman AM. BAG3: a new player in the heart failure paradigm. *Heart Fail Rev* 2015; **20**: 423-434 [PMID: [25925243](#) DOI: [10.1007/s10741-015-9487-6](#)]
 - 43 **Sutanto H**, Lyon A, Lumens J, Schotten U, Dobrev D, Heijman J. Cardiomyocyte calcium handling in health and disease: Insights from in vitro and in silico studies. *Prog Biophys Mol Biol* 2020; **157**: 54-75 [PMID: [32188566](#) DOI: [10.1016/j.biombio.2020.02.008](#)]
 - 44 **Sheng JJ**, Feng HZ, Pinto JR, Wei H, Jin JP. Increases of desmin and α -actinin in mouse cardiac myofibrils as a response to diastolic dysfunction. *J Mol Cell Cardiol* 2016; **99**: 218-229 [PMID: [26529187](#) DOI: [10.1016/j.yjmcc.2015.10.035](#)]
 - 45 **Sun N**, Yazawa M, Liu J, Han L, Sanchez-Freire V, Abilez OJ, Navarrete EG, Hu S, Wang L, Lee A, Pavlovic A, Lin S, Chen R, Hajjar RJ, Snyder MP, Dolmetsch RE, Butte MJ, Ashley EA, Longaker MT, Robbins RC, Wu JC. Patient-specific induced pluripotent stem cells as a model for familial dilated cardiomyopathy. *Sci Transl Med* 2012; **4**: 130ra47 [PMID: [22517884](#) DOI: [10.1126/scitranslmed.3003552](#)]
 - 46 **Li W**, Lu M, Banerjee S, Zhong J, Ye A, Molter J, Yu X. Ex vivo diffusion tensor MRI reflects microscopic structural remodeling associated with aging and disease progression in normal and cardiomyopathic Syrian hamsters. *NMR Biomed* 2009; **22**: 819-825 [PMID: [19434665](#) DOI: [10.1002/nbm.1394](#)]
 - 47 **Cetinkaya A**, Berge B, Sen-Hild B, Troidl K, Gajawada P, Kubin N, Valeske K, Schranz D, Akintürk H, Schönborg M, Kubin T, Choi YH, Richter M. Radixin Relocalization and Nonmuscle α -Actinin Expression Are Features of Remodeling Cardiomyocytes in Adult Patients with Dilated Cardiomyopathy. *Dis Markers* 2020; **2020**: 9356738 [PMID: [32774516](#) DOI: [10.1155/2020/9356738](#)]
 - 48 **Park KC**, Gaze DC, Collinson PO, Marber MS. Cardiac troponins: from myocardial infarction to chronic disease. *Cardiovasc Res* 2017; **113**: 1708-1718 [PMID: [29016754](#) DOI: [10.1093/cvr/cvx183](#)]
 - 49 **Schultheiss HP**, Fairweather D, Caforio ALP, Escher F, Hershberger RE, Lipshultz SE, Liu PP, Matsumori A, Mazzanti A, McMurray J, Priori SG. Dilated cardiomyopathy. *Nat Rev Dis Primers* 2019; **5**: 32 [PMID: [31073128](#) DOI: [10.1038/s41572-019-0084-1](#)]
 - 50 **Chang AN**, Parvatiyar MS, Potter JD. Troponin and cardiomyopathy. *Biochem Biophys Res Commun* 2008; **369**: 74-81 [PMID: [18157941](#) DOI: [10.1016/j.bbrc.2007.12.081](#)]
 - 51 **Baldeviano GC**, Barin JG, Talor MV, Srinivasan S, Bedja D, Zheng D, Gabrielson K, Iwakura Y, Rose NR, Cihakova D. Interleukin-17A is dispensable for myocarditis but essential for the progression to dilated cardiomyopathy. *Circ Res* 2010;

- 106: 1646-1655 [PMID: [20378858](#) DOI: [10.1161/CIRCRESAHA.109.213157](#)]
- 52 **Wang ZH**, Liao YH, Yuan J, Jin XJ, Yu M, Chen RZ, Xu DJ, Wei J, Wan J, Zhao DC, Han HY, Li B, Tian G, Hu G, Xu J. Continued Elevation of Plasma IL-4 and IL-17 Predicts the Progression from VMC to DCM. *Dis Markers* 2020; **2020**: 9385472 [PMID: [31998421](#) DOI: [10.1155/2020/9385472](#)]
- 53 **Liu X**, Zhang W, Han Z. Decreased circulating follicular regulatory T cells in patients with dilated cardiomyopathy. *Braz J Med Biol Res* 2021; **54**: e11232 [PMID: [34669781](#) DOI: [10.1590/1414-431X2021e11232](#)]
- 54 **Nagao K**, Inada T, Tamura A, Kajitani K, Shimamura K, Yukawa H, Aida K, Sowa N, Nishiga M, Horie T, Makita T, Ono K, Tanaka M. Circulating markers of collagen types I, III, and IV in patients with dilated cardiomyopathy: relationships with myocardial collagen expression. *ESC Heart Fail* 2018; **5**: 1044-1051 [PMID: [30273997](#) DOI: [10.1002/ehf2.12360](#)]
- 55 **Mihailovici AR**, Deliu RC, Mărgăritescu C, Simionescu CE, Donoiu I, Istrătoae O, Tudorașcu DR, Târtea EA, Gheonea DI. Collagen I and III, MMP-1 and TIMP-1 immunoexpression in dilated cardiomyopathy. *Rom J Morphol Embryol* 2017; **58**: 777-781 [PMID: [29250654](#)]
- 56 **Tsoutsman T**, Wang X, Garchow K, Riser B, Twigg S, Semsarian C. CCN2 plays a key role in extracellular matrix gene expression in severe hypertrophic cardiomyopathy and heart failure. *J Mol Cell Cardiol* 2013; **62**: 164-178 [PMID: [23756156](#) DOI: [10.1016/j.yjmcc.2013.05.019](#)]
- 57 **Podewski EK**, Hilfiker-Kleiner D, Hilfiker A, Morawietz H, Lichtenberg A, Wollert KC, Drexler H. Alterations in Janus kinase (JAK)-signal transducers and activators of transcription (STAT) signaling in patients with end-stage dilated cardiomyopathy. *Circulation* 2003; **107**: 798-802 [PMID: [12591746](#) DOI: [10.1161/01.cir.0000057545.82749.ff](#)]
- 58 **Li Q**, Ye WX, Huang ZJ, Zhang Q, He YF. Effect of IL-6-mediated STAT3 signaling pathway on myocardial apoptosis in mice with dilated cardiomyopathy. *Eur Rev Med Pharmacol Sci* 2019; **23**: 3042-3050 [PMID: [31002169](#) DOI: [10.26355/eurev_201904_17586](#)]
- 59 **Wang Y**, Li M, Xu L, Liu J, Wang D, Li Q, Wang L, Li P, Chen S, Liu T. Expression of Bcl-2 and microRNAs in cardiac tissues of patients with dilated cardiomyopathy. *Mol Med Rep* 2017; **15**: 359-365 [PMID: [27922664](#) DOI: [10.3892/mmr.2016.5977](#)]
- 60 **Besler C**, Urban D, Watzka S, Lang D, Rommel KP, Kandolf R, Klingel K, Thiele H, Linke A, Schuler G, Adams V, Lurz P. Endomyocardial miR-133a levels correlate with myocardial inflammation, improved left ventricular function, and clinical outcome in patients with inflammatory cardiomyopathy. *Eur J Heart Fail* 2016; **18**: 1442-1451 [PMID: [27292200](#) DOI: [10.1002/ehf.579](#)]
- 61 **Satoh M**, Minami Y, Takahashi Y, Tabuchi T, Nakamura M. A cellular microRNA, let-7i, is a novel biomarker for clinical outcome in patients with dilated cardiomyopathy. *J Card Fail* 2011; **17**: 923-929 [PMID: [22041329](#) DOI: [10.1016/j.cardfail.2011.07.012](#)]
- 62 **Carrillo-Salinas FJ**, Ngwenyama N, Anastasiou M, Kaur K, Alcaide P. Heart Inflammation: Immune Cell Roles and Roads to the Heart. *Am J Pathol* 2019; **189**: 1482-1494 [PMID: [31108102](#) DOI: [10.1016/j.ajpath.2019.04.009](#)]
- 63 **Noutsias M**, Rohde M, Göldner K, Block A, Blunert K, Hemaïdan L, Hummel M, Blohm JH, Lassner D, Kühl U, Schultheiss HP, Volk HD, Kotsch K. Expression of functional T-cell markers and T-cell receptor Vbeta repertoire in endomyocardial biopsies from patients presenting with acute myocarditis and dilated cardiomyopathy. *Eur J Heart Fail* 2011; **13**: 611-618 [PMID: [21422001](#) DOI: [10.1093/eurjhf/hfr014](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

