

# World Journal of *Radiology*

*World J Radiol* 2022 April 28; 14(4): 70-106



**MINIREVIEWS**

- 70 Focal liver lesions in cirrhosis: Role of contrast-enhanced ultrasonography  
*Bartolotta TV, Randazzo A, Bruno E, Taibbi A*

**ORIGINAL ARTICLE****Retrospective Cohort Study**

- 82 Decreased cross-sectional muscle area in male patients with clear cell renal cell carcinoma and peritumoral collateral vessels  
*Greco F, Beomonte Zobel B, Mallio CA*

**Retrospective Study**

- 91 Outcome of percutaneous drainage for septic complications coexisted with COVID-19  
*Deif MA, Mounir AM, Abo-Hedibah SA, Abdel Khalek AM, Elmokadem AH*

**LETTER TO THE EDITOR**

- 104 Follow-up computed tomography scan in post-COVID-19 pneumonia  
*Chohan A, Choudhury S, Dadhwal R, Vakil AP, Franco R, Taweeseed PT*

**ABOUT COVER**

Editorial Board Member of *World Journal of Radiology*, Xian-Li Lv, MD, Associate Professor, Department of Neurointervention, Beijing Tsinghua Changgung Hospital, School of Clinical Medicine, Tsinghua University, Beijing 102218, China. lxl02301@btch.edu.cn

**AIMS AND SCOPE**

The primary aim of *World Journal of Radiology (WJR, World J Radiol)* is to provide scholars and readers from various fields of radiology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

*WJR* mainly publishes articles reporting research results and findings obtained in the field of radiology and covering a wide range of topics including state of the art information on cardiopulmonary imaging, gastrointestinal imaging, genitourinary imaging, musculoskeletal imaging, neuroradiology/head and neck imaging, nuclear medicine and molecular imaging, pediatric imaging, vascular and interventional radiology, and women's imaging.

**INDEXING/ABSTRACTING**

The *WJR* is now abstracted and indexed in Emerging Sources Citation Index (Web of Science), PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2021 edition of Journal Citation Reports® cites the 2020 Journal Citation Indicator (JCI) for *WJR* as 0.51.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Wen-Wen Qi, Production Department Director: Xu Guo; Editorial Office Director: Jia-Ping Yan.

**NAME OF JOURNAL**

*World Journal of Radiology*

**ISSN**

ISSN 1949-8470 (online)

**LAUNCH DATE**

January 31, 2009

**FREQUENCY**

Monthly

**EDITORS-IN-CHIEF**

Thomas J Vogl

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/1949-8470/editorialboard.htm>

**PUBLICATION DATE**

April 28, 2022

**COPYRIGHT**

© 2022 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

## Follow-up computed tomography scan in post-COVID-19 pneumonia

Asad Chohan, Saiara Choudhury, Rahul Dadhwal, Abhay P Vakil, Rene Franco, Pahnwat Tonya Taweeseedt

**Specialty type:** Radiology, nuclear medicine and medical imaging

**Provenance and peer review:**

Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): A  
Grade B (Very good): B  
Grade C (Good): C  
Grade D (Fair): 0  
Grade E (Poor): 0

**P-Reviewer:** Arteaga-Livias K, Peru; Naswhan AJ, Qatar; Valencia GA, Peru

**Received:** December 17, 2021

**Peer-review started:** December 17, 2021

**First decision:** February 21, 2022

**Revised:** February 24, 2022

**Accepted:** March 26, 2022

**Article in press:** March 26, 2022

**Published online:** April 28, 2022



**Asad Chohan, Saiara Choudhury, Rahul Dadhwal, Abhay P Vakil, Rene Franco, Pahnwat Tonya Taweeseedt,** Pulmonary Medicine, Corpus Christi Medical Center, Corpus Christi, TX 78411, United States

**Corresponding author:** Pahnwat Tonya Taweeseedt, MD, Academic Fellow, Pulmonary Medicine, Corpus Christi Medical Center, 3315 S Alameda St, Corpus Christi, TX 78411, United States. [pahnwatt@gmail.com](mailto:pahnwatt@gmail.com)

### Abstract

The coronavirus disease 2019 (COVID-19) global pandemic can be a severe illness that leads to morbidity and mortality. With the increasing number of COVID-19 pneumonia survivors, several long-term changes may persist, including abnormal imaging of lung parenchyma. In addition to the clinical course, it is vital to follow up on pulmonary imaging during the post-infectious period, which is not routinely required in other common pulmonary diagnoses. Computed tomography (CT) scan of the chest is an effective and diagnostic tool for pneumonia which gives an insight into structural abnormalities within the lungs, complications, and possible progression of the disease. Several studies have monitored COVID-19 pneumonia and its complications using serial CT chest imaging from the initial phase of infection, hospitalization, and post-discharge. Nonetheless, long-term follow-up imaging data in post-COVID-19 is still limited. We have summarized the findings utilizing a systematic review of the literature regarding COVID-19 pneumonia imaging, including long-term follow-up.

**Key Words:** COVID-19; Pneumonia; Computed tomography; Evolution; Progression

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Changes seen in computed tomography imaging related to coronavirus disease 2019 (COVID-19) pneumonia appear to progress and peak around two weeks post-hospitalization. Overall improvement and complete resolution of COVID-19 pneumonia-related changes imaging can be seen in the majority of the patients with long-term follow-up. We have summarized the findings utilizing a systematic review of the literature regarding COVID-19 pneumonia imaging, including long-term follow-up.

**Citation:** Chohan A, Choudhury S, Dadhwal R, Vakil AP, Franco R, Taweeseedt PT. Follow-up computed tomography scan in post-COVID-19 pneumonia. *World J Radiol* 2022; 14(4): 104-106

**URL:** <https://www.wjgnet.com/1949-8470/full/v14/i4/104.htm>

**DOI:** <https://dx.doi.org/10.4329/wjr.v14.i4.104>

## TO THE EDITOR

We read the article titled “Review on radiological evolution of COVID-19 pneumonia using computed tomography” by Casartelli *et al*[1] with keen interest. A chest computed tomography (CT) scan can be a useful diagnostic tool in a high-prevalence or pandemic situation, especially with clinical correlation. Risk stratification and assessing the progression of disease are also effective uses of CT chest imaging in coronavirus disease 2019 (COVID-19) patients. Given the global spread of COVID-19 and the magnitude of both direct and indirect effects of the disease, a CT scan of the chest can help in long-term prognostication in patients who survive.

Multiple studies have concluded that with disease progression, certain initial CT findings in COVID-19 can evolve with a specific pattern and regularity. COVID-19 pneumonia-related changes seen on CT chest imaging typically progress rapidly, plateau, and subsequently start to resolve thereafter. Changes in CT imaging vary widely from six to seventeen days but typically stabilize within the first two weeks of COVID-19 pneumonia. In the short term, some of the features seem to recur, with scans mostly showing consolidations and ground-glass opacities (GGO). Besides GGO, chest CT characteristics that indicate the reparation, including subpleural, linear opacities, and fibrotic changes, were also reported. A sign termed “fishing net on trees” has also been reported. Some reports have also mentioned interseptal thickening and fibrous streaks[1,2]. Three weeks post-discharge, GGO and fibrous stripes have been seen, while after four weeks, mostly linear opacities remained. The “tinted” sign and bronchovascular bundle distortion have also been mentioned. The bronchovascular bundle distortion could possibly be a result of inflammatory destruction or subsegmental atelectasis. The latter two signs mentioned above may signify the gradual resolution of inflammation with re-expansion of alveoli based on previous reports. This review included reports with follow-up durations of up to four months[1].

In a study conducted by Pan *et al*[3], two hundred nine patients with COVID-19 infection, who had been admitted to the hospital, undertook serial chest CT at three, seven, and twelve months. One-year CT chest follow-up revealed residual linear lesions, multiple areas of reticular opacities/cysts, and complete resolution in 12%, 13%, and 75%, respectively[3]. In another study conducted by Guan *et al*[4], CT results of 69 patients who had COVID-19 infection were assessed in three different phases: Initial CT, peak CT, and CT prior to discharge. Peak CT in this study was the highest attenuation of the density without alteration in size during COVID-19 progression or the maximal size of lesion on CT which is the most common pattern. The intervals were closely correlated to lobe scores and CT appearances; the higher the lobe score, the longer the intervals. The lobe score was calculated according to the percentage of the lesion in one lobe with the zero equals to no lesion, one equals more than 0% to less than 25%, two equals 25% to less than 50%, three equals 50% to less than 75%, and four equals to 75% or more. While the utilization of lobe score may be beneficial, further studies are necessary to assess its effectiveness on a larger scale.

The duration of initial interval is inversely correlated with the amount of consolidations, air bronchograms, and irregular lines[3]. The intervals will be longer if irregular and reticular lines are seen on the peak CT and pre-discharge CT. After that, COVID-19 pneumonia lesions on the CT chest may resolve completely, while GGO, irregular and reticular lines may remain[3]. In a similar study conducted by Chen *et al*[5], 41 patients were followed with chest CT during the hospital stay and at two weeks, one month, three months, six months, and one year after discharge. The study concluded that patients showed continuous improvement on lung CT scans during the 1-year follow-up time; however residual lesions (GGO and reticular patterns) may still be found, which are associated with lung volume parameters and risk of developing lung opacities[5]. Liu *et al*[6] retrospectively evaluated chest CT follow-ups on 51 patients with COVID-19 performed on the day prior to discharge, two weeks post-discharge, and four weeks post-discharge. The results of this study indicated that changes seen were significantly reduced, including density reduction on follow-up scans as compared to the scans done at the time of discharge.

Unlike the systematic review by Casartelli *et al*[1], these results showed that 64.7% of discharged patients progressed to complete resolution of previously seen lung lesions at 4-wk follow-up, indicating that damaged lung tissue could heal in patients with COVID-19 pneumonia[5]. In another study conducted by Liu *et al*[7], 41 patients diagnosed with COVID-19 were followed up after seven months with chest CT and cardiopulmonary exercise testing. The predominant chest CT patterns at seven months included parenchymal bands (41%), interlobular septal thickening (32%), and traction bronchiectasis (29%). Sixty-one percent of the patients achieved complete radiological resolution, while 29% went on to develop pulmonary fibrosis. Those patients who went on to develop fibrotic lung disease appeared to have an increased risk due to older age and comorbid conditions[7].

While CT scan of the chest is an effective tool in COVID-19 patients, the side effects to patients of repeat irradiation need to be kept in mind and the use of low dose CT to follow up these patients can be considered. In conclusion, CT scans of the chest are an effective diagnostic tool which can provide insight into the structural pathology of pulmonary disease, its progression, and its association with long-term effects. Future studies should be utilized to define its utility in determining long-term progression in patients with COVID-19 pneumonia.

---

## FOOTNOTES

---

**Author contributions:** Chohan A and Taweeseed PT wrote the letter; Choudhury S wrote the letter; Franco R, Dadhwal R, Vakil AP and Taweeseed PT revised the letter.

**Conflict-of-interest statement:** All authors declare no conflicts-of-interest related to this article.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** United States

**ORCID number:** Asad Chohan 0000-0002-1801-9792; Saiara Choudhury 0000-0001-7225-4905; Rahul Dadhwal 0000-0001-6963-1466; Abhay P Vakil 0000-0003-4947-0233; Rene Franco 0000-0002-0684-7734; Pahnwat Tonya Taweeseed 0000-0002-5791-6920.

**S-Editor:** Wang JJ

**L-Editor:** A

**P-Editor:** Wang JJ

---

## REFERENCES

---

- 1 **Casartelli C**, Perrone F, Balbi M, Alfieri V, Milanese G, Buti S, Silva M, Sverzellati N, Bersanelli M. Review on radiological evolution of COVID-19 pneumonia using computed tomography. *World J Radiol* 2021; **13**: 294-306 [PMID: 34630915 DOI: 10.4329/wjr.v13.i9.294]
- 2 **Mejia-Zambrano H**. Radiological and functional pulmonary complications in patients recovered from COVID-19. *Microb Infect Chemother* 2021 [DOI: 10.54034/mic.e1217]
- 3 **Pan F**, Yang L, Liang B, Ye T, Li L, Liu D, Wang J, Hesketh RL, Zheng C. Chest CT Patterns from Diagnosis to 1 Year of Follow-up in Patients with COVID-19. *Radiology* 2022; **302**: 709-719 [PMID: 34609153 DOI: 10.1148/radiol.2021211199]
- 4 **Guan CS**, Lv ZB, Li JJ, Du YN, Chen H, Cui T, Guo N, Chen BD, Xie RM. CT appearances, patterns of progression, and follow-up of COVID-19: evaluation on thin-section CT. *Insights Imaging* 2021; **12**: 73 [PMID: 34110540 DOI: 10.1186/s13244-021-01019-0]
- 5 **Chen Y**, Ding C, Yu L, Guo W, Feng X, Su J, Xu T, Ren C, Shi D, Wu W, Yi P, Liu J, Tao J, Lang G, Li Y, Xu M, Sheng J, Li L, Xu K. One-year follow-up of chest CT findings in patients after SARS-CoV-2 infection. *BMC Med* 2021; **19**: 191 [PMID: 34365975 DOI: 10.1186/s12916-021-02056-8]
- 6 **Liu C**, Ye L, Xia R, Zheng X, Yuan C, Wang Z, Lin R, Shi D, Gao Y, Yao J, Sun Q, Wang X, Jin M. Chest Computed Tomography and Clinical Follow-Up of Discharged Patients with COVID-19 in Wenzhou City, Zhejiang, China. *Ann Am Thorac Soc* 2020; **17**: 1231-1237 [PMID: 32692945 DOI: 10.1513/AnnalsATS.202004-324OC]
- 7 **Liu M**, Lv F, Huang Y, Xiao K. Follow-Up Study of the Chest CT Characteristics of COVID-19 Survivors Seven Months After Recovery. *Front Med (Lausanne)* 2021; **8**: 636298 [PMID: 33732719 DOI: 10.3389/fmed.2021.636298]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA  
**Telephone:** +1-925-3991568  
**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
**Help Desk:** <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>

