

# World Journal of *Clinical Cases*

*World J Clin Cases* 2022 July 6; 10(19): 6341-6758



**MINIREVIEWS**

- 6341** Review of clinical characteristics, immune responses and regulatory mechanisms of hepatitis E-associated liver failure  
*Chen C, Zhang SY, Chen L*
- 6349** Current guidelines for *Helicobacter pylori* treatment in East Asia 2022: Differences among China, Japan, and South Korea  
*Cho JH, Jin SY*
- 6360** Review of epidermal growth factor receptor-tyrosine kinase inhibitors administration to non-small-cell lung cancer patients undergoing hemodialysis  
*Lan CC, Hsieh PC, Huang CY, Yang MC, Su WL, Wu CW, Wu YK*

**ORIGINAL ARTICLE****Case Control Study**

- 6370** Pregnancy-related psychopathology: A comparison between pre-COVID-19 and COVID-19-related social restriction periods  
*Chieffo D, Avallone C, Serio A, Kotzalidis GD, Balocchi M, De Luca I, Hirsch D, Gonzalez del Castillo A, Lanzotti P, Marano G, Rinaldi L, Lanzone A, Mercuri E, Mazza M, Sani G*
- 6385** Intestinal mucosal barrier in functional constipation: Dose it change?  
*Wang JK, Wei W, Zhao DY, Wang HF, Zhang YL, Lei JP, Yao SK*

**Retrospective Cohort Study**

- 6399** Identification of risk factors for surgical site infection after type II and type III tibial pilon fracture surgery  
*Hu H, Zhang J, Xie XG, Dai YK, Huang X*

**Retrospective Study**

- 6406** Total knee arthroplasty in Ranawat II valgus deformity with enlarged femoral valgus cut angle: A new technique to achieve balanced gap  
*Lv SJ, Wang XJ, Huang JF, Mao Q, He BJ, Tong PJ*
- 6417** Preliminary evidence in treatment of eosinophilic gastroenteritis in children: A case series  
*Chen Y, Sun M*
- 6428** Self-made wire loop snare successfully treats gastric persimmon stone under endoscopy  
*Xu W, Liu XB, Li SB, Deng WP, Tong Q*
- 6437** Neoadjuvant transcatheter arterial chemoembolization and systemic chemotherapy for the treatment of undifferentiated embryonal sarcoma of the liver in children  
*He M, Cai JB, Lai C, Mao JQ, Xiong JN, Guan ZH, Li LJ, Shu Q, Ying MD, Wang JH*

- 6446 Effect of cold snare polypectomy for small colorectal polyps

*Meng QQ, Rao M, Gao PJ*

- 6456 Field evaluation of COVID-19 rapid antigen test: Are rapid antigen tests less reliable among the elderly?

*Tabain I, Cucevic D, Skreb N, Mrzljak A, Ferencak I, Hruskar Z, Mistic A, Kuzle J, Skoda AM, Jankovic H, Vilibic-Cavlek T*

### Observational Study

- 6464 Tracheobronchial intubation using flexible bronchoscopy in children with Pierre Robin sequence: Nursing considerations for complications

*Ye YL, Zhang CF, Xu LZ, Fan HF, Peng JZ, Lu G, Hu XY*

- 6472 Family relationship of nurses in COVID-19 pandemic: A qualitative study

*Çelik MY, Kiliç M*

### META-ANALYSIS

- 6483 Diagnostic accuracy of  $\geq 16$ -slice spiral computed tomography for local staging of colon cancer: A systematic review and meta-analysis

*Liu D, Sun LM, Liang JH, Song L, Liu XP*

### CASE REPORT

- 6496 Delayed-onset endophthalmitis associated with *Achromobacter* species developed in acute form several months after cataract surgery: Three case reports

*Kim TH, Lee SJ, Nam KY*

- 6501 Sustained dialysis with misplaced peritoneal dialysis catheter outside peritoneum: A case report

*Shen QQ, Behera TR, Chen LL, Attia D, Han F*

- 6507 Arteriovenous thrombotic events in a patient with advanced lung cancer following bevacizumab plus chemotherapy: A case report

*Kong Y, Xu XC, Hong L*

- 6514 Endoscopic ultrasound radiofrequency ablation of pancreatic insulinoma in elderly patients: Three case reports

*Rossi G, Petrone MC, Capurso G, Partelli S, Falconi M, Arcidiacono PG*

- 6520 Acute choroidal involvement in lupus nephritis: A case report and review of literature

*Yao Y, Wang HX, Liu LW, Ding YL, Sheng JE, Deng XH, Liu B*

- 6529 Triple A syndrome-related achalasia treated by per-oral endoscopic myotomy: Three case reports

*Liu FC, Feng YL, Yang AM, Guo T*

- 6536 Choroidal thickening with serous retinal detachment in BRAF/MEK inhibitor-induced uveitis: A case report

*Kiraly P, Groznic AL, Valentinčić NV, Mekjavić PJ, Urbančić M, Ocvirk J, Mesti T*

- 6543 Esophageal granular cell tumor: A case report

*Chen YL, Zhou J, Yu HL*

- 6548** Hem-o-lok clip migration to the common bile duct after laparoscopic common bile duct exploration: A case report  
*Liu DR, Wu JH, Shi JT, Zhu HB, Li C*
- 6555** Chidamide and sintilimab combination in diffuse large B-cell lymphoma progressing after chimeric antigen receptor T therapy  
*Hao YY, Chen PP, Yuan XG, Zhao AQ, Liang Y, Liu H, Qian WB*
- 6563** Relapsing polychondritis with isolated tracheobronchial involvement complicated with Sjogren's syndrome: A case report  
*Chen JY, Li XY, Zong C*
- 6571** Acute methanol poisoning with bilateral diffuse cerebral hemorrhage: A case report  
*Li J, Feng ZJ, Liu L, Ma YJ*
- 6580** Immunoabsorption therapy for Klinefelter syndrome with antiphospholipid syndrome in a patient: A case report  
*Song Y, Xiao YZ, Wang C, Du R*
- 6587** Roxadustat for treatment of anemia in a cancer patient with end-stage renal disease: A case report  
*Zhou QQ, Li J, Liu B, Wang CL*
- 6595** Imaging-based diagnosis for extraskeletal Ewing sarcoma in pediatrics: A case report  
*Chen ZH, Guo HQ, Chen JJ, Zhang Y, Zhao L*
- 6602** Unusual course of congenital complete heart block in an adult: A case report  
*Su LN, Wu MY, Cui YX, Lee CY, Song JX, Chen H*
- 6609** Penile metastasis from rectal carcinoma: A case report  
*Sun JJ, Zhang SY, Tian JJ, Jin BY*
- 6617** Isolated cryptococcal osteomyelitis of the ulna in an immunocompetent patient: A case report  
*Ma JL, Liao L, Wan T, Yang FC*
- 6626** Magnetic resonance imaging features of intrahepatic extramedullary hematopoiesis: Three case reports  
*Luo M, Chen JW, Xie CM*
- 6636** Giant retroperitoneal liposarcoma treated with radical conservative surgery: A case report and review of literature  
*Lieto E, Cardella F, Erario S, Del Sorbo G, Reginelli A, Galizia G, Urraro F, Panarese I, Auricchio A*
- 6647** Transplanted kidney loss during colorectal cancer chemotherapy: A case report  
*Pośpiech M, Kolonko A, Nieszporek T, Kozak S, Kozaczka A, Karkoszka H, Winder M, Chudek J*
- 6656** Massive gastrointestinal bleeding after endoscopic rubber band ligation of internal hemorrhoids: A case report  
*Jiang YD, Liu Y, Wu JD, Li GP, Liu J, Hou XH, Song J*

- 6664** Mills' syndrome is a unique entity of upper motor neuron disease with N-shaped progression: Three case reports  
*Zhang ZY, Ouyang ZY, Zhao GH, Fang JJ*
- 6672** Entire process of electrocardiogram recording of Wellens syndrome: A case report  
*Tang N, Li YH, Kang L, Li R, Chu QM*
- 6679** Retroperitoneal tumor finally diagnosed as a bronchogenic cyst: A case report and review of literature  
*Gong YY, Qian X, Liang B, Jiang MD, Liu J, Tao X, Luo J, Liu HJ, Feng YG*
- 6688** Successful treatment of Morbihan disease with total glucosides of paeony: A case report  
*Zhou LF, Lu R*
- 6695** Ant sting-induced whole-body pustules in an inebriated male: A case report  
*Chen SQ, Yang T, Lan LF, Chen XM, Huang DB, Zeng ZL, Ye XY, Wan CL, Li LN*
- 6702** Plastic surgery for giant metastatic endometrioid adenocarcinoma in the abdominal wall: A case report and review of literature  
*Wang JY, Wang ZQ, Liang SC, Li GX, Shi JL, Wang JL*
- 6710** Delayed-release oral mesalamine tablet mimicking a small jejunal gastrointestinal stromal tumor: A case report  
*Frosio F, Rausa E, Marra P, Boutron-Ruault MC, Lucianetti A*
- 6716** Concurrent alcoholic cirrhosis and malignant peritoneal mesothelioma in a patient: A case report  
*Liu L, Zhu XY, Zong WJ, Chu CL, Zhu JY, Shen XJ*
- 6722** Two smoking-related lesions in the same pulmonary lobe of squamous cell carcinoma and pulmonary Langerhans cell histiocytosis: A case report  
*Gencer A, Ozcibik G, Karakas FG, Sarbay I, Batur S, Borekci S, Turna A*
- 6728** Proprotein convertase subtilisin/kexin type 9 inhibitor non responses in an adult with a history of coronary revascularization: A case report  
*Yang L, Xiao YY, Shao L, Ouyang CS, Hu Y, Li B, Lei LF, Wang H*
- 6736** Multimodal imaging study of lipemia retinalis with diabetic retinopathy: A case report  
*Zhang SJ, Yan ZY, Yuan LF, Wang YH, Wang LF*
- 6744** Primary squamous cell carcinoma of the liver: A case report  
*Kang LM, Yu DP, Zheng Y, Zhou YH*
- 6750** Tumor-to-tumor metastasis of clear cell renal cell carcinoma to contralateral synchronous pheochromocytoma: A case report  
*Wen HY, Hou J, Zeng H, Zhou Q, Chen N*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Abdulqadir Jeprel Naswhan, MSc, RN, Director, Research Scientist, Senior Lecturer, Senior Researcher, Nursing for Education and Practice Development, Hamad Medical Corporation, Doha 576214, Qatar. anashwan@hamad.qa

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

*WJCC* mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The *WJCC* is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for *WJCC* as 1.337; IF without journal self cites: 1.301; 5-year IF: 1.742; Journal Citation Indicator: 0.33; Ranking: 119 among 169 journals in medicine, general and internal; and Quartile category: Q3. The *WJCC*'s CiteScore for 2020 is 0.8 and Scopus CiteScore rank 2020: General Medicine is 493/793.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Xu Guo*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

July 6, 2022

**COPYRIGHT**

© 2022 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

## Current guidelines for *Helicobacter pylori* treatment in East Asia 2022: Differences among China, Japan, and South Korea

Jun-Hyung Cho, So-Young Jin

**Specialty type:** Gastroenterology and hepatology

**Provenance and peer review:** Invited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): B  
Grade C (Good): C  
Grade D (Fair): D  
Grade E (Poor): 0

**P-Reviewer:** Herawati F, Indonesia; Megraud F, France

**Received:** January 13, 2022

**Peer-review started:** January 13, 2022

**First decision:** March 8, 2022

**Revised:** March 11, 2022

**Accepted:** May 8, 2022

**Article in press:** May 8, 2022

**Published online:** July 6, 2022



**Jun-Hyung Cho**, Digestive Disease Center, Soonchunhyang University Hospital, Seoul 04401, South Korea

**So-Young Jin**, Department of Pathology, Soonchunhyang University Hospital, Seoul 04401, South Korea

**Corresponding author:** Jun-Hyung Cho, MD, PhD, Associate Professor, Digestive Disease Center, Soonchunhyang University Hospital, 59, Daesagwan-ro, Yongsan-gu, Seoul 04401, South Korea. [chojhmd@naver.com](mailto:chojhmd@naver.com)

### Abstract

*Helicobacter pylori* (*H. pylori*) infection is highly prevalent in East Asia. The overall seroprevalence rate of *H. pylori* infection is 44.2% in China, 37.6%-43.2% in Japan, and 51.0% in South Korea. *H. pylori* can cause peptic ulcer disease and gastric cancer. East Asian countries have high rates of gastric cancer (age-standardized incidence rate: 20-30 per 100000). The Kyoto global consensus report emphasized that *H. pylori* gastritis should be considered the main cause for the development of gastric cancer. *H. pylori* treatment guidelines in China, Japan, and South Korea have recently been revised according to data from each of those countries. However, emerging antibiotic resistance is an important barrier to *H. pylori* eradication. The recommended *H. pylori* treatment regimens differ among those three East Asian countries. In this review, recent guidelines and up-to-date research on *H. pylori* treatment regimens from China, Japan, and South Korea are discussed.

**Key Words:** *Helicobacter pylori*; Treatment; Antibiotic resistance; China; Japan; South Korea

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Since 2000, the standard triple regimen containing clarithromycin (CAM) has been used as a legacy therapy to eradicate *Helicobacter pylori* (*H. pylori*). Resistance to CAM by *H. pylori* has increased to > 15% in East Asia. First-line eradication rates below 80% are strongly associated with CAM-resistant *H. pylori* strain emergence. *H. pylori* treatment guidelines in China, Japan, and South Korea were revised according to new data. In China, adding bismuth to *H. pylori* regimens was recommended as an empirical first-line treatment. In Japan, *H. pylori* treatment success increased when the potassium-competitive acid blocker (P-CAB) was introduced. In South Korea, tailored *H. pylori* eradication based on molecular testing for CAM resistance is used as the first-line treatment option. Dual therapy involving frequent administration of high-dose amoxicillin has shown good efficacy for *H. pylori* eradication in clinical trials. Furthermore, P-CABs, with their rapid and strong acid-suppressing activity, may contribute to successful *H. pylori* treatment in future.

**Citation:** Cho JH, Jin SY. Current guidelines for *Helicobacter pylori* treatment in East Asia 2022: Differences among China, Japan, and South Korea. *World J Clin Cases* 2022; 10(19): 6349-6359

**URL:** <https://www.wjgnet.com/2307-8960/full/v10/i19/6349.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v10.i19.6349>


## INTRODUCTION

*Helicobacter pylori* (*H. pylori*) can cause peptic ulcer disease and gastric cancer[1]. In *H. pylori*-infected stomachs, gastric atrophy and intestinal metaplasia are linked to gastric cancer development[2]. *H. pylori* is present in more than half of the world's population[3]. The overall seroprevalence rate of *H. pylori* infection is 44.2% in China, 37.6%-43.2% in Japan, and 51.0% in South Korea[4-6]. Gastric cancer is the fourth most common cause of cancer-related mortality worldwide[7]. Almost half of the incident cases and deaths occur in East Asia. In 2019, the age-standardized incidence rate of gastric cancer per 100000 was 30.64 in China, 28.29 in Japan, and 28.67 in South Korea[8].

The Kyoto Global Consensus recommends *H. pylori* treatment to prevent gastric cancer in countries with a high incidence thereof[9]. In Japan, eradication therapy for all *H. pylori*-positive subjects has been covered by the national insurance system since 2013[10]. In China, *H. pylori* treatment was strongly recommended for preventing primary gastric cancer in a recent consensus report[11]. In 2018, the South Korean government insurance system started to cover eradication therapy for *H. pylori* gastritis. However, primary antibiotic resistance of *H. pylori* has increased in East Asia, so obtaining successful therapeutic outcomes using antibiotic regimens is challenging[12]. To prevent primary gastric cancer, *H. pylori* should be successfully eradicated. Based on a comparison of recent guidelines, this review focuses on the current status of *H. pylori* treatment in China, Japan, and South Korea in terms of *H. pylori* resistance to antibiotics, recommended *H. pylori* treatment regimens, and up-to-date results of *H. pylori* therapy in the three countries.

## H. PYLORI RESISTANCE TO ANTIBIOTICS IN EAST ASIA

Since 2000, the standard triple regimen containing clarithromycin (CAM) has been used as a legacy therapy to eradicate *H. pylori*[13]. Macrolides have been widely used to treat other infectious diseases. Accordingly, the number of CAM-resistant *H. pylori* strains has increased rapidly in many countries over the past decade[14]. In East Asia, CAM resistance of *H. pylori* has increased by > 15%. Therefore, first-line *H. pylori* eradication rates using triple therapy have decreased to < 80%[15]. *H. pylori* resistance rates to metronidazole (MDZ) and fluoroquinolone are also high in East Asia (Figure 1). In China, primary resistance of *H. pylori* to CAM, MDZ, and levofloxacin (LVFX) is high and has increased over time (28.9%, 63.8%, and 28.0%, respectively)[16]. These patterns of *H. pylori* resistance are similar to those in South Korea. A recent nationwide study reported *H. pylori* resistance rates to CAM, MDZ, and LVFX of 17.8%, 29.5%, and 37.0%, respectively[17]. In Japan, the CAM resistance rate has increased gradually from 7% in 2000 to 38.5% in 2014[18]. The resistance rate of *H. pylori* to LVFX is relatively high at approximately 15%[19]. In Japan, MDZ resistance is lower (< 10%) than in other Asian countries. Low MDZ resistance against *H. pylori* is thought to be correlated with antibiotic consumption in the community. The use of MDZ is strictly regulated in Japan, where it has been approved for the treatment of only selected diseases, such as trichomoniasis[20]. In contrast, the *H. pylori* resistance rates to amoxicillin (AMX) and tetracycline (TET) are equally low among the three countries (3.1% and 3.9% in China, 3.0% and 2.0% in Japan, and 9.5% and 0% in South Korea, respectively)[21].



**Figure 1** Prevalence of primary *Helicobacter pylori* resistance to antibiotics in China, Japan, and South Korea. CAM: Clarithromycin; MDZ: Metronidazole; LVFX: Levofloxacin; AMX: Amoxicillin; TET: Tetracycline.

## RECOMMENDED *H. PYLORI* REGIMENS

Some differences in *H. pylori* treatment regimens exist among the guidelines of China, Japan, and South Korea. Bismuth-based *H. pylori* regimens are strongly recommended in China. In Japan, potassium-competitive acid blockers (P-CABs) are widely used to eradicate *H. pylori*. The current South Korean guidelines are similar to those of Western countries. Notably, molecular testing to detect CAM resistance is recommended as the initial treatment option.

### China

According to the Fifth Chinese Consensus for *H. pylori* Management of 2016, seven *H. pylori* therapies containing bismuth salt are recommended as empirical regimens[11]. The various antibiotic combinations include two of the following six antimicrobial agents: AMX, CAM, MDZ, TET, LVFX, and furazolidone (FZD). The recommended eradication regimens for *H. pylori* include AMX and CAM, AMX and MDZ, AMX and LVFX, AMX and FZD, AMX and TET, MDZ and TET, and FZD and TET (Table 1). The standard dose of a proton pump inhibitor (PPI) and 220 mg of bismuth are prescribed twice daily with the two antibiotics. Bismuth has long been used to treat peptic ulcer disease, dyspepsia, parasite infections, and infectious diarrhea[22]. The antibacterial effects of bismuth include inhibition of protein and cell wall synthesis in *H. pylori*. The main role of bismuth is to increase the eradication rate by 30%-40% in resistant *H. pylori* strains[23].

Zhang *et al*[24] reported on a 14-d modified bismuth quadruple therapy (BQT) containing CAM or MDZ that was effective against *H. pylori* in a region with high resistance to CAM (26.5%) and MDZ (45.5%). CAM- and MDZ-containing regimens displayed high eradication rates of 88.8% and 88.9% in the intention-to-treat (ITT) analysis, and 94.9% and 96.9% in the per-protocol (PP) analysis, respectively. In an MDZ-containing regimen, 1 g of AMX twice daily was used as a substitute for 500 mg of TET four times daily. Notably, a modified BQT containing AMX and MDZ was demonstrated to be effective (> 90%) in MDZ-resistant *H. pylori* strains. A high dosage (1500 mg/d) of MDZ for 14 d can overcome *H. pylori* resistance[25]. Chen *et al*[26] performed a comparative study between the classic BQT and a modified one containing AMX and MDZ as a rescue *H. pylori* therapy. In comparison with the classic BQT, the modified BQT achieved a similar eradication rate (87.2%-95.3% vs 88.5%-93.7%). Adverse events occurred less frequently with the modified than the classic BQT regimen (34.0% vs 51.9%, *P* = 0.001).

FZD is a nitrofurantoin antibiotic effective against gram-negative and -positive bacteria[27]. Qiao *et al*[28] reported that an FZD-based BQT showed a similar first-line *H. pylori* eradication rate in PP analysis to a CAM-based BQT (95.8% vs 93.4%). The resistance rates of *H. pylori* to AMX, TET, and FZD remain low

**Table 1 Recommended regimens according to the *Helicobacter pylori* treatment guidelines in China, Japan, and South Korea**

Country	Regimen	Drugs	Duration (d)	
China	1	PPI bid + bismuth 220 mg bid + AMX 1000 mg bid + CAM 500 mg bid	14	
	2	PPI bid + bismuth 220 mg bid + AMX 1000 mg bid + MDZ 400 mg tid or qid	14	
	3	PPI bid + bismuth 220 mg bid + AMX 1000 mg bid + LVFX 500 mg qd or 200 mg bid	14	
	4	PPI bid + bismuth 220 mg bid + AMX 1000 mg bid + FZD 100 mg bid or tid	14	
	5	PPI bid + bismuth 220 mg bid + AMX 1000 mg bid + TET 500 mg tid or qid	14	
	6	PPI bid + bismuth 220 mg bid + MDZ 400 mg tid or qid + TET 500 mg tid or qid	14	
	7	PPI bid + bismuth 220 mg bid + FZD 100 mg bid + TET 500 mg tid or qid	14	
Japan	First-line	PPI or vonoprazan 20 mg bid + AMX 750 mg bid + CAM 200 mg bid	7	
	Second-line	PPI bid + AMX 750 mg bid + MDZ 250 mg bid	7	
	Third-line	PPI bid + AMX 750 mg or MDZ 250 mg bid + STFX 100 mg bid	7	
South Korea	First-line	Standard triple therapy	PPI bid + AMX 1000 mg bid + CAM 500 mg bid	14
		Sequential therapy	PPI bid + AMX 1000 mg bid (5 d), then CAM 500 mg bid + MDZ 500 mg bid (5 d)	10
		Concomitant therapy	PPI bid + AMX 1000 mg bid + CAM 500 mg bid + MDZ 500 mg bid	10
		Tailored therapy	Standard triple therapy (CAM-sensitive) or classic BQT (CAM-resistant)	7-14
	Second-line	Classic BQT	PPI bid + bismuth 120 mg qid + MDZ 500 mg tid + TET 500 mg qid	10-14
	Third-line	-	PPI bid + AMX 1000 mg bid + LVFX 500 mg qd or 250 mg bid	10-14

AMX: Amoxicillin; BQT: Bismuth quadruple therapy; CAM: Clarithromycin; FZD: Furazolidone; LVFX: Levofloxacin; MDZ: Metronidazole; PPI: Proton pump inhibitor; STFX: Sitafloxacin; TET: Tetracycline.

in China at < 5%[29]. Accordingly, a modified BQT containing FZD/AMX and FZD/TET achieved a > 90% *H. pylori* eradication rate in patients who did not respond to the previous treatment[30]. Compared to the classic BQT, two FZD-containing regimens resulted in less frequent adverse events. The *H. pylori* eradication efficacy of regimens employing AMX and TET may not be inferior to other regimens[31]. However, TET and FZD have limited availability[32].

Interestingly, the recommended *H. pylori* therapies were not categorized into first- and second-line regimens. This strategy is markedly different from Japanese and South Korean guidelines, in which the first- and second-line regimens are divided. Any of the seven regimens can be prescribed to patients to eradicate *H. pylori*. After failed initial therapy, a second-line regimen is selected from among the remaining regimens. Among the antibiotic combinations, the LVFX-containing regimen is not recommended as an initial treatment due to a high resistance rate[33]. It can be considered an alternative to rescue therapy in combination with bismuth salt. AMX, TET, and FZD can be reused after treatment failure because these drugs rarely produce secondary resistance. Repeat MDZ prescription requires an optimized dose (at least 1500-1600 mg/d). In contrast, reuse of CAM and LVFX should be avoided after failed eradication.

### Japan

Since *H. pylori* treatment for chronic gastritis was approved by the Japanese national health insurance system in 2013, prescriptions for eradication therapy have markedly increased. Approximately 8.5 million *H. pylori*-positive patients received eradication regimens from 2013 to 2019[34]. The revised 2016 guidelines were similar to previous ones[35]. As a first-line regimen, standard triple therapy was still recommended to eradicate *H. pylori* (Table 1). The AMX and CAM dosages were lower than those of China and South Korea. The recommended AMX and CAM doses are 750 and 200 mg twice daily, respectively. The 200 mg dose of CAM twice daily has similar efficacy to 400 mg twice daily, and no

significant difference was found between 7- and 14-d treatment durations. However, adverse drug events, such as dysgeusia, occurred more frequently during the 14-d therapy.

In Japan, the successful eradication rates for all types of *H. pylori* therapy decreased to < 80% in 2014. Since the P-CAB was launched in February 2015, the eradication rates of triple therapy, including AMX and CAM with P-CAB, have been significantly higher than those with conventional PPIs[36]. The novel P-CAB vonoprazan (VPZ) exerts a rapid and sustained suppressive effect on gastric acid for optimal *H. pylori* treatment[37]. The proportion of regimens including VPZ has increased rapidly, and is now 80% [34]. As first-line treatment, standard triple therapy containing 20 mg VPZ twice daily resulted in higher *H. pylori* eradication rates than those of a PPI-based regimen (86.4%-91.2% vs 71.7%-79.4%)[38,39]. There was no significant difference between the VPZ- and PPI-based triple regimens for eradicating CAM-susceptible *H. pylori* strains. *H. pylori* eradication rates were 87.3% and 76.5% ( $P = 0.21$ ) in the ITT analysis, and 88.9% and 86.7% ( $P = 0.77$ ) in the PP analysis, respectively[40]. However, VPZ was superior to the PPI-based triple regimen for CAM-resistant *H. pylori* strains (73.2%-87.5% vs 40.0%-53.8%)[41].

CAM resistance is a strong contributor to *H. pylori* eradication failure after first-line treatment[42]. Therefore, MDZ-based triple therapy with PPI/VPZ is recommended as the second-line eradication therapy. MDZ (250 mg) is prescribed twice daily for 7 d. The MDZ-based *H. pylori* treatment results in a higher eradication rate (> 90%) than CAM-based triple therapy because of the low MDZ resistance in Japan[43,44]. In a meta-analysis by Shinozaki *et al*[45], VPZ was more efficacious for second-line *H. pylori* eradication compared to conventional PPIs. Seven-day VPZ, AMX, and MDZ triple therapy can be a strong candidate as an empirical *H. pylori* regimen. Sitafloxacin (STFX)-based triple therapy has been recommended as a third-line therapy in combination with AMX or MDZ[46]. STFX is a new quinolone antibacterial agent expected to be efficacious due to its low minimum inhibitory concentration (MIC) for *H. pylori*, even for LVFX-resistant strains[47]. STFX has a good *H. pylori* eradication effect in combination with AMX or MDZ, even for third-line treatment[48].

### South Korea

First-line *H. pylori* therapies consist of three empirical regimens and one tailored eradication regimen (Table 1)[49]. One of the following empirical regimens can be chosen: standard triple therapy for 14 d, non-BQT for 10 d, or classic BQT for 10-14 d. Standard triple therapy consists of a standard PPI dose, 1000 mg of AMX, and 500 mg of CAM, twice daily for 14 d. The eradication rate of 14-d standard triple therapy is superior to those of the 7- and 10-d therapeutic regimens. The pooled eradication rate of 14-d therapy was 78.1%, which is unacceptable for successful *H. pylori* eradication.

Non-BQT has been divided into concomitant and sequential therapy. A PPI, 500 mg of CAM, 1 g of AMX, and 500 mg of MDZ are prescribed twice daily for 10 d for the concomitant therapy. The sequential therapy consists of a PPI and 1 g of AMX twice daily for 5 d, followed by the PPI, 500 mg of CAM, and 500 mg of MDZ twice daily for 5 d. Concomitant and sequential therapy resulted in good *H. pylori* eradication rates of 94.2% and 91.7% in a modified ITT analysis, and 95.6% and 91.4% in the PP analysis, respectively[50]. Sequential therapy was superior to 7- and 10-d standard triple therapy as a first-line treatment[51]. However, the role of sequential therapy in *H. pylori* eradication is diminishing according to Western guidelines[52]. Lee *et al*[53] reported a higher eradication rate for concomitant than sequential therapy (94.4% vs 84.4%,  $P = 0.018$ ). A recent nationwide study demonstrated that the *H. pylori* eradication rate of concomitant therapy is significantly higher than that of sequential therapy (91.8% vs 86.1%,  $P < 0.001$ )[54]. Antibiotic overuse and the emergence of multidrug-resistant *H. pylori* may be problematic for concomitant therapy[55].

As the last of the empirical first-line therapies, the classic BQT (standard dose PPI twice a day, MDZ 500 mg three times a day, TET 500 mg, and bismuth 120 mg four times a day) is administered for 10-14 d. However, the guidelines report high rates of adverse drug events. The relatively high rates of adverse events and high pill burden are problematic[56]. According to the South Korean nationwide registry database, the prescription rate of the classic BQT as a first-line therapy is currently low (2.63%)[57]. Thus, the classic BQT can be considered as a rescue therapy for other regimens.

Tailored *H. pylori* eradication shows promise for achieving more successful outcomes before treatment, compared to conventional therapies[58]. Antimicrobial susceptibility testing using a culture method has been recommended for effective first-line *H. pylori* eradication in regions with a high rate of antibiotic resistance[14]. However, susceptibility-guided therapy is time-consuming and requires specific expertise in clinical practice[59]. As an alternative, molecular methods for detecting CAM resistance are available for tailored *H. pylori* eradication in South Korea[60]. *H. pylori* resistance against CAM is mediated by point mutations in 23S ribosomal RNA[61]. The treatment regimen is selected based on the presence of the A2142G and A2143G point mutations that cause CAM resistance[62]. *H. pylori*-infected subjects without point mutations are treated with the 7-d standard triple regimen. Lee *et al*[63] demonstrated that the 7-d triple therapy was as effective as the 14-d therapy in patients without a point mutation. The classic BQT is recommended when A2142G and/or A2143G point mutations are detected. In contrast, MDZ-based triple therapy has an unacceptably low eradication rate in CAM-resistant *H. pylori* strains[64]. Tailored *H. pylori* eradication using molecular testing was more efficacious as a first-line treatment than standard CAM-based triple therapy (97.0% vs 81.8%)[65]. A recent study showed that the *H. pylori* eradication rate was similar between tailored eradication using DPO-PCR and

the classic BQT (96.0% vs 95.7%,  $P = 0.9$ )[66]. Adverse drug events occurred less frequently with tailored eradication than BQT (12.0% vs 43.7%,  $P < 0.001$ ).

## UP-TO-DATE RESEARCH ON *H. PYLORI* THERAPY

*H. pylori* is an infectious disease; as such, the general principles regarding the treatment of infectious diseases, such as antimicrobial stewardship, are relevant to *H. pylori*[67]. Accordingly, recent research has focused on the optimal drug regimen in terms of dose, treatment duration, and minimal adverse events[68]. AMX, a beta-lactam antibiotic, has a half-life of approximately one hour, exhibiting time-dependent killing[69]. Therefore, frequent dosing has clinical advantages which help to maintain the plasma concentration higher than the MIC. The time above the MIC can reach 24 hours if 500 mg of AMX is dosed four times a day[70]. Along with efficacious antibiotics, sufficient and continuous acid suppression is required for successful eradication[71]. When *H. pylori* enters a replicative state between pH 6 and 8, the pathogen becomes highly susceptible to antibiotics, such as AMX. Recently, P-CABs have been found to increase the intragastric pH to 6 or more, which improves antibiotic stability and bioavailability[72].

### Dual therapy

The primary and secondary *H. pylori* resistance rates to AMX are low. In 1989, PPI-AMX dual therapy (DT) was first used to eradicate *H. pylori*[73]. However, the dosage of AMX, dosing intervals, and duration of therapy differed among previous investigators. Thereafter, a satisfactory *H. pylori* eradication using DT has not been achieved consistently[74]. DT has good *H. pylori* eradication efficacy, and is gaining increasing attention worldwide. In 2015, Yang *et al*[75] introduced a modified 14-d DT by increasing the dosage and frequency of administration (second-generation PPI and 750 mg of AMX four times daily). In a meta-analysis by Gao *et al*[76], DT and other commonly used regimens achieved similar efficacies in the ITT analysis (83.2% vs 85.3%,  $P = 0.87$ ) and PP analysis (87.5% vs 90.1%,  $P = 0.33$ ). In the DT group, drug-related adverse events occurred less frequently compared to the current mainstream therapy recommended by guidelines (12.9% vs 28.0%,  $P < 0.001$ ).

In a meta-analysis, Li *et al*[77] included randomized controlled studies in which both a PPI and AMX were administered four times daily. As a first-line *H. pylori* treatment, administering high-dose DT resulted in a higher eradication rate than other regimens in ITT analysis (89.8% vs 84.2%,  $P = 0.04$ ) and PP analysis (92.9% vs 88.3%,  $P = 0.06$ ). Zou *et al*[78] reported that 14-d DT had a higher *H. pylori* eradication rate than 10-d DT in ITT (89.7% vs 78.4%,  $P = 0.039$ ) and PP (92.9% vs 80.0%,  $P = 0.014$ ) analyses. The dosage and treatment duration of AMX were recommended as 3 g/d for 14 d to optimize the DT, respectively. As a first-line treatment, high-dose, high-frequency DT was effective and safe for treating *H. pylori* infections in elderly patients and those with multiple comorbidities[79]. Successful *H. pylori* eradication was achieved in 90.9% of patients, and adverse events (11.1%) were mainly mild.

### P-CABS

P-CABs are highly active drugs targeting  $H^+$ ,  $K^+$ -ATPase in the gastric acid secretion of parietal cells. The mechanism of action is different from that of PPIs. Conventional PPIs require 3-5 d to achieve maximal and steady-state gastric acid inhibition, whereas P-CABs increase the intragastric pH to nearly 7 within four hours[80].

As an alternative to a PPI, VPZ was efficacious when combined with DT. Furuta *et al*[81] compared the first-line *H. pylori* eradication rate of VA-DT (20 mg of VPZ twice daily and 500 mg of AMX three times daily for 7 d) and CAM-based triple therapy using VPZ. The eradication rates using the VPZ-based dual and triple therapies were 92.9% and 91.9% ( $P = 0.728$ ) in the ITT analysis and 94.4% and 92.7% ( $P = 0.715$ ) in the PP analysis, respectively. VA-DT showed a comparable *H. pylori* eradication rate without the need for CAM. In a randomized trial by Suzuki *et al*[82], the eradication rate of VA-DT (20 mg of VPZ and 750 mg of AMX twice daily for 7 d) was similar to that for VPZ-based triple therapy (84.5% vs 89.2%,  $P = 0.203$  in the ITT analysis; 87.1% vs 90.2%,  $P = 0.372$  in the PP analysis). In the subgroup analysis, the eradication rates of VA-DT in CAM-resistant *H. pylori* strains were significantly higher than those of VPZ-based triple therapy (92.3% vs 76.2%,  $P = 0.048$ ). When CAM-resistant *H. pylori* infection was treated with VPZ and AMX, CAM was not beneficial. Thus, extending the treatment duration of VA-DT to 14 d may be a promising alternative *H. pylori* treatment.

## CONCLUSION

At present, the availability of specific drugs and reagents differs among China, Japan, and South Korea. Bismuth is not licensed for use in Japan, whereas classic and modified BQTs are recommended in China and South Korea. Since 2015, VPZ was introduced to eradicate *H. pylori* in Japan. In contrast, few studies have focused on VPZ outside of Japan. In South Korea, molecular tests for CAM-resistant *H. pylori* are

commercially available. In Japan, a low rate of MDZ resistance results from the limited use of MDZ by the national health insurance system. Unlike in the other two countries, MDZ has not been approved as a component of first-line *H. pylori* regimens in Japan. Therefore, the strategy for the treatment of *H. pylori* infection might be selected based on the antibiotic resistance rate and medical policy in each country.

In China, adding bismuth to all *H. pylori* regimens is recommended as the empirical first-line treatment. Clinical trials with DT involving frequent administration of high-dose AMX have been widely performed in the Chinese population. In Japan, *H. pylori* treatment success has increased since VPZ was launched. Furthermore, VPZ may play an important role in DT by optimizing the intragastric environment for AMX action. PPIs may be replaced by VPZ due to its rapid and strong acid suppression. In South Korea, tailored eradication can be used as a first-line *H. pylori* treatment option based on the presence of a point mutation. The advantages of the *H. pylori* regimens used in China, Japan, and South Korea need to be combined in future research.

---

## FOOTNOTES

**Author contributions:** Cho JH designed research; Cho JH and Jin SY performed research and literature review; Cho JH wrote the paper; Cho JH and Jin SY contributed critical revision and editing.

**Supported by** the Soonchunhyang University Research Fund, No. 10210061.

**Conflict-of-interest statement:** The authors declare that they have no competing interests.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** South Korea

**ORCID number:** Jun-Hyung Cho 0000-0003-2075-2333; So-Young Jin 0000-0002-9900-8322.

**Corresponding Author's Membership in Professional Societies:** The Korean Society of Gastroenterology.

**S-Editor:** Yan JP

**L-Editor:** A

**P-Editor:** Yan JP

---

## REFERENCES

- 1 **Crowe SE.** *Helicobacter pylori* Infection. *N Engl J Med* 2019; **380**: 1158-1165 [PMID: 30893536 DOI: 10.1056/NEJMcp1710945]
- 2 **Wang F, Meng W, Wang B, Qiao L.** *Helicobacter pylori*-induced gastric inflammation and gastric cancer. *Cancer Lett* 2014; **345**: 196-202 [PMID: 23981572 DOI: 10.1016/j.canlet.2013.08.016]
- 3 **Hooi JKY, Lai WY, Ng WK, Suen MMY, Underwood FE, Tanyingoh D, Malfertheiner P, Graham DY, Wong VWS, Wu JCY, Chan FKL, Sung JJY, Kaplan GG, Ng SC.** Global Prevalence of *Helicobacter pylori* Infection: Systematic Review and Meta-Analysis. *Gastroenterology* 2017; **153**: 420-429 [PMID: 28456631 DOI: 10.1053/j.gastro.2017.04.022]
- 4 **Ren S, Cai P, Liu Y, Wang T, Zhang Y, Li Q, Gu Y, Wei L, Yan C, Jin G.** Prevalence of *Helicobacter pylori* infection in China: A systematic review and meta-analysis. *J Gastroenterol Hepatol* 2021 Dec 3 [PMID: 34862656 DOI: 10.1111/jgh.15751]
- 5 **Ueda J, Goshio M, Inui Y, Matsuda T, Sakakibara M, Mabe K, Nakajima S, Shimoyama T, Yasuda M, Kawai T, Murakami K, Kamada T, Mizuno M, Kikuchi S, Lin Y, Kato M.** Prevalence of *Helicobacter pylori* infection by birth year and geographic area in Japan. *Helicobacter* 2014; **19**: 105-110 [PMID: 24506211 DOI: 10.1111/hel.12110]
- 6 **Lee JH, Choi KD, Jung HY, Baik GH, Park JK, Kim SS, Kim BW, Hong SJ, Lim H, Shin CM, Lee SH, Jeon SW, Kim JH, Choi CW, Jung HK, Choi SC, Cho JW, Lee WS, Na SY, Sung JK, Song KH, Chung JW, Yun SC;** Korean College of Helicobacter and Upper Gastrointestinal Research. Seroprevalence of *Helicobacter pylori* in Korea: A multicenter, nationwide study conducted in 2015 and 2016. *Helicobacter* 2018; **23**: e12463 [PMID: 29345022 DOI: 10.1111/hel.12463]
- 7 **International Agency for Research on Cancer.** Globocan 2020. Cancer today. Available from: <https://gco.iarc.fr/today/home>
- 8 **Yang X, Zhang T, Zhang H, Sang S, Chen H, Zuo X.** Temporal trend of gastric cancer burden along with its risk factors in China from 1990 to 2019, and projections until 2030: comparison with Japan, South Korea, and Mongolia. *Biomark Res* 2021; **9**: 84 [PMID: 34784961 DOI: 10.1186/s40364-021-00340-6]
- 9 **Sugano K, Tack J, Kuipers EJ, Graham DY, El-Omar EM, Miura S, Haruma K, Asaka M, Uemura N, Malfertheiner P;**

- faculty members of Kyoto Global Consensus Conference. Kyoto global consensus report on *Helicobacter pylori* gastritis. *Gut* 2015; **64**: 1353-1367 [PMID: 26187502 DOI: 10.1136/gutjnl-2015-309252]
- 10 Asaka M, Kato M, Sakamoto N. Roadmap to eliminate gastric cancer with *Helicobacter pylori* eradication and consecutive surveillance in Japan. *J Gastroenterol* 2014; **49**: 1-8 [PMID: 24162382 DOI: 10.1007/s00535-013-0897-8]
  - 11 Liu WZ, Xie Y, Lu H, Cheng H, Zeng ZR, Zhou LY, Chen Y, Wang JB, Du YQ, Lu NH; Chinese Society of Gastroenterology, Chinese Study Group on *Helicobacter pylori* and Peptic Ulcer. Fifth Chinese National Consensus Report on the management of *Helicobacter pylori* infection. *Helicobacter* 2018; **23**: e12475 [PMID: 29512258 DOI: 10.1111/hel.12475]
  - 12 Thung I, Aramin H, Vavinskaya V, Gupta S, Park JY, Crowe SE, Valasek MA. Review article: the global emergence of *Helicobacter pylori* antibiotic resistance. *Aliment Pharmacol Ther* 2016; **43**: 514-533 [PMID: 26694080 DOI: 10.1111/apt.13497]
  - 13 Papastergiou V, Georgopoulos SD, Karatapanis S. Treatment of *Helicobacter pylori* infection: Past, present and future. *World J Gastrointest Pathophysiol* 2014; **5**: 392-399 [PMID: 25400982 DOI: 10.4291/wjgp.v5.i4.392]
  - 14 Shiotani A, Roy P, Lu H, Graham DY. *Helicobacter pylori* diagnosis and therapy in the era of antimicrobial stewardship. *Therap Adv Gastroenterol* 2021; **14**: 17562848211064080 [PMID: 34987609 DOI: 10.1177/17562848211064080]
  - 15 Huang CC, Tsai KW, Tsai TJ, Hsu PI. Update on the first-line treatment for *Helicobacter pylori* infection - a continuing challenge from an old enemy. *Biomark Res* 2017; **5**: 23 [PMID: 28702193 DOI: 10.1186/s40364-017-0103-x]
  - 16 Hu Y, Zhu Y, Lu NH. Primary Antibiotic Resistance of *Helicobacter pylori* in China. *Dig Dis Sci* 2017; **62**: 1146-1154 [PMID: 28315035 DOI: 10.1007/s10620-017-4536-8]
  - 17 Lee JH, Ahn JY, Choi KD, Jung HY, Kim JM, Baik GH, Kim BW, Park JC, Jung HK, Cho SJ, Shin CM, Choi YJ, Lee SH, Kim JH, Lee WS, Sung JK, Chung JW, Cheung DY, Lee H, Min YW, Kim JJ, Kim SY; Korean College of Helicobacter; Upper Gastrointestinal Research. Nationwide antibiotic resistance mapping of *Helicobacter pylori* in Korea: A prospective multicenter study. *Helicobacter* 2019; **24**: e12592 [PMID: 31111572 DOI: 10.1111/hel.12592]
  - 18 Okamura T, Suga T, Nagaya T, Arakura N, Matsumoto T, Nakayama Y, Tanaka E. Antimicrobial resistance and characteristics of eradication therapy of *Helicobacter pylori* in Japan: a multi-generational comparison. *Helicobacter* 2014; **19**: 214-220 [PMID: 24758533 DOI: 10.1111/hel.12124]
  - 19 Suzuki H, Mori H. World trends for *H. pylori* eradication therapy and gastric cancer prevention strategy by *H. pylori* test-and-treat. *J Gastroenterol* 2018; **53**: 354-361 [PMID: 29138921 DOI: 10.1007/s00535-017-1407-1]
  - 20 Murata M, Sugimoto M, Mizuno H, Kanno T, Satoh K. Clarithromycin Versus Metronidazole in First-Line *Helicobacter Pylori* Triple Eradication Therapy Based on Resistance to Antimicrobial Agents: Meta-Analysis. *J Clin Med* 2020; **9**: 543 [PMID: 32079208 DOI: 10.3390/jcm9020543]
  - 21 Kuo YT, Liou JM, El-Omar EM, Wu JY, Leow AHR, Goh KL, Das R, Lu H, Lin JT, Tu YK, Yamaoka Y, Wu MS; Asian Pacific Alliance on Helicobacter and Microbiota. Primary antibiotic resistance in *Helicobacter pylori* in the Asia-Pacific region: a systematic review and meta-analysis. *Lancet Gastroenterol Hepatol* 2017; **2**: 707-715 [PMID: 28781119 DOI: 10.1016/S2468-1253(17)30219-4]
  - 22 Lu H, Zhang W, Graham DY. Bismuth-containing quadruple therapy for *Helicobacter pylori*: lessons from China. *Eur J Gastroenterol Hepatol* 2013; **25**: 1134-1140 [PMID: 23778309 DOI: 10.1097/MEG.0b013e3283633b57]
  - 23 Hu Y, Zhu Y, Lu NH. Recent progress in *Helicobacter pylori* treatment. *Chin Med J (Engl)* 2020; **133**: 335-343 [PMID: 31929363 DOI: 10.1097/CM9.0000000000000618]
  - 24 Zhang W, Chen Q, Liang X, Liu W, Xiao S, Graham DY, Lu H. Bismuth, lansoprazole, amoxicillin and metronidazole or clarithromycin as first-line *Helicobacter pylori* therapy. *Gut* 2015; **64**: 1715-1720 [PMID: 26338726 DOI: 10.1136/gutjnl-2015-309900]
  - 25 Graham DY, Dore MP, Lu H. Understanding treatment guidelines with bismuth and non-bismuth quadruple *Helicobacter pylori* eradication therapies. *Expert Rev Anti Infect Ther* 2018; **16**: 679-687 [PMID: 30102559 DOI: 10.1080/14787210.2018.1511427]
  - 26 Chen Q, Zhang W, Fu Q, Liang X, Liu W, Xiao S, Lu H. Rescue Therapy for *Helicobacter pylori* Eradication: A Randomized Non-Inferiority Trial of Amoxicillin or Tetracycline in Bismuth Quadruple Therapy. *Am J Gastroenterol* 2016; **111**: 1736-1742 [PMID: 27670603 DOI: 10.1038/ajg.2016.443]
  - 27 Graham DY, Lu H. Furazolidone in *Helicobacter pylori* therapy: misunderstood and often unfairly maligned drug told in a story of French bread. *Saudi J Gastroenterol* 2012; **18**: 1-2 [PMID: 22249084 DOI: 10.4103/1319-3767.91724]
  - 28 Qiao C, Li Y, Liu J, Ji C, Qu J, Hu J, Ji R, Wan M, Lin B, Lin M, Qi Q, Zuo X. Clarithromycin versus furazolidone for naïve *Helicobacter pylori* infected patients in a high clarithromycin resistance area. *J Gastroenterol Hepatol* 2021; **36**: 2383-2388 [PMID: 33691344 DOI: 10.1111/jgh.15468]
  - 29 Liu DS, Wang YH, Zhu ZH, Zhang SH, Zhu X, Wan JH, Lu NH, Xie Y. Characteristics of *Helicobacter pylori* antibiotic resistance: data from four different populations. *Antimicrob Resist Infect Control* 2019; **8**: 192 [PMID: 31798838 DOI: 10.1186/s13756-019-0632-1]
  - 30 Liang X, Xu X, Zheng Q, Zhang W, Sun Q, Liu W, Xiao S, Lu H. Efficacy of bismuth-containing quadruple therapies for clarithromycin-, metronidazole-, and fluoroquinolone-resistant *Helicobacter pylori* infections in a prospective study. *Clin Gastroenterol Hepatol* 2013; **11**: 802-807.e1 [PMID: 23376004 DOI: 10.1016/j.cgh.2013.01.008]
  - 31 Lv ZF, Wang FC, Zheng HL, Wang B, Xie Y, Zhou XJ, Lv NH. Meta-analysis: is combination of tetracycline and amoxicillin suitable for *Helicobacter pylori* infection? *World J Gastroenterol* 2015; **21**: 2522-2533 [PMID: 25741163 DOI: 10.3748/wjg.v21.i8.2522]
  - 32 Zhang J, Han C, Lu WQ, Wang N, Wu SR, Wang YX, Ma JP, Wang JH, Hao C, Yuan DH, Liu N, Shi YQ. A randomized, multicenter and noninferiority study of amoxicillin plus berberine vs tetracycline plus furazolidone in quadruple therapy for *Helicobacter pylori* rescue treatment. *J Dig Dis* 2020; **21**: 256-263 [PMID: 32348007 DOI: 10.1111/1751-2980.12870]
  - 33 Lee YC, Dore MP, Graham DY. Diagnosis and Treatment of *Helicobacter pylori* Infection. *Annu Rev Med* 2022; **73**: 183-195 [PMID: 35084993 DOI: 10.1146/annurev-med-042220-020814]
  - 34 Deguchi H, Uda A, Murakami K. Current Status of *Helicobacter pylori* Diagnosis and Eradication Therapy in Japan Using a Nationwide Database. *Digestion* 2020; **101**: 441-449 [PMID: 31216549 DOI: 10.1159/000500819]

- 35 **Kato M**, Ota H, Okuda M, Kikuchi S, Satoh K, Shimoyama T, Suzuki H, Handa O, Furuta T, Mabe K, Murakami K, Sugiyama T, Uemura N, Takahashi S. Guidelines for the management of *Helicobacter pylori* infection in Japan: 2016 Revised Edition. *Helicobacter* 2019; **24**: e12597 [PMID: 31111585 DOI: 10.1111/hel.12597]
- 36 **Kiyotoki S**, Nishikawa J, Sakaida I. Efficacy of Vonoprazan for *Helicobacter pylori* Eradication. *Intern Med* 2020; **59**: 153-161 [PMID: 31243237 DOI: 10.2169/internalmedicine.2521-18]
- 37 **Akazawa Y**, Fukuda D, Fukuda Y. Vonoprazan-based therapy for *Helicobacter pylori* eradication: experience and clinical evidence. *Therap Adv Gastroenterol* 2016; **9**: 845-852 [PMID: 27803739 DOI: 10.1177/1756283X16668093]
- 38 **Suzuki S**, Gotoda T, Kusano C, Iwatsuka K, Moriyama M. The Efficacy and Tolerability of a Triple Therapy Containing a Potassium-Competitive Acid Blocker Compared With a 7-Day PPI-Based Low-Dose Clarithromycin Triple Therapy. *Am J Gastroenterol* 2016; **111**: 949-956 [PMID: 27185079 DOI: 10.1038/ajg.2016.182]
- 39 **Sue S**, Kuwashima H, Iwata Y, Oka H, Arima I, Fukuchi T, Sanga K, Inokuchi Y, Ishii Y, Kanno M, Terada M, Amano H, Naito M, Iwase S, Okazaki H, Komatsu K, Kokawa A, Kawana I, Morimoto M, Saito T, Kunishi Y, Ikeda A, Takahashi D, Miwa H, Sasaki T, Tamura T, Kondo M, Shibata W, Maeda S. The Superiority of Vonoprazan-based First-line Triple Therapy with Clarithromycin: A Prospective Multi-center Cohort Study on *Helicobacter pylori* Eradication. *Intern Med* 2017; **56**: 1277-1285 [PMID: 28566587 DOI: 10.2169/internalmedicine.56.7833]
- 40 **Sue S**, Ogushi M, Arima I, Kuwashima H, Nakao S, Naito M, Komatsu K, Kaneko H, Tamura T, Sasaki T, Kondo M, Shibata W, Maeda S. Vonoprazan- vs proton-pump inhibitor-based first-line 7-day triple therapy for clarithromycin-susceptible *Helicobacter pylori*: A multicenter, prospective, randomized trial. *Helicobacter* 2018; **23**: e12456 [PMID: 29271026 DOI: 10.1111/hel.12456]
- 41 **Li M**, Oshima T, Horikawa T, Tozawa K, Tomita T, Fukui H, Watari J, Miwa H. Systematic review with meta-analysis: Vonoprazan, a potent acid blocker, is superior to proton-pump inhibitors for eradication of clarithromycin-resistant strains of *Helicobacter pylori*. *Helicobacter* 2018; **23**: e12495 [PMID: 29873436 DOI: 10.1111/hel.12495]
- 42 **Savoldi A**, Carrara E, Graham DY, Conti M, Tacconelli E. Prevalence of Antibiotic Resistance in *Helicobacter pylori*: A Systematic Review and Meta-analysis in World Health Organization Regions. *Gastroenterology* 2018; **155**: 1372-1382 [PMID: 29990487 DOI: 10.1053/j.gastro.2018.07.007]
- 43 **Nishizawa T**, Maekawa T, Watanabe N, Harada N, Hosoda Y, Yoshinaga M, Yoshio T, Ohta H, Inoue S, Toyokawa T, Yamashita H, Saito H, Kuwai T, Katayama S, Masuda E, Miyabayashi H, Kimura T, Nishizawa Y, Takahashi M, Suzuki H. Clarithromycin Versus Metronidazole as First-line *Helicobacter pylori* Eradication: A Multicenter, Prospective, Randomized Controlled Study in Japan. *J Clin Gastroenterol* 2015; **49**: 468-471 [PMID: 24921211 DOI: 10.1097/MCG.000000000000165]
- 44 **Adachi T**, Matsui S, Watanabe T, Okamoto K, Okamoto A, Kono M, Yamada M, Nagai T, Komeda Y, Minaga K, Kamata K, Yamao K, Takenaka M, Asakuma Y, Sakurai T, Nishida N, Kashida H, Kudo M. Comparative Study of Clarithromycin-versus Metronidazole-Based Triple Therapy as First-Line Eradication for *Helicobacter pylori*. *Oncology* 2017; **93** Suppl 1: 15-19 [PMID: 29258111 DOI: 10.1159/000481224]
- 45 **Shinozaki S**, Kobayashi Y, Osawa H, Sakamoto H, Hayashi Y, Lefor AK, Yamamoto H. Effectiveness and Safety of Vonoprazan versus Proton Pump Inhibitors for Second-Line *Helicobacter pylori* Eradication Therapy: Systematic Review and Meta-Analysis. *Digestion* 2021; **102**: 319-325 [PMID: 31914442 DOI: 10.1159/000504939]
- 46 **Murakami K**, Furuta T, Ando T, Nakajima T, Inui Y, Oshima T, Tomita T, Mabe K, Sasaki M, Suganuma T, Nomura H, Satoh K, Hori S, Inoue S, Tomokane T, Kudo M, Inaba T, Take S, Ohkusa T, Yamamoto S, Mizuno S, Kamoshida T, Amagai K, Iwamoto J, Miwa J, Kodama M, Okimoto T, Kato M, Asaka M; Japan GAST Study Group. Multi-center randomized controlled study to establish the standard third-line regimen for *Helicobacter pylori* eradication in Japan. *J Gastroenterol* 2013; **48**: 1128-1135 [PMID: 23307042 DOI: 10.1007/s00535-012-0731-8]
- 47 **Sugimoto M**, Sahara S, Ichikawa H, Kagami T, Uotani T, Furuta T. High *Helicobacter pylori* cure rate with sitafloxacin-based triple therapy. *Aliment Pharmacol Ther* 2015; **42**: 477-483 [PMID: 26075959 DOI: 10.1111/apt.13280]
- 48 **Nishizawa T**, Munkjargal M, Ebinuma H, Toyoshima O, Suzuki H. Sitafloxacin for Third-Line *Helicobacter pylori* Eradication: A Systematic Review. *J Clin Med* 2021; **10**: 2722 [PMID: 34202993 DOI: 10.3390/jcm10122722]
- 49 **Jung HK**, Kang SJ, Lee YC, Yang HJ, Park SY, Shin CM, Kim SE, Lim HC, Kim JH, Nam SY, Shin WG, Park JM, Choi IJ, Kim JG, Choi M; Korean College of *Helicobacter* and Upper Gastrointestinal Research. Evidence-Based Guidelines for the Treatment of *Helicobacter pylori* Infection in Korea 2020. *Gut Liver* 2021; **15**: 168-195 [PMID: 33468712 DOI: 10.5009/gnl20288]
- 50 **Park SM**, Kim JS, Kim BW, Ji JS, Choi H. Randomized clinical trial comparing 10- or 14-day sequential therapy and 10- or 14-day concomitant therapy for the first line empirical treatment of *Helicobacter pylori* infection. *J Gastroenterol Hepatol* 2017; **32**: 589-594 [PMID: 27505301 DOI: 10.1111/jgh.13510]
- 51 **Kim BJ**, Lee H, Lee YC, Jeon SW, Kim GH, Kim HS, Sung JK, Lee DH, Kim HU, Park MI, Choi IJ, Yoon SM, Kim SW, Baik GH, Lee JY, Kim JI, Kim SG, Kim J, Lee J, Kim JG, Kim JJ; Korean College of *Helicobacter* Upper Gastrointestinal Research. Ten-Day Concomitant, 10-Day Sequential, and 7-Day Triple Therapy as First-Line Treatment for *Helicobacter pylori* Infection: A Nationwide Randomized Trial in Korea. *Gut Liver* 2019; **13**: 531-540 [PMID: 31505907 DOI: 10.5009/gnl19136]
- 52 **Fallone CA**, Moss SF, Malfertheiner P. Reconciliation of Recent *Helicobacter pylori* Treatment Guidelines in a Time of Increasing Resistance to Antibiotics. *Gastroenterology* 2019; **157**: 44-53 [PMID: 30998990 DOI: 10.1053/j.gastro.2019.04.011]
- 53 **Lee HJ**, Kim JI, Lee JS, Jun EJ, Oh JH, Cheung DY, Chung WC, Kim BW, Kim SS. Concomitant therapy achieved the best eradication rate for *Helicobacter pylori* among various treatment strategies. *World J Gastroenterol* 2015; **21**: 351-359 [PMID: 25574111 DOI: 10.3748/wjg.v21.i1.351]
- 54 **Lee BE**, Kim JS, Kim BW, Kim JH, Kim JI, Chung JW, Jeon SW, Lee JH, Kim N, Lee JY, Seo SY, Park SY, Kim SE, Joo MK, Song HJ, Kim KB, Bang CS, Kim HJ. Consistency of *Helicobacter pylori* eradication rates of first-line concomitant and sequential therapies in Korea: A nationwide multicenter retrospective study for the last 10 years. *Helicobacter* 2021; **26**: e12780 [PMID: 33596342 DOI: 10.1111/hel.12780]
- 55 **White B**, Winte M, DeSipio J, Phadtare S. Clinical Factors Implicated in Antibiotic Resistance in *Helicobacter pylori*

- Patients. *Microorganisms* 2022; **10**: 322 [PMID: 35208776 DOI: 10.3390/microorganisms10020322]
- 56 **Graham DY**, Lee SY. How to Effectively Use Bismuth Quadruple Therapy: The Good, the Bad, and the Ugly. *Gastroenterol Clin North Am* 2015; **44**: 537-563 [PMID: 26314667 DOI: 10.1016/j.gtc.2015.05.003]
- 57 **Kim BJ**, Yang CH, Song HJ, Jeon SW, Kim GH, Kim HS, Kim TH, Shim KN, Chung IK, Park MI, Choi IJ, Kim JH, Kim BW, Baik GH, Han SW, Seo HE, Jung WT, Hwan Oh J, Kim SG, Lee JH, Park SK, Park BJ, Yang BR, Lee J, Kim JG. Online registry for nationwide database of *Helicobacter pylori* eradication in Korea: Correlation of antibiotic use density with eradication success. *Helicobacter* 2019; **24**: e12646 [PMID: 31368629 DOI: 10.1111/hel.12646]
- 58 **Chen H**, Dang Y, Zhou X, Liu B, Liu S, Zhang G. Tailored Therapy Versus Empiric Chosen Treatment for *Helicobacter pylori* Eradication: A Meta-Analysis. *Medicine (Baltimore)* 2016; **95**: e2750 [PMID: 26886617 DOI: 10.1097/MD.0000000000002750]
- 59 **Gisbert JP**. Empirical or susceptibility-guided treatment for *Helicobacter pylori* infection? *Therap Adv Gastroenterol* 2020; **13**: 1756284820968736 [PMID: 33240392 DOI: 10.1177/1756284820968736]
- 60 **Nishizawa T**, Suzuki H. Mechanisms of *Helicobacter pylori* antibiotic resistance and molecular testing. *Front Mol Biosci* 2014; **1**: 19 [PMID: 25988160 DOI: 10.3389/fmolb.2014.00019]
- 61 **Kim SY**, Park JM, Lim CH, Lee HA, Shin GY, Choe Y, Cho YK, Choi MG. Types of 23S Ribosomal RNA Point Mutations and Therapeutic Outcomes for *Helicobacter pylori*. *Gut Liver* 2021; **15**: 528-536 [PMID: 33376228 DOI: 10.5009/gnl20225]
- 62 **Cho JH**, Jin SY. Optimized diagnosis of *Helicobacter pylori* and tailored eradication therapy for preventing gastric cancer: a proposal for SHAKE strategy. *Expert Rev Gastroenterol Hepatol* 2020; **14**: 553-564 [PMID: 32410515 DOI: 10.1080/17474124.2020.1770594]
- 63 **Lee JW**, Kim SJ, Choi CW, Kim HJ, Kang DH, Kim HW, Park SB, Nam HS, Ryu DG. Seven-day triple therapy is sufficient to eradicate infection caused by *Helicobacter pylori* without 23S rRNA point mutation. *Medicine (Baltimore)* 2021; **100**: e26133 [PMID: 34032763 DOI: 10.1097/MD.0000000000026133]
- 64 **Seo SI**, Do BJ, Kang JG, Kim HS, Jang MK, Kim HY, Shin WG. *Helicobacter pylori* Eradication According to Sequencing-Based 23S Ribosomal RNA Point Mutation Associated with Clarithromycin Resistance. *J Clin Med* 2019; **9**: 54 [PMID: 31881688 DOI: 10.3390/jcm9010054]
- 65 **Kim JL**, Cho SJ, Chung SJ, Lee A, Choi J, Chung H, Kim SG. Empiric Versus Clarithromycin Resistance-Guided Therapy for *Helicobacter pylori* Based on Polymerase Chain Reaction Results in Patients With Gastric Neoplasms or Gastric Mucosa-Associated Lymphoid Tissue Lymphoma: A Randomized Controlled Trial. *Clin Transl Gastroenterol* 2020; **11**: e00194 [PMID: 33094958 DOI: 10.14309/ctg.0000000000000194]
- 66 **Choi YI**, Chung JW, Park DK, Kim KO, Kwon KA, Kim YJ, Seo JY. Tailored eradication vs empirical bismuth-containing quadruple therapy for first-line *Helicobacter pylori* eradication: A comparative, open trial. *World J Gastroenterol* 2019; **25**: 6743-6751 [PMID: 31857776 DOI: 10.3748/wjg.v25.i46.6743]
- 67 **Dyar OJ**, Huttner B, Schouten J, Pulcini C; ESGAP (ESCMID Study Group for Antimicrobial stewardship). What is antimicrobial stewardship? *Clin Microbiol Infect* 2017; **23**: 793-798 [PMID: 28882725 DOI: 10.1016/j.cmi.2017.08.026]
- 68 **Graham DY**, Dore MP. *Helicobacter pylori* therapy: a paradigm shift. *Expert Rev Anti Infect Ther* 2016; **14**: 577-585 [PMID: 27077447 DOI: 10.1080/14787210.2016.1178065]
- 69 **Marcus EA**, Inatomi N, Nagami GT, Sachs G, Scott DR. The effects of varying acidity on *Helicobacter pylori* growth and the bactericidal efficacy of ampicillin. *Aliment Pharmacol Ther* 2012; **36**: 972-979 [PMID: 23009227 DOI: 10.1111/apt.12059]
- 70 **Furuta T**, Graham DY. Pharmacologic aspects of eradication therapy for *Helicobacter pylori* Infection. *Gastroenterol Clin North Am* 2010; **39**: 465-480 [PMID: 20951912 DOI: 10.1016/j.gtc.2010.08.007]
- 71 **Takahashi N**, Take Y. Tegoprazan, a Novel Potassium-Competitive Acid Blocker to Control Gastric Acid Secretion and Motility. *J Pharmacol Exp Ther* 2018; **364**: 275-286 [PMID: 29180359 DOI: 10.1124/jpet.117.244202]
- 72 **Oshima T**, Miwa H. Potent Potassium-competitive Acid Blockers: A New Era for the Treatment of Acid-related Diseases. *J Neurogastroenterol Motil* 2018; **24**: 334-344 [PMID: 29739175 DOI: 10.5056/jnm18029]
- 73 **Yun J**, Wu Z, Qi G, Han T, Zhang D. The high-dose amoxicillin-proton pump inhibitor dual therapy in eradication of *Helicobacter pylori* infection. *Expert Rev Gastroenterol Hepatol* 2021; **15**: 149-157 [PMID: 32960107 DOI: 10.1080/17474124.2021.1826306]
- 74 **Graham DY**, Lu H, Shiotani A. Failure of optimized dual proton pump inhibitor amoxicillin therapy: What now? *Saudi J Gastroenterol* 2017; **23**: 265-267 [PMID: 28937019 DOI: 10.4103/sjg.SJG\_292\_17]
- 75 **Yang JC**, Lin CJ, Wang HL, Chen JD, Kao JY, Shun CT, Lu CW, Lin BR, Shieh MJ, Chang MC, Chang YT, Wei SC, Lin LC, Yeh WC, Kuo JS, Tung CC, Leong YL, Wang TH, Wong JM. High-dose dual therapy is superior to standard first-line or rescue therapy for *Helicobacter pylori* infection. *Clin Gastroenterol Hepatol* 2015; **13**: 895-905 [PMID: 25460556 DOI: 10.1016/j.cgh.2014.10.036]
- 76 **Gao CP**, Zhang D, Zhang T, Wang JX, Han SX, Graham DY, Lu H. PPI-amoxicillin dual therapy for *Helicobacter pylori* infection: An update based on a systematic review and meta-analysis. *Helicobacter* 2020; **25**: e12692 [PMID: 32314468 DOI: 10.1111/hel.12692]
- 77 **Li C**, Shi Y, Suo B, Tian X, Zhou L, Song Z. PPI-amoxicillin dual therapy four times daily is superior to guidelines recommended regimens in the *Helicobacter pylori* eradication therapy within Asia: A systematic review and meta-analysis. *Helicobacter* 2021; **26**: e12816 [PMID: 34002433 DOI: 10.1111/hel.12816]
- 78 **Zou PY**, Hu J, Zhao JT, Zhao Z, Mei H, Yang J, Zhu YJ, Zhang Y, Lan CH. 10-Day and 14-day high-dose dual therapy for the treatment of *Helicobacter pylori*: A propensity score matching analysis. *Helicobacter* 2021; **26**: e12833 [PMID: 34318966 DOI: 10.1111/hel.12833]
- 79 **Gao W**, Ye H, Deng X, Wang C, Xu Y, Li Y, Zhang X, Cheng H. Rabeprazole-amoxicillin dual therapy as first-line treatment for *H pylori* eradication in special patients: A retrospective, real-life study. *Helicobacter* 2020; **25**: e12717 [PMID: 32548932 DOI: 10.1111/hel.12717]
- 80 **Miftahussurur M**, Pratama Putra B, Yamaoka Y. The Potential Benefits of Vonoprazan as *Helicobacter pylori* Infection Therapy. *Pharmaceuticals (Basel)* 2020; **13**: 276 [PMID: 32998241 DOI: 10.3390/ph13100276]

- 81 **Furuta T**, Yamade M, Kagami T, Uotani T, Suzuki T, Higuchi T, Tani S, Hamaya Y, Iwaizumi M, Miyajima H, Umemura K, Osawa S, Sugimoto K. Dual Therapy with Vonoprazan and Amoxicillin Is as Effective as Triple Therapy with Vonoprazan, Amoxicillin and Clarithromycin for Eradication of *Helicobacter pylori*. *Digestion* 2020; **101**: 743-751 [PMID: [31434101](#) DOI: [10.1159/000502287](#)]
- 82 **Suzuki S**, Gotoda T, Kusano C, Ikehara H, Ichijima R, Ohyauchi M, Ito H, Kawamura M, Ogata Y, Ohtaka M, Nakahara M, Kawabe K. Seven-day vonoprazan and low-dose amoxicillin dual therapy as first-line *Helicobacter pylori* treatment: a multicentre randomised trial in Japan. *Gut* 2020; **69**: 1019-1026 [PMID: [31915235](#) DOI: [10.1136/gutjnl-2019-319954](#)]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA  
**Telephone:** +1-925-3991568  
**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
**Help Desk:** <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>

