

World Journal of *Gastroenterology*

World J Gastroenterol 2022 July 7; 28(25): 2782-3007



REVIEW

- 2782 Inflammation, microbiome and colorectal cancer disparity in African-Americans: Are there bugs in the genetics?
Ahmad S, Ashktorab H, Brim H, Housseau F
- 2802 Altered gut microbiota patterns in COVID-19: Markers for inflammation and disease severity
Chakraborty C, Sharma AR, Bhattacharya M, Dhama K, Lee SS
- 2823 Long noncoding RNAs in hepatitis B virus replication and oncogenesis
Li HC, Yang CH, Lo SY

MINIREVIEWS

- 2843 Characteristics of inflammatory bowel diseases in patients with concurrent immune-mediated inflammatory diseases
Akiyama S, Fukuda S, Steinberg JM, Suzuki H, Tsuchiya K
- 2854 Correlation of molecular alterations with pathological features in hepatocellular carcinoma: Literature review and experience of an Italian center
Maloberti T, De Leo A, Sanza V, Gruppioni E, Altamari A, Riefolo M, Visani M, Malvi D, D'Errico A, Tallini G, Vasuri F, de Biase D
- 2867 Micelles as potential drug delivery systems for colorectal cancer treatment
Fatfat Z, Fatfat M, Gali-Muhtasib H
- 2881 Incretin based therapy and pancreatic cancer: Realising the reality
Suryadevara V, Roy A, Sahoo J, Kamalanathan S, Naik D, Mohan P, Kalayarasan R
- 2890 Non-alcoholic fatty liver disease and the impact of genetic, epigenetic and environmental factors in the offspring
Wajsbrot NB, Leite NC, Salles GF, Villela-Nogueira CA
- 2900 Role of transcribed ultraconserved regions in gastric cancer and therapeutic perspectives
Gao SS, Zhang ZK, Wang XB, Ma Y, Yin GQ, Guo XB
- 2910 Multiple roles for cholinergic signaling in pancreatic diseases
Yang JM, Yang XY, Wan JH

ORIGINAL ARTICLE**Basic Study**

- 2920 Fecal gene detection based on next generation sequencing for colorectal cancer diagnosis
He SY, Li YC, Wang Y, Peng HL, Zhou CL, Zhang CM, Chen SL, Yin JF, Lin M

- 2937 Mechanism and therapeutic strategy of hepatic *TM6SF2*-deficient non-alcoholic fatty liver diseases *via in vivo* and *in vitro* experiments

Li ZY, Wu G, Qiu C, Zhou ZJ, Wang YP, Song GH, Xiao C, Zhang X, Deng GL, Wang RT, Yang YL, Wang XL

- 2955 Upregulated adenosine 2A receptor accelerates post-infectious irritable bowel syndrome by promoting CD4+ T cells' T helper 17 polarization

Dong LW, Ma ZC, Fu J, Huang BL, Liu FJ, Sun D, Lan C

Retrospective Study

- 2968 Four-year experience with more than 1000 cases of total laparoscopic liver resection in a single center

Lan X, Zhang HL, Zhang H, Peng YF, Liu F, Li B, Wei YG

SCIENTOMETRICS

- 2981 Mapping the global research landscape on nutrition and the gut microbiota: Visualization and bibliometric analysis

Zyoud SH, Shakhshir M, Abushanab AS, Al-Jabi SW, Koni A, Shahwan M, Jairoun AA, Abu Taha A

CASE REPORT

- 2994 Early gastric cancer presenting as a typical submucosal tumor cured by endoscopic submucosal dissection: A case report

Cho JH, Lee SH

LETTER TO THE EDITOR

- 3001 Acupuncture and moxibustion for treatment of Crohn's disease: A brief review

Xie J, Huang Y, Wu HG, Li J

CORRECTION

- 3004 Correction to "Aberrant methylation of secreted protein acidic and rich in cysteine gene and its significance in gastric cancer"

Shao S, Zhou NM, Dai DQ

- 3006 Correction to "Gut microbiota dysbiosis in Chinese children with type 1 diabetes mellitus: An observational study"

Liu X, Cheng YW, Shao L, Sun SH, Wu J, Song QH, Zou HS, Ling ZX

ABOUT COVER

Editorial Board Member of *World Journal of Gastroenterology*, Hideyuki Chiba, MD, PhD, Director, Department of Gastroenterology, Omori Red Cross Hospital, 4-30-1, Chuo, Ota-Ku, Tokyo 143-8527, Japan. h.chiba04@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Gastroenterology (WJG, World J Gastroenterol)* is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. *WJG* mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The *WJG* is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central, and Scopus. The 2021 edition of Journal Citation Report® cites the 2020 impact factor (IF) for *WJG* as 5.742; Journal Citation Indicator: 0.79; IF without journal self cites: 5.590; 5-year IF: 5.044; Ranking: 28 among 92 journals in gastroenterology and hepatology; and Quartile category: Q2. The *WJG*'s CiteScore for 2020 is 6.9 and Scopus CiteScore rank 2020: Gastroenterology is 19/136.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*; **Production Department Director:** *Xiang Li*; **Editorial Office Director:** *Jia-Ru Fan*.

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Andrzej S Tarnawski

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

PUBLICATION DATE

July 7, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Multiple roles for cholinergic signaling in pancreatic diseases

Jun-Min Yang, Xiao-Yu Yang, Jian-Hua Wan

Specialty type: Gastroenterology and hepatology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B, B, B
Grade C (Good): C, C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Aktekin A, Turkey; Kitamura K, Japan; Luyer MDP, Netherlands; Matsuo Y, Japan; Susak YM, Ukraine

Received: February 20, 2022

Peer-review started: February 20, 2022

First decision: May 9, 2022

Revised: May 18, 2022

Accepted: June 13, 2022

Article in press: June 13, 2022

Published online: July 7, 2022



Jun-Min Yang, Xiao-Yu Yang, Jian-Hua Wan, Department of Gastroenterology, The First Affiliated Hospital of Nanchang University, Nanchang 330006, Jiangxi Province, China

Corresponding author: Jian-Hua Wan, MD, PhD, Doctor, Department of Gastroenterology, The First Affiliated Hospital of Nanchang University, No. 17 Yongwaizheng Street, Nanchang 330006, Jiangxi Province, China. 18942337437@163.com

Abstract

Cholinergic nerves are widely distributed throughout the human body and participate in various physiological activities, including sensory, motor, and visceral activities, through cholinergic signaling. Cholinergic signaling plays an important role in pancreatic exocrine secretion. A large number of studies have found that cholinergic signaling overstimulates pancreatic acinar cells through muscarinic receptors, participates in the onset of pancreatic diseases such as acute pancreatitis and chronic pancreatitis, and can also inhibit the progression of pancreatic cancer. However, cholinergic signaling plays a role in reducing pain and inflammation through nicotinic receptors, but enhances the proliferation and invasion of pancreatic tumor cells. This review focuses on the progression of cholinergic signaling and pancreatic diseases in recent years and reveals the role of cholinergic signaling in pancreatic diseases.

Key Words: Acetylcholine; Muscarinic receptors; Nicotinic receptors; Pancreatic exocrine; Pancreatitis; Pancreatic cancer

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The pancreas is a nerve-rich organ that lies behind the peritoneum and is surrounded by many nerve plexuses. Studies have found that cholinergic signaling is involved in the physiological function of the pancreas and the pathological process of pancreatic diseases due to its action on different receptors. Perhaps starting with cholinergic receptors could uncover potential therapeutic targets for pancreatic diseases.

Citation: Yang JM, Yang XY, Wan JH. Multiple roles for cholinergic signaling in pancreatic diseases. *World J Gastroenterol* 2022; 28(25): 2910-2919

URL: <https://www.wjgnet.com/1007-9327/full/v28/i25/2910.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v28.i25.2910>

INTRODUCTION

Acetylcholine is a neurotransmitter released by all cholinergic neurons that plays an important role in the peripheral and central nervous systems. The vagus nerve is the longest nerve in the human body and provides innervation to most organs, especially the organs of the digestive system. Cholinergic signaling released by vagus nerve activation mainly acts on cholinergic receptors. According to pharmacological properties, cholinergic receptors are divided into muscarinic receptors (M receptors) and nicotinic receptors (N receptors). Among them, M receptors are widely distributed in smooth muscles and glands and participate in the secretion of glands. M receptors are divided into 5 subtypes of M1-M5, among which M1, M3, and M5 receptors are coupled to Gq protein and M2 and M4 receptors are coupled to Gi protein[1,2]. Each of these five subtypes of M receptors has a unique distribution pattern and is expressed in many areas of the central nervous system and peripheral tissues[1,3-5]. N receptors are ligand-gated ion channels, a pentamer made up of 5 identical homologous subunits (including α 1-10, β 1- β 5, γ , δ , and ϵ) and retain the potential to be activated by the appropriate agonist[6-8].

The pancreas is an important digestive organ of the human body and plays an important role in the digestion and absorption of food. Pancreatic diseases include acute pancreatitis (AP), chronic pancreatitis (CP), and pancreatic cancer (PCa). Recently, the incidence of pancreatic disease has been increasing year by year, posing a serious threat to human life and health[9-11]. Recent studies have found that cholinergic neuromodulation is involved in the occurrence and development of pancreatic diseases[12-14]. This article reviews the relationship between the three main types of pancreatic diseases (AP, CP, PCa) and cholinergic signaling.

CHOLINERGIC SIGNALING AND PANCREATIC EXOCRINE FUNCTION

The exocrine system of the pancreas includes three different stages, namely, the cephalic stage, the stomach stage, and the intestinal stage[15]. The total amount of daily exocrine activity of the human pancreas is 1-2 L, and approximately 20% occurs in the cephalic stage, which is under the control of the vagus nerve[16-18]. Studies on humans and animals have shown that the cholinergic system regulates pancreatic exocrine secretion through the vagus nerve reflex. These reflexes originate in the dorsal motor nucleus of the vagus nerve in the medulla oblongata. The cranial preganglionic nerve fibers exit the vagus nerve and terminate at the intrapancreatic ganglia to form synapses through cholinergic preganglionic fibers[19,20].

The intrapancreatic ganglion are the integration center of pancreatic exocrine secretion, and terminal axons from these ganglia innervate approximately every acinar artery. Although preganglionic neurotransmission is mediated by acetylcholine through nicotinic and muscarinic receptors, postpancreatic innervation can be mediated by a variety of neurotransmitters, including acetylcholine, which acts on muscarinic receptors in pancreatic acinar cells[21,22]. Cholinergic agonists produce a pancreatic secretory response similar to that of cephalic stimulation, whereas cholinergic antagonists or resection of the vagus nerve can block the cephalic response. These results show that the acetylcholine released by the efferents of the vagus nerve is the primary mechanism by which sensory input leads to the regulation of pancreatic exocrine secretion[23-26]. The cholinergic system also plays an important role in the gastric and intestinal stages of pancreatic secretion regulation[27,28], which are regulated by nerves and bodily fluids. In the intestinal phase, exocrine secretion of the pancreas is mainly regulated by cholecystokinin (CCK) and other gastrointestinal hormones. However, the cholinergic system also plays a role in the secretion of human pancreatic enzymes stimulated by CCK[21,29]. The postprandial physiological dose of CCK mainly acts on the afferent pathway of the vagus nerve in the gastric and duodenal mucosa and stimulates pancreatic exocrine secretion through the cholinergic efferent nerve [19].

Studies have found that human pancreatic acinar cells preferentially express the M3 receptor[30] and are highly expressed in acinar cells[31]. Acetylcholine acts on the M3 receptor and couples to the Gq protein to activate phospholipase C and promote the release of intracellular calcium ions by initiating the phosphatidyl C-inositol triphosphate cascade and promoting the secretion of pancreatic acinar cells [2,32,33]. According to drug blocking studies, the M3 receptor is a muscarinic receptor that stimulates exocrine secretion from the pancreas[34-36]. However, some studies have shown that the M1 receptor can also control pancreatic exocrine secretion[37,38]. The inhibitory effect of drugs blocking the M1 receptor on the secretion of amylase in isolated pancreatic acinar cells was significantly greater than that of drugs blocking the M3 receptor[39]. Carbachol-induced amylase secretion was significantly impaired in the isolated acinars with M1 and M3 muscarinic receptor single knockout (KO) mice, and amylase secretion was eliminated in the acinar preparation of M1 and M3 muscarinic receptor double KO mice. Therefore, it is proposed that cholinergic signaling stimulates the secretion of pancreatic amylase and is mediated by the combination of M1 and M3 receptors[40].

CHOLINERGIC SIGNALING AND AP

AP is a common pancreatic disease in clinical practice. AP has a rapid onset and progress, and high morbidity and mortality worldwide[11]. There are many causes of AP, such as biliary, hypertriglyceridemia, and chronic alcohol consumption, among which biliary is the most common factor[41]. However, the specific mechanism underlying the cause of pancreatitis is not clear. When the human body is exposed to cholinergic agonists, such as scorpion stings or organophosphorus pesticide poisoning, some patients will develop AP. This is direct evidence that cholinergic signaling stimulation is associated with the occurrence of human AP[42-45]. Organophosphorus pesticides can inhibit cholinesterase activity and cause a large amount of acetylcholine to accumulate in nerve endings. Similarly, scorpion toxin is a neurotoxic protein that can cause AP by activating the nerve pathway that releases acetylcholine[46,47].

Pancreatic duct ligation in the rat model is often used to simulate clinical AP caused by gallstone obstruction[48], but the severity of experimental pancreatitis is low except when used in the possum model. However, some researchers have found in a pancreatic duct ligation rat model that cholinergic signaling stimulation can aggravate AP inflammation, suggesting that cholinergic signaling may be involved in the pathogenesis of AP caused by cholangiopancreatic duct obstruction[49]. This may be due to an increase in M3 receptor expression induced by rat pancreatic duct ligation, which amplifies the overstimulation of cholinergic signaling on acinar cells and aggravates the intracellular stress response[50]. Before the first clinical onset of acute alcoholic pancreatitis, patients usually have a history of alcohol abuse for many years. Chronic alcohol intake can affect the exocrine regulation of the pancreas by interfering with the cholinergic and trypsin pathways[51]. Long-term use of ethanol feeding can significantly reduce pancreatic acetylcholinesterase activity in rats, while the expression level of pancreatic cholinergic M receptors has not changed, which increases the level and duration of acetylcholine in the pancreas and leads to excessive cholinergic signaling stimulation and damage to pancreatic acinar cells[52,53]. Ethanol-treated pancreatic acinar cells can aggravate the pancreatic injury response caused by the cholinergic agonist carbachol, and its effect may be mediated by protein kinase C downstream signaling of cholinergic receptors[54]. The cholinergic receptor antagonist atropine can improve pancreatitis induced by the combination of alcohol and cerulein suggesting that the cholinergic signaling pathway is involved in the pathogenesis of pancreatitis[52]. Therefore, acetylcholine may play a key role in the pathogenesis of acute alcoholic pancreatitis[12].

The M3 receptor is highly expressed in human pancreatic acinar cells[31]. Wan *et al*[55] used chemical genetic technology, in which designer receptors are exclusively activated by designer drugs, to express mutant M3 receptors in mouse acinar cells, causing them to lose their response to acetylcholine but can be activated by the specific drug clozapine-N-oxide (CNO). CNO can induce AP in mutant M3 receptor mice and cause more extensive acinar cell necrosis and inflammation. In addition, the use of M3 receptor antagonists can improve the severity of AP induced by cerulein in wild-type mice (Figure 1) [55].

These results indicate that the activation of the M3 receptor by cholinergic nerve terminals releasing acetylcholine may be one of the pathogenesises of AP. Muscarinic receptor agonists stimulate the activation of trypsinogen and nuclear factor-kappaB, which are two key signaling pathways in the pathogenesis of pancreatitis[56]. Reducing inflammation has always been a major goal in the treatment of AP, and many anti-inflammatory drugs have shown beneficial responses in experimental pancreatitis [57]. However, due to the lack of an in-depth understanding of its pathogenesis, there is currently no effective prevention and treatment strategy[58]. Through in-depth research on the cholinergic signaling pathway, it may be possible to block cholinergic signaling early to treat AP, especially in the prevention of pancreatitis after endoscopic retrograde cholangiopancreatography.

CHOLINERGIC SIGNALING AND CP

CP is characterized by chronic inflammation and fibrosis of the pancreas caused by multiple factors. The incidence and prevalence of CP are increasing each year, but there is no current specific treatment[10]. The most common causes of CP are excessive drinking, smoking or genetic mutations[59]. Alcohol is considered to be one of the main risk factors for CP. A total of 40%-70% of CP patients become sick due to excessive alcohol consumption[10]. At the same time, excessive alcohol consumption increases the risk of PCa in individuals[60].

Recurrent episodes of AP have been associated with the progression of CP, which is more common in patients with alcoholism. It has been reported that a certain extent of chronic pancreatic damage was already present at the time of AP episodes[41]. The dose-response relationship between alcohol consumption for AP and CP is linear in males[61]. Alcohol-induced pancreatitis may be caused by the alcohol-induced increased viscosity of pancreatic secretions, which blocks the pancreatic duct, and by premature activation of trypsinogen in acinar cells[62].

The hypertonicity of intrapancreatic cholinergic neurons caused by chronic alcoholism may be involved in the pathogenesis of CP[13]. In CP patients, ethanol can cause excessive sensitivity of the




Figure 1 M3 receptor activation on pancreatic acinar cells causes acute and chronic pancreatitis. Ach: Acetylcholine; CCK: Cholecystokinin.

pancreatic parasympathetic nerve pathways after a meal[63]. This may be due to chronic alcoholism interrupting the autonomic nerve suppression reflex, leading to the decentralization of the intermediate autonomic nerve (located in the gastric antrum and duodenum) and the intrapancreatic ganglia. Due to an increase in the activity of these autonomic nerves, cholinergic signaling in the pancreas increases, resulting in excessive protein secretion and obstruction of pancreatic juice flow[64,65]. Histopathological analysis of pancreatic tissue samples from patients with CP (including alcoholic pancreatitis) demonstrated that the density of cholinergic fibers in the pancreas of patients with CP was slightly increased compared with normal pancreatic tissue samples[66]. Therefore, the impaired interaction between cholinergic signaling and their receptors on pancreatic acinar cells may be a mechanism of the pathogenesis of alcoholic pancreatitis.

Compared with AP, few CP models use injury mechanisms that may be related to the pathogenesis of human diseases, and the clinical relevance of the pathogenesis of most CP models is unclear[48]. However, Wan *et al*[55] used a mutant M3 receptor mouse-induced CP model and observed typical CP features, such as extensive chronic inflammation, fibrosis, adipose tissue infiltration, and pancreatic atrophy (Figure 1)[55]. The use of this M3 receptor model may increase our understanding of the pathophysiological process of human CP and identify specific treatments.

Long-term and recurrent pain is a common characteristic of CP[10,67]. A large number of studies have shown that cholinergic nerves have a significant analgesic effect on chronic neuropathic pain, inflammatory pain, and visceral pain[68-70]. Choline transporter (CHT1) is considered to be the rate-limiting step of neuronal acetylcholine synthesis and is essential for the effective recovery of acetylcholine[71,72]. CHT1 is upregulated in CP-induced pain models. CHT1 specific inhibitor, hemicholinium-3, can significantly enhance CP-induced hyperalgesia and reduce the amount of acetylcholine in the dorsal root ganglia of the pancreas in a dose-dependent manner[73]. Further research found that acetylcholine reduces pain and inflammation through $\alpha 7$ nicotinic acetylcholine receptors ($\alpha 7$ nAChRs) [70,74-77]. The activation of $\alpha 7$ nAChRs enhances the autophagic flux of acinar cells mediated by the transcription factor EB pathway and promotes lysosomal degradation to inhibit acinar cell damage, thereby protecting experimental pancreatitis. It has been suggested that cholinergic signaling activation of endogenous $\alpha 7$ nAChRs in the pancreas may be an endogenous protective mechanism in the process of pancreatitis[78]. Therefore, the use of $\alpha 7$ nAChRs to develop analgesic and anti-inflammatory drugs will be a promising target, especially for the treatment of CP.

CHOLINERGIC SIGNALING AND PCa

PCa is a gastrointestinal tumor with a poor prognosis, and more than 90% of PCa are pancreatic ductal adenocarcinoma (PDAC). According to reports, the 5-year survival rate for PCa patients in the United States from 2009 to 2015 was only 9%[79]. With the continuous increase in morbidity and mortality, PCa is expected to become the second leading cause of cancer-related deaths by 2030[80].

CP has been identified as a risk factor for PCa[81,82]. A meta-analysis found that CP increases the risk of PCa, and five years after the diagnosis of CP the risk of PCa increased nearly eightfold[83]. Pain is a characteristic feature of CP and PCa. Studies have found that increased nerve fiber density and hypertrophy are typical features of CP and PCa, and this pathological change appears to enhance and produce pancreatic neuropathic pain[84]. In vitro, myenteric plexus and dorsal root ganglia neurons were isolated from neonatal rats with CP and PCa, and they exhibited strong neurite outgrowth, more complex branching patterns, and somatic hypertrophy. They findings suggest that the intrapancreatic microenvironment in CP and PCa appears to be a key factor in the generation of pancreatic neuropathy and neural plasticity[85].

In-depth research on the tumor microenvironment found that perineural invasion is an important feature of PCa, which can lead to local tumor recurrence and poor prognosis[86]. As the degree of invasion increases, the survival rate of PCa patients is significantly reduced[87]. However, cholinergic signaling appears to play a tumor-suppressing role in PCa[14,88]. Regarding the relationship between heart rate variability (HRV) as an indicator of vagus nerve activity and the overall survival of patients with advanced PCa, it was found that higher vagus nerve activity represented by a higher initial HRV was significantly associated with a lower risk of death from PCa and was not affected by confounding factors such as age and cancer treatment[89]. The activation of M receptors by cholinergic signals can inhibit the progression of pancreatic tumors. Inhibition of the downstream EGFR/MAPK and PI3K/AKT signaling pathways of PCa cells through the M1 receptor signaling pathway inhibits tumor stem cells, CD11b⁺ cells, tumor necrosis factor- α levels, and liver metastasis[14].

However, M3 receptor expression seems to be a biomarker for the poor prognosis of PDAC. Compared with patients with low M3 expression, patients with high M3 expression have a worse prognosis and shorter survival time, but the study did not find a statistically significant relationship between M3 receptor expression and peripheral nerve infiltration in PCa[90]. The presence of nonneuronal acetylcholine secreted by fibroblasts and pancreatic stellate cells in the microenvironment of pancreatic tumors activates the M3 receptor, leading to further tumor progression[91].

Oxidative stress and inflammatory signaling contribute to the development of pancreatitis. Español *et al*[92] reported that long-term inflammatory stimulation by LPS plus interferon- γ (IFN- γ) could induce muscarinic receptor expression which mainly involved the M3 and M5 receptors. It could also upregulate the expression of NOS and COX-2 to enhance the effect of carbachol on NO and PGE₂ production[92]. Transgenic overexpression of COX-2 in the pancreas induces CP and the formation of preinvasive ductal tumors[93,94]. Muscarinic receptors could serve as promising therapeutic targets for pancreatic inflammation and could prevent the transformation of CP to PCa.

The increase in acetylcholine levels by nAChRs can inhibit histone deacetylase 1-mediated CCL5 chemokines. This could weaken the ability of PDAC cells to recruit CD8⁺ T cells and directly inhibit the production of IFN- γ by CD8⁺ T cells. This effect is conducive to Th2 differentiation and could thereby promote tumor growth[86]. Smoking is one of the main causes of PCa[95], and approximately 21% of PCa deaths are attributed to smoking[96]. Studies have shown that nicotine (nAChR agonist) can increase the proliferation activity and self-renewal ability of PCa stem cells by activating the sonic hedgehog signaling pathway[97]. $\alpha 7$ nAChRs have also been shown to upregulate mucin-4 by coactivating the JAK2/STAT3 downstream signaling cascade *via* the MEK/ERK1/2 pathway, thereby increasing the migration and invasion capabilities of PCa cells[98]. Because the pancreas is widely innervated by many different neurons, the relationship between PCa and the nervous system is complicated[99]. Cholinergic signaling produces different effects based on the different receptor mechanisms involved and may become a potential therapeutic target for PCa in the future.

CONCLUSION

Cholinergic signaling participates in the physiological function of the pancreas and the pathological process of pancreatic diseases. Although the mechanism by which cholinergic signaling regulates pancreatic diseases is still unclear, an increasing number of studies have shown that cholinergic signaling plays a key role in the occurrence and development of pancreatic diseases. There are many animal models of pancreatitis that can be used to help us study the pathogenesis and pathophysiological process of pancreatitis. Among them, rodent models are most commonly used to study acute and CP, but experimental pancreatitis is not necessarily the most relevant to human diseases[48]. Due to the lack of an in-depth understanding of its pathogenesis, there is no effective prevention and treatment strategy at present. Activation of the M3 receptor may be one of the causes of pancreatitis since muscarinic receptors are widely expressed in various glands and smooth muscles throughout the body. The mutant M3 receptor model provides an alternative method for the study of pancreatitis[55]. The pancreas is an organ rich in nervous tissue and lies behind the peritoneum and is surrounded by many nerve plexus. Cholinergic signaling has different effects due to its action on different receptors, which leads to a complicated pathogenesis of pancreatic diseases such as AP, CP, and pancreatic tumors (Table 1). Reducing or enhancing the signal downstream of the receptor may provide a potential therapeutic target in the future.

Table 1 Role of different cholinergic receptors in pancreatic diseases

Pancreatic diseases	Receptor type	Relevant mechanism	Effect	Ref.
Acute pancreatitis	M3	Receptor overexpression	Acinar cell hypersecretion	[49,50,52,55]
Chronic pancreatitis	M3	Cholinergic signaling increases	Acinar cell hypersecretion	[13,64-66]
	M3/M5	Receptor overexpression	Induce fibroblast proliferation	[92]
	$\alpha 7$	Enhances the autophagic flux of acinar cells	Inhibit acinar cell damage	[70,74-77]
Pancreatic cancer	M1	Inhibition of the EGFR/MAPK and PI3K/AKT signaling pathways	Inhibit the progression of pancreatic tumors	[14]
	M3	Receptor overexpression	Induction of preinvasive ductal tumor formation	[90,92-94]
	$\alpha 7$	Activating the JAK2/STAT3 signaling pathway	Increasing the migration and invasion capabilities of tumor cells	[86,97,98]

FOOTNOTES

Author contributions: Yang JM and Yang XY conceived the study, participated in the design, and drafted the manuscript; Wan JH edited and checked the manuscript; all of the authors have read and approved the final manuscript.

Supported by National Natural Science Foundation of China, No. 82100686.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Jian-Hua Wan 0000-0002-1364-3844.

S-Editor: Fan JR

L-Editor: A

P-Editor: Fan JR

REFERENCES

- Caulfield MP, Birdsall NJ. International Union of Pharmacology. XVII. Classification of muscarinic acetylcholine receptors. *Pharmacol Rev* 1998; **50**: 279-290 [PMID: 9647869]
- Qin K, Dong C, Wu G, Lambert NA. Inactive-state preassembly of G(q)-coupled receptors and G(q) heterotrimers. *Nat Chem Biol* 2011; **7**: 740-747 [PMID: 21873996 DOI: 10.1038/nchembio.642]
- Wess J. Molecular biology of muscarinic acetylcholine receptors. *Crit Rev Neurobiol* 1996; **10**: 69-99 [PMID: 8853955 DOI: 10.1615/critrevneurobiol.v10.i1.40]
- Volpicelli LA, Levey AI. Muscarinic acetylcholine receptor subtypes in cerebral cortex and hippocampus. *Prog Brain Res* 2004; **145**: 59-66 [PMID: 14650906 DOI: 10.1016/S0079-6123(03)45003-6]
- Abrams P, Andersson KE, Buccafusco JJ, Chapple C, de Groat WC, Fryer AD, Kay G, Latices A, Nathanson NM, Pasricha PJ, Wein AJ. Muscarinic receptors: their distribution and function in body systems, and the implications for treating overactive bladder. *Br J Pharmacol* 2006; **148**: 565-578 [PMID: 16751797 DOI: 10.1038/sj.bjp.0706780]
- Wang J, Lindstrom J. Orthosteric and allosteric potentiation of heteromeric neuronal nicotinic acetylcholine receptors. *Br J Pharmacol* 2018; **175**: 1805-1821 [PMID: 28199738 DOI: 10.1111/bph.13745]
- Wonnacott S, Bermudez I, Millar NS, Tzartos SJ. Nicotinic acetylcholine receptors. *Br J Pharmacol* 2018; **175**: 1785-1788 [PMID: 29878346 DOI: 10.1111/bph.14209]
- Hurst R, Rollema H, Bertrand D. Nicotinic acetylcholine receptors: from basic science to therapeutics. *Pharmacol Ther* 2013; **137**: 22-54 [PMID: 22925690 DOI: 10.1016/j.pharmthera.2012.08.012]
- Mizrahi JD, Surana R, Valle JW, Shroff RT. Pancreatic cancer. *Lancet* 2020; **395**: 2008-2020 [PMID: 32593337 DOI: 10.1016/S0140-6736(20)30974-0]
- Beyer G, Habtezion A, Werner J, Lerch MM, Mayerle J. Chronic pancreatitis. *Lancet* 2020; **396**: 499-512 [PMID: 32593337 DOI: 10.1016/S0140-6736(20)30974-0]

- 32798493 DOI: [10.1016/S0140-6736\(20\)31318-0](https://doi.org/10.1016/S0140-6736(20)31318-0)]
- 11 **Petrov MS**, Yadav D. Global epidemiology and holistic prevention of pancreatitis. *Nat Rev Gastroenterol Hepatol* 2019; **16**: 175-184 [PMID: [30482911](https://pubmed.ncbi.nlm.nih.gov/30482911/) DOI: [10.1038/s41575-018-0087-5](https://doi.org/10.1038/s41575-018-0087-5)]
 - 12 **Gr nroos JM**, Kaila T, Aho HJ, Nevalainen TJ. Decrease in the number of muscarinic receptors in rat pancreas after chronic alcohol intake. *Pharmacol Toxicol* 1989; **64**: 356-359 [PMID: [2748543](https://pubmed.ncbi.nlm.nih.gov/2748543/) DOI: [10.1111/j.1600-0773.1989.tb00664.x](https://doi.org/10.1111/j.1600-0773.1989.tb00664.x)]
 - 13 **Tiscornia OM**, Dreiling DA. Physiopathogenic hypothesis of alcoholic pancreatitis: supranormal ecbolic stimulation of the "pancreon" units secondary to the loss of the negative component of pancreas innervation. *Pancreas* 1987; **2**: 604-612 [PMID: [3313381](https://pubmed.ncbi.nlm.nih.gov/3313381/)]
 - 14 **Renz BW**, Tanaka T, Sunagawa M, Takahashi R, Jiang Z, Macchini M, Dantes Z, Valenti G, White RA, Middelhoff MA, Ilmer M, Oberstein PE, Angele MK, Deng H, Hayakawa Y, Westphalen CB, Werner J, Remotti H, Reichert M, Taylor YH, Nagar K, Friedman RA, Iuga AC, Olive KP, Wang TC. Cholinergic Signaling via Muscarinic Receptors Directly and Indirectly Suppresses Pancreatic Tumorigenesis and Cancer Stemness. *Cancer Discov* 2018; **8**: 1458-1473 [PMID: [30185628](https://pubmed.ncbi.nlm.nih.gov/30185628/) DOI: [10.1158/2159-8290.CD-18-0046](https://doi.org/10.1158/2159-8290.CD-18-0046)]
 - 15 **Konturek SJ**, Pepera J, Zabielski K, Konturek PC, Pawlik T, Szlachcic A, Hahn EG. Brain-gut axis in pancreatic secretion and appetite control. *J Physiol Pharmacol* 2003; **54**: 293-317 [PMID: [14566070](https://pubmed.ncbi.nlm.nih.gov/14566070/)]
 - 16 **Anagnostides A**, Chadwick VS, Selden AC, Maton PN. Sham feeding and pancreatic secretion. Evidence for direct vagal stimulation of enzyme output. *Gastroenterology* 1984; **87**: 109-114 [PMID: [6724252](https://pubmed.ncbi.nlm.nih.gov/6724252/)]
 - 17 **Konturek SJ**, Zabielski R, Konturek JW, Czarniecki J. Neuroendocrinology of the pancreas; role of brain-gut axis in pancreatic secretion. *Eur J Pharmacol* 2003; **481**: 1-14 [PMID: [14637169](https://pubmed.ncbi.nlm.nih.gov/14637169/) DOI: [10.1016/j.ejphar.2003.08.042](https://doi.org/10.1016/j.ejphar.2003.08.042)]
 - 18 **Defilippi C**, Solomon TE, Valenzuela JE. Pancreatic secretory response to sham feeding in humans. *Digestion* 1982; **23**: 217-223 [PMID: [6183160](https://pubmed.ncbi.nlm.nih.gov/6183160/) DOI: [10.1159/000198753](https://doi.org/10.1159/000198753)]
 - 19 **Owyang C**. Physiological mechanisms of cholecystokinin action on pancreatic secretion. *Am J Physiol* 1996; **271**: G1-G7 [PMID: [8760100](https://pubmed.ncbi.nlm.nih.gov/8760100/) DOI: [10.1152/ajpgi.1996.271.1.G1](https://doi.org/10.1152/ajpgi.1996.271.1.G1)]
 - 20 **Owyang C**, Logsdon CD. New insights into neurohormonal regulation of pancreatic secretion. *Gastroenterology* 2004; **127**: 957-969 [PMID: [15362050](https://pubmed.ncbi.nlm.nih.gov/15362050/) DOI: [10.1053/j.gastro.2004.05.002](https://doi.org/10.1053/j.gastro.2004.05.002)]
 - 21 **Niebergall-Roth E**, Singer MV. Central and peripheral neural control of pancreatic exocrine secretion. *J Physiol Pharmacol* 2001; **52**: 523-538 [PMID: [11787756](https://pubmed.ncbi.nlm.nih.gov/11787756/)]
 - 22 **Singer MV**, Niebergall-Roth E. Secretion from acinar cells of the exocrine pancreas: role of enteropancreatic reflexes and cholecystokinin. *Cell Biol Int* 2009; **33**: 1-9 [PMID: [18948215](https://pubmed.ncbi.nlm.nih.gov/18948215/) DOI: [10.1016/j.cellbi.2008.09.008](https://doi.org/10.1016/j.cellbi.2008.09.008)]
 - 23 **Becker S**, Niebel W, Singer MV. Nervous control of gastric and pancreatic secretory response to 2-deoxy-D-glucose in the dog. *Digestion* 1988; **39**: 187-196 [PMID: [3209002](https://pubmed.ncbi.nlm.nih.gov/3209002/) DOI: [10.1159/000199624](https://doi.org/10.1159/000199624)]
 - 24 **Holtmann G**, Singer MV, Kriebel R, St cker KH, Goebell H. Differential effects of acute mental stress on interdigestive secretion of gastric acid, pancreatic enzymes, and gastroduodenal motility. *Dig Dis Sci* 1989; **34**: 1701-1707 [PMID: [2582983](https://pubmed.ncbi.nlm.nih.gov/2582983/) DOI: [10.1007/BF01540047](https://doi.org/10.1007/BF01540047)]
 - 25 **Chariot J**, de la Tour J, Anglade P, Roz  C. Cholinergic mechanisms in the pancreas after extrinsic denervation in the rat. *Am J Physiol* 1987; **252**: G755-G761 [PMID: [3591943](https://pubmed.ncbi.nlm.nih.gov/3591943/) DOI: [10.1152/ajpgi.1987.252.6.G755](https://doi.org/10.1152/ajpgi.1987.252.6.G755)]
 - 26 **Park HS**, Lee YL, Kwon HY, Chey WY, Park HJ. Significant cholinergic role in secretin-stimulated exocrine secretion in isolated rat pancreas. *Am J Physiol* 1998; **274**: G413-G418 [PMID: [9486197](https://pubmed.ncbi.nlm.nih.gov/9486197/) DOI: [10.1152/ajpgi.1998.274.2.G413](https://doi.org/10.1152/ajpgi.1998.274.2.G413)]
 - 27 **You CH**, Rominger JM, Chey WY. Effects of atropine on the action and release of secretin in humans. *Am J Physiol* 1982; **242**: G608-G611 [PMID: [7091334](https://pubmed.ncbi.nlm.nih.gov/7091334/) DOI: [10.1152/ajpgi.1982.242.6.G608](https://doi.org/10.1152/ajpgi.1982.242.6.G608)]
 - 28 **Blair EL**, Brown JC, Harper AA, Scratcherd T. A gastric phase of pancreatic secretion. *J Physiol* 1966; **184**: 812-824 [PMID: [5912208](https://pubmed.ncbi.nlm.nih.gov/5912208/) DOI: [10.1113/jphysiol.1966.sp007949](https://doi.org/10.1113/jphysiol.1966.sp007949)]
 - 29 **Gabrylewicz A**, Kulesza E, Konturek SJ. Comparison of loxiglumide, a cholecystokinin receptor antagonist, and atropine on hormonal and meal-stimulated pancreatic secretion in man. *Scand J Gastroenterol* 1990; **25**: 731-738 [PMID: [2396088](https://pubmed.ncbi.nlm.nih.gov/2396088/) DOI: [10.3109/00365529008997600](https://doi.org/10.3109/00365529008997600)]
 - 30 **Lugea A**, Waldron RT, Mareninova OA, Shalbuva N, Deng N, Su HY, Thomas DD, Jones EK, Messenger SW, Yang J, Hu C, Gukovsky I, Liu Z, Groblewski GE, Gukovskaya AS, Gorelick FS, Pandol SJ. Human Pancreatic Acinar Cells: Proteomic Characterization, Physiologic Responses, and Organellar Disorders in ex Vivo Pancreatitis. *Am J Pathol* 2017; **187**: 2726-2743 [PMID: [28935577](https://pubmed.ncbi.nlm.nih.gov/28935577/) DOI: [10.1016/j.ajpath.2017.08.017](https://doi.org/10.1016/j.ajpath.2017.08.017)]
 - 31 **Ji B**, Bi Y, Simeone D, Mortensen RM, Logsdon CD. Human pancreatic acinar cells lack functional responses to cholecystokinin and gastrin. *Gastroenterology* 2001; **121**: 1380-1390 [PMID: [11729117](https://pubmed.ncbi.nlm.nih.gov/11729117/) DOI: [10.1053/gast.2001.29557](https://doi.org/10.1053/gast.2001.29557)]
 - 32 **Futatsugi A**, Nakamura T, Yamada MK, Ebisui E, Nakamura K, Uchida K, Kitaguchi T, Takahashi-Iwanaga H, Noda T, Aruga J, Mikoshiba K. IP3 receptor types 2 and 3 mediate exocrine secretion underlying energy metabolism. *Science* 2005; **309**: 2232-2234 [PMID: [16195467](https://pubmed.ncbi.nlm.nih.gov/16195467/) DOI: [10.1126/science.1114110](https://doi.org/10.1126/science.1114110)]
 - 33 **Wakui M**, Osipchuk YV, Petersen OH. Receptor-activated cytoplasmic Ca²⁺ spiking mediated by inositol trisphosphate is due to Ca₂(+)-induced Ca₂⁺ release. *Cell* 1990; **63**: 1025-1032 [PMID: [1701691](https://pubmed.ncbi.nlm.nih.gov/1701691/) DOI: [10.1016/0092-8674\(90\)90505-9](https://doi.org/10.1016/0092-8674(90)90505-9)]
 - 34 **Louie DS**, Owyang C. Muscarinic receptor subtypes on rat pancreatic acini: secretion and binding studies. *Am J Physiol* 1986; **251**: G275-G279 [PMID: [2426974](https://pubmed.ncbi.nlm.nih.gov/2426974/) DOI: [10.1152/ajpgi.1986.251.2.G275](https://doi.org/10.1152/ajpgi.1986.251.2.G275)]
 - 35 **Iwatsuki K**, Horiuchi A, Yonekura H, Homma N, Haruta K, Chiba S. Subtypes of muscarinic receptors in pancreatic exocrine secretion in anesthetized dog. *Pancreas* 1989; **4**: 339-345 [PMID: [2734277](https://pubmed.ncbi.nlm.nih.gov/2734277/) DOI: [10.1097/00006676-198906000-00011](https://doi.org/10.1097/00006676-198906000-00011)]
 - 36 **van Zwam AJ**, Willems PH, Rodrigues de Miranda JF, de Pont JJ, van Ginneken CA. Binding characteristics of the muscarinic receptor subtype in rabbit pancreas. *J Recept Res* 1990; **10**: 119-135 [PMID: [2262932](https://pubmed.ncbi.nlm.nih.gov/2262932/) DOI: [10.3109/10799899009064661](https://doi.org/10.3109/10799899009064661)]
 - 37 **Singer MV**, Teyssen S, K ppers U. Influence of the M1-receptor antagonists telenzepine and pirenzepine on pancreatic secretory response to intraduodenal tryptophan in dogs. *Digestion* 1991; **48**: 34-42 [PMID: [1868967](https://pubmed.ncbi.nlm.nih.gov/1868967/) DOI: [10.1159/000200661](https://doi.org/10.1159/000200661)]
 - 38 **Teyssen S**, Niebergall E, Chari ST, Singer MV. Comparison of two dose-response techniques to study the pancreatic

- secretory response to intraduodenal tryptophan in the absence and presence of the M1-receptor antagonist telenzepine. *Pancreas* 1995; **10**: 368-373 [PMID: 7792293 DOI: 10.1097/00006676-199505000-00008]
- 39 **Schmid SW**, Modlin IM, Tang LH, Stoch A, Rhee S, Nathanson MH, Scheele GA, Gorelick FS. Telenzepine-sensitive muscarinic receptors on rat pancreatic acinar cells. *Am J Physiol* 1998; **274**: G734-G741 [PMID: 9575856 DOI: 10.1152/ajpgi.1998.274.4.G734]
- 40 **Gautam D**, Han SJ, Heard TS, Cui Y, Miller G, Bloodworth L, Wess J. Cholinergic stimulation of amylase secretion from pancreatic acinar cells studied with muscarinic acetylcholine receptor mutant mice. *J Pharmacol Exp Ther* 2005; **313**: 995-1002 [PMID: 15764735 DOI: 10.1124/jpet.105.084855]
- 41 **Lankisch PG**, Apte M, Banks PA. Acute pancreatitis. *Lancet* 2015; **386**: 85-96 [PMID: 25616312 DOI: 10.1016/S0140-6736(14)60649-8]
- 42 **Bartholomew C**. Acute scorpion pancreatitis in Trinidad. *Br Med J* 1970; **1**: 666-668 [PMID: 5443968 DOI: 10.1136/bmj.1.5697.666]
- 43 **Singh S**, Bhardwaj U, Verma SK, Bhalla A, Gill K. Hyperamylasemia and acute pancreatitis following anticholinesterase poisoning. *Hum Exp Toxicol* 2007; **26**: 467-471 [PMID: 17698941 DOI: 10.1177/0960327107076814]
- 44 **Roeyen G**, Chapelle T, Jorens P, de Beeck BO, Ysebaert D. Necrotizing pancreatitis due to poisoning with organophosphate pesticides. *Acta Gastroenterol Belg* 2008; **71**: 27-29 [PMID: 18396746]
- 45 **Marsh WH**, Vukov GA, Conradi EC. Acute pancreatitis after cutaneous exposure to an organophosphate insecticide. *Am J Gastroenterol* 1988; **83**: 1158-1160 [PMID: 3421227]
- 46 **Becerril B**, Marangoni S, Possani LD. Toxins and genes isolated from scorpions of the genus Tityus. *Toxicon* 1997; **35**: 821-835 [PMID: 9241777 DOI: 10.1016/S0041-0101(96)00198-5]
- 47 **Fletcher PL Jr**, Fletcher MD, Possani LD. Characteristics of pancreatic exocrine secretion produced by venom from the Brazilian scorpion, Tityus serrulatus. *Eur J Cell Biol* 1992; **58**: 259-270 [PMID: 1385124]
- 48 **Lerch MM**, Gorelick FS. Models of acute and chronic pancreatitis. *Gastroenterology* 2013; **144**: 1180-1193 [PMID: 23622127 DOI: 10.1053/j.gastro.2012.12.043]
- 49 **Samuel I**, Chaudhary A, Fisher RA, Joehl RJ. Exacerbation of acute pancreatitis by combined cholinergic stimulation and duct obstruction. *Am J Surg* 2005; **190**: 721-724 [PMID: 16226947 DOI: 10.1016/j.amjsurg.2005.07.010]
- 50 **Samuel I**, Zaheer S, Fisher RA, Zaheer A. Cholinergic receptor induction and JNK activation in acute pancreatitis. *Am J Surg* 2003; **186**: 569-574 [PMID: 14599627 DOI: 10.1016/j.amjsurg.2003.07.016]
- 51 **Grönroos JM**, Aho HJ, Nevalainen TJ. Cholinergic hypothesis of alcoholic pancreatitis. *Dig Dis* 1992; **10**: 38-45 [PMID: 1551245 DOI: 10.1159/000171342]
- 52 **Lugea A**, Gong J, Nguyen J, Nieto J, French SW, Pandolfi SJ. Cholinergic mediation of alcohol-induced experimental pancreatitis. *Alcohol Clin Exp Res* 2010; **34**: 1768-1781 [PMID: 20626730 DOI: 10.1111/j.1530-0277.2010.01264.x]
- 53 **Grönroos JM**, Kaila T, Hietaranta AJ. Alcohol, pancreatic muscarinic receptors and acute pancreatitis. *Exp Toxicol Pathol* 1994; **45**: 503-505 [PMID: 8054828 DOI: 10.1016/S0940-2993(11)80513-0]
- 54 **Satoh A**, Gukovskaya AS, Reeve JR Jr, Shimosegawa T, Pandolfi SJ. Ethanol sensitizes NF-kappaB activation in pancreatic acinar cells through effects on protein kinase C-epsilon. *Am J Physiol Gastrointest Liver Physiol* 2006; **291**: G432-G438 [PMID: 16574982 DOI: 10.1152/ajpgi.00579.2005]
- 55 **Wan J**, Wang J, Wagner LE 2nd, Wang OH, Gui F, Chen J, Zhu X, Haddock AN, Edenfield BH, Haight B, Mukhopadhyay D, Wang Y, Yule DI, Bi Y, Ji B. Pancreas-specific CHRM3 activation causes pancreatitis in mice. *JCI Insight* 2021; **6** [PMID: 34314386 DOI: 10.1172/jci.insight.132585]
- 56 **Zhan X**, Wang F, Bi Y, Ji B. Animal models of gastrointestinal and liver diseases. Animal models of acute and chronic pancreatitis. *Am J Physiol Gastrointest Liver Physiol* 2016; **311**: G343-G355 [PMID: 27418683 DOI: 10.1152/ajpgi.00372.2015]
- 57 **Sah RP**, Saluja A. Molecular mechanisms of pancreatic injury. *Curr Opin Gastroenterol* 2011; **27**: 444-451 [PMID: 21844752 DOI: 10.1097/MOG.0b013e328349e346]
- 58 **Kambhampati S**, Park W, Habtezion A. Pharmacologic therapy for acute pancreatitis. *World J Gastroenterol* 2014; **20**: 16868-16880 [PMID: 25493000 DOI: 10.3748/wjg.v20.i45.16868]
- 59 **Singh VK**, Yadav D, Garg PK. Diagnosis and Management of Chronic Pancreatitis: A Review. *JAMA* 2019; **322**: 2422-2434 [PMID: 31860051 DOI: 10.1001/jama.2019.19411]
- 60 **Wang YT**, Gou YW, Jin WW, Xiao M, Fang HY. Association between alcohol intake and the risk of pancreatic cancer: a dose-response meta-analysis of cohort studies. *BMC Cancer* 2016; **16**: 212 [PMID: 26968702 DOI: 10.1186/s12885-016-2241-1]
- 61 **Samokhvalov AV**, Rehm J, Roerecke M. Alcohol Consumption as a Risk Factor for Acute and Chronic Pancreatitis: A Systematic Review and a Series of Meta-analyses. *EBioMedicine* 2015; **2**: 1996-2002 [PMID: 26844279 DOI: 10.1016/j.ebiom.2015.11.023]
- 62 **Klochov A**, Kudaravalli P, Lim Y, Sun Y. Alcoholic Pancreatitis. StatPearls. Treasure Island (FL), 2021
- 63 **Hirano H**, Shimosegawa T, Meguro T, Shiga N, Koizumi M, Toyota T. Effects of ethanol on meal-stimulated secretion of pancreatic polypeptide and cholecystokinin: comparison of healthy volunteers, heavy drinkers, and patients with chronic pancreatitis. *J Gastroenterol* 1996; **31**: 86-93 [PMID: 8808434 DOI: 10.1007/BF01211192]
- 64 **Gregg JA**, Sharma MM. Endoscopic measurement of pancreatic juice secretory flow rates and pancreatic secretory pressures after secretin administration in human controls and in patients with acute relapsing pancreatitis, chronic pancreatitis, and pancreatic cancer. *Am J Surg* 1978; **136**: 569-574 [PMID: 707739 DOI: 10.1016/0002-9610(78)90312-4]
- 65 **Renner IG**, Rinderknecht H, Valenzuela JE, Douglas AP. Studies of pure pancreatic secretions in chronic alcoholic subjects without pancreatic insufficiency. *Scand J Gastroenterol* 1980; **15**: 241-244 [PMID: 7384747 DOI: 10.3109/00365528009181462]
- 66 **Ceyhan GO**, Demir IE, Rauch U, Bergmann F, Müller MW, Büchler MW, Friess H, Schäfer KH. Pancreatic neuropathy results in "neural remodeling" and altered pancreatic innervation in chronic pancreatitis and pancreatic cancer. *Am J Gastroenterol* 2009; **104**: 2555-2565 [PMID: 19568227 DOI: 10.1038/ajg.2009.380]
- 67 **Di Sebastiano P**, di Mola FF, Bockman DE, Friess H, Büchler MW. Chronic pancreatitis: the perspective of pain

- generation by neuroimmune interaction. *Gut* 2003; **52**: 907-911 [PMID: 12740353 DOI: 10.1136/gut.52.6.907]
- 68 **Zhang XL**, Albers KM, Gold MS. Inflammation-induced increase in nicotinic acetylcholine receptor current in cutaneous nociceptive DRG neurons from the adult rat. *Neuroscience* 2015; **284**: 483-499 [PMID: 25453771 DOI: 10.1016/j.neuroscience.2014.10.018]
- 69 **Joshi SK**, Mikusa JP, Weaver B, Honore P. Morphine and ABT-594 (a nicotinic acetylcholine agonist) exert centrally mediated antinociception in the rat cyclophosphamide cystitis model of visceral pain. *J Pain* 2008; **9**: 146-156 [PMID: 18088559 DOI: 10.1016/j.jpain.2007.09.004]
- 70 **Freitas K**, Ghosh S, Ivy Carroll F, Lichtman AH, Imad Damaj M. Effects of $\alpha 7$ positive allosteric modulators in murine inflammatory and chronic neuropathic pain models. *Neuropharmacology* 2013; **65**: 156-164 [PMID: 23079470 DOI: 10.1016/j.neuropharm.2012.08.022]
- 71 **Okuda T**, Haga T. High-affinity choline transporter. *Neurochem Res* 2003; **28**: 483-488 [PMID: 12675135 DOI: 10.1023/a:1022809003997]
- 72 **Yamamura HI**, Snyder SH. Choline: high-affinity uptake by rat brain synaptosomes. *Science* 1972; **178**: 626-628 [PMID: 5086398 DOI: 10.1126/science.178.4061.626]
- 73 **Luo D**, Chen L, Yu B. Inhibition of the high affinity choline transporter enhances hyperalgesia in a rat model of chronic pancreatitis. *Biochem Biophys Res Commun* 2017; **488**: 204-210 [PMID: 28483526 DOI: 10.1016/j.bbrc.2017.05.036]
- 74 **van Westerloo DJ**, Giebelen IA, Florquin S, Bruno MJ, Larosa GJ, Ulloa L, Tracey KJ, van der Poll T. The vagus nerve and nicotinic receptors modulate experimental pancreatitis severity in mice. *Gastroenterology* 2006; **130**: 1822-1830 [PMID: 16697744 DOI: 10.1053/j.gastro.2006.02.022]
- 75 **Schneider L**, Jabrailova B, Soliman H, Hofer S, Strobel O, Hackert T, Büchler MW, Werner J. Pharmacological cholinergic stimulation as a therapeutic tool in experimental necrotizing pancreatitis. *Pancreas* 2014; **43**: 41-46 [PMID: 24212240 DOI: 10.1097/MPA.0b013e3182a85c21]
- 76 **Damaj MI**, Meyer EM, Martin BR. The antinociceptive effects of $\alpha 7$ nicotinic agonists in an acute pain model. *Neuropharmacology* 2000; **39**: 2785-2791 [PMID: 11044748 DOI: 10.1016/s0028-3908(00)00139-8]
- 77 **Rowley TJ**, McKinstry A, Greenidge E, Smith W, Flood P. Antinociceptive and anti-inflammatory effects of choline in a mouse model of postoperative pain. *Br J Anaesth* 2010; **105**: 201-207 [PMID: 20511332 DOI: 10.1093/bja/aeq113]
- 78 **Li B**, Wu J, Bao J, Han X, Shen S, Ye X, Dai J, Wu Z, Niu M, He Y, Ni J, Wen L, Wang X, Hu G. Activation of $\alpha 7$ nACh receptor protects against acute pancreatitis through enhancing TFEB-regulated autophagy. *Biochim Biophys Acta Mol Basis Dis* 2020; **1866**: 165971 [PMID: 32950676 DOI: 10.1016/j.bbadis.2020.165971]
- 79 **Siegel RL**, Miller KD, Jemal A. Cancer statistics, 2020. *CA Cancer J Clin* 2020; **70**: 7-30 [PMID: 31912902 DOI: 10.3322/caac.21590]
- 80 **Rahib L**, Smith BD, Aizenberg R, Rosenzweig AB, Fleshman JM, Matrisian LM. Projecting cancer incidence and deaths to 2030: the unexpected burden of thyroid, liver, and pancreas cancers in the United States. *Cancer Res* 2014; **74**: 2913-2921 [PMID: 24840647 DOI: 10.1158/0008-5472.CAN-14-0155]
- 81 **Lowenfels AB**, Maisonneuve P, Cavallini G, Ammann RW, Lankisch PG, Andersen JR, Dimagno EP, Andrén-Sandberg A, Domellöf L. Pancreatitis and the risk of pancreatic cancer. International Pancreatitis Study Group. *N Engl J Med* 1993; **328**: 1433-1437 [PMID: 8479461 DOI: 10.1056/NEJM199305203282001]
- 82 **Malka D**, Hammel P, Maire F, Rufat P, Madeira I, Pessione F, Lévy P, Ruzsniwski P. Risk of pancreatic adenocarcinoma in chronic pancreatitis. *Gut* 2002; **51**: 849-852 [PMID: 12427788 DOI: 10.1136/gut.51.6.849]
- 83 **Kirkegård J**, Mortensen FV, Cronin-Fenton D. Chronic Pancreatitis and Pancreatic Cancer Risk: A Systematic Review and Meta-analysis. *Am J Gastroenterol* 2017; **112**: 1366-1372 [PMID: 28762376 DOI: 10.1038/ajg.2017.218]
- 84 **Ceyhan GO**, Bergmann F, Kadihasanoglu M, Altintas B, Demir IE, Hinz U, Müller MW, Giese T, Büchler MW, Giese NA, Friess H. Pancreatic neuropathy and neuropathic pain--a comprehensive pathomorphological study of 546 cases. *Gastroenterology* 2009; **136**: 177-186.e1 [PMID: 18992743 DOI: 10.1053/j.gastro.2008.09.029]
- 85 **Demir IE**, Ceyhan GO, Rauch U, Altintas B, Klotz M, Müller MW, Büchler MW, Friess H, Schäfer KH. The microenvironment in chronic pancreatitis and pancreatic cancer induces neuronal plasticity. *Neurogastroenterol Motil* 2010; **22**: 480-490, e112 [PMID: 19912545 DOI: 10.1111/j.1365-2982.2009.01428.x]
- 86 **Yang MW**, Tao LY, Jiang YS, Yang JY, Huo YM, Liu DJ, Li J, Fu XL, He R, Lin C, Liu W, Zhang JF, Hua R, Li Q, Jiang SH, Hu LP, Tian GA, Zhang XX, Niu N, Lu P, Shi J, Xiao GG, Wang LW, Xue J, Zhang ZG, Sun YW. Perineural Invasion Reprograms the Immune Microenvironment through Cholinergic Signaling in Pancreatic Ductal Adenocarcinoma. *Cancer Res* 2020; **80**: 1991-2003 [PMID: 32098780 DOI: 10.1158/0008-5472.CAN-19-2689]
- 87 **Liebl F**, Demir IE, Mayer K, Schuster T, D'Haese JG, Becker K, Langer R, Bergmann F, Wang K, Rosenberg R, Novotny AR, Feith M, Reim D, Friess H, Ceyhan GO. The impact of neural invasion severity in gastrointestinal malignancies: a clinicopathological study. *Ann Surg* 2014; **260**: 900-7; discussion 907 [PMID: 25379860 DOI: 10.1097/SLA.0000000000000968]
- 88 **Kamiya A**, Hiyama T, Fujimura A, Yoshikawa S. Sympathetic and parasympathetic innervation in cancer: therapeutic implications. *Clin Auton Res* 2021; **31**: 165-178 [PMID: 32926324 DOI: 10.1007/s10286-020-00724-y]
- 89 **De Couck M**, Maréchal R, Moorthamers S, Van Laethem JL, Gidron Y. Vagal nerve activity predicts overall survival in metastatic pancreatic cancer, mediated by inflammation. *Cancer Epidemiol* 2016; **40**: 47-51 [PMID: 26618335 DOI: 10.1016/j.canep.2015.11.007]
- 90 **Zhang L**, Xiu D, Zhan J, He X, Guo L, Wang J, Tao M, Fu W, Zhang H. High expression of muscarinic acetylcholine receptor 3 predicts poor prognosis in patients with pancreatic ductal adenocarcinoma. *Oncotargets Ther* 2016; **9**: 6719-6726 [PMID: 27826198 DOI: 10.2147/OTT.S111382]
- 91 **Pfützing PL**, Fangmann L, Wang K, Demir E, Gürlevik E, Fleischmann-Mundt B, Brooks J, D'Haese JG, Teller S, Hecker A, Jesinghaus M, Jäger C, Ren L, Istvanffy R, Kühnel F, Friess H, Ceyhan GO, Demir IE. Indirect cholinergic activation slows down pancreatic cancer growth and tumor-associated inflammation. *J Exp Clin Cancer Res* 2020; **39**: 289 [PMID: 33357230 DOI: 10.1186/s13046-020-01796-4]
- 92 **Español AJ**, Maddaleno MO, Lombardi MG, Cella M, Martínez Pulido P, Sales ME. Treatment with LPS plus INF- γ induces the expression and function of muscarinic acetylcholine receptors, modulating NIH3T3 cell proliferation:

- participation of NOS and COX. *Br J Pharmacol* 2014; **171**: 5154-5167 [PMID: 24990429 DOI: 10.1111/bph.12834]
- 93 **Colby JK**, Klein RD, McArthur MJ, Conti CJ, Kiguchi K, Kawamoto T, Riggs PK, Pavone AI, Sawicki J, Fischer SM. Progressive metaplastic and dysplastic changes in mouse pancreas induced by cyclooxygenase-2 overexpression. *Neoplasia* 2008; **10**: 782-796 [PMID: 18670639 DOI: 10.1593/neo.08330]
- 94 **Müller-Decker K**, Fürstenberger G, Annan N, Kucher D, Pohl-Arnold A, Steinbauer B, Esposito I, Chiblak S, Friess H, Schirmacher P, Berger I. Preinvasive duct-derived neoplasms in pancreas of keratin 5-promoter cyclooxygenase-2 transgenic mice. *Gastroenterology* 2006; **130**: 2165-2178 [PMID: 16762637 DOI: 10.1053/j.gastro.2006.03.053]
- 95 **Vincent A**, Herman J, Schulick R, Hruban RH, Goggins M. Pancreatic cancer. *Lancet* 2011; **378**: 607-620 [PMID: 21620466 DOI: 10.1016/S0140-6736(10)62307-0]
- 96 **GBD 2017 Pancreatic Cancer Collaborators**. . The global, regional, and national burden of pancreatic cancer and its attributable risk factors in 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet Gastroenterol Hepatol* 2019; **4**: 934-947 [PMID: 31648972 DOI: 10.1016/S2468-1253(19)30347-4]
- 97 **Al-Wadei MH**, Banerjee J, Al-Wadei HA, Schuller HM. Nicotine induces self-renewal of pancreatic cancer stem cells via neurotransmitter-driven activation of sonic hedgehog signalling. *Eur J Cancer* 2016; **52**: 188-196 [PMID: 26689865 DOI: 10.1016/j.ejca.2015.10.003]
- 98 **Momi N**, Ponnusamy MP, Kaur S, Rachagani S, Kunigal SS, Chellappan S, Ouellette MM, Batra SK. Nicotine/cigarette smoke promotes metastasis of pancreatic cancer through $\alpha 7nAChR$ -mediated MUC4 upregulation. *Oncogene* 2013; **32**: 1384-1395 [PMID: 22614008 DOI: 10.1038/onc.2012.163]
- 99 **Hutchings C**, Phillips JA, Djamgoz MBA. Nerve input to tumours: Pathophysiological consequences of a dynamic relationship. *Biochim Biophys Acta Rev Cancer* 2020; **1874**: 188411 [PMID: 32828885 DOI: 10.1016/j.bbcan.2020.188411]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

