

World Journal of *Gastrointestinal Endoscopy*

World J Gastrointest Endosc 2022 September 16; 14(9): 512-580



MINIREVIEWS

- 512 Simulation-based mastery learning in gastrointestinal endoscopy training
Maulahela H, Annisa NG, Konstantin T, Syam AF, Soetikno R

ORIGINAL ARTICLE**Case Control Study**

- 524 Endoscopic ultrasound elastography for malignant pancreatic masses and associated lymph nodes: Critical evaluation of strain ratio cutoff value
Puga-Tejada M, Del Valle R, Oleas R, Egas-Izquierdo M, Arevalo-Mora M, Baquerizo-Burgos J, Ospina J, Soria-Alcivar M, Pitanga-Lukashok H, Robles-Medranda C

Retrospective Study

- 536 Screening for hilar biliary invasion in ampullary cancer patients
Takagi T, Sugimoto M, Suzuki R, Konno N, Asama H, Sato Y, Irie H, Nakamura J, Takasumi M, Hashimoto M, Kato T, Kobashi R, Yanagita T, Hashimoto Y, Marubashi S, Hikichi T, Ohira H
- 547 Endoscopic therapy using a self-expandable metallic stent with an anti-migration system for postorthotopic liver transplantation anastomotic biliary stricture
Pinheiro LW, Martins FP, De Paulo GA, Contini MLC, Ferrari AP, Della Libera E

Observational Study

- 555 Clinical profile, diagnostic yield, and procedural outcomes of single balloon enteroscopy: A tertiary care hospital experience
Inam M, Karim MM, Tariq U, Ismail FW

Prospective Study

- 564 Role of endoscopic ultrasound in evaluation of patients with missed common bile duct stones
Eissa M, Okasha HH, Abbasy M, Khamis AK, Abdellatef A, Rady MA

CASE REPORT

- 575 Isolated esophageal tuberculosis: A case report
Diallo I, Touré O, Sarr ES, Sow A, Ndiaye B, Diawara PS, Dial CM, Mbengue A, Fall F

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Endoscopy*, Giuseppe Di Buono, MD, PhD, Doctor, Lecturer, Research Assistant Professor, Department of Surgical, Oncological and Oral Sciences, University of Palermo, Palermo 90127, Italy. giuseppe.dibuono@unipa.it

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Endoscopy* (*WJGE*, *World J Gastrointest Endosc*) is to provide scholars and readers from various fields of gastrointestinal endoscopy with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGE mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal endoscopy and covering a wide range of topics including capsule endoscopy, colonoscopy, double-balloon enteroscopy, duodenoscopy, endoscopic retrograde cholangiopancreatography, endosonography, esophagoscopy, gastrointestinal endoscopy, gastroscopy, laparoscopy, natural orifice endoscopic surgery, proctoscopy, and sigmoidoscopy.

INDEXING/ABSTRACTING

The *WJGE* is now abstracted and indexed in Emerging Sources Citation Index (Web of Science), PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 edition of Journal Citation Reports® cites the 2021 Journal Citation Indicator (JCI) for *WJGE* as 0.33.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yi-Xuan Cai; **Production Department Director:** Xu Guo; **Editorial Office Director:** Jia-Ping Yan.

NAME OF JOURNAL

World Journal of Gastrointestinal Endoscopy

ISSN

ISSN 1948-5190 (online)

LAUNCH DATE

October 15, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Anastasios Koulaouzidis, Bing Hu, Sang Chul Lee, Joo Young Cho

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-5190/editorialboard.htm>

PUBLICATION DATE

September 16, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Isolated esophageal tuberculosis: A case report

Ibrahima Diallo, Omar Touré, Elhadji Souleymane Sarr, Abdoul Sow, Bineta Ndiaye, Papa Silman Diawara, Cherif Mouhamed Dial, Ababacar Mbengue, Fatou Fall

Specialty type: Gastroenterology and hepatology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): B

Grade C (Good): C

Grade D (Fair): 0

Grade E (Poor): E

P-Reviewer: Anandan H, India; Meng FZ, China; Novita BD, Indonesia

Received: May 3, 2022

Peer-review started: May 5, 2022

First decision: June 16, 2022

Revised: July 11, 2022

Accepted: August 6, 2022

Article in press: August 6, 2022

Published online: September 16, 2022



Ibrahima Diallo, Hepato-Gastroenterology, Hopital Principal de Dakar, Dakar 3006, Senegal

Omar Touré, Abdoul Sow, Bineta Ndiaye, Fatou Fall, Hepatogastroenterology, Hopital Principal de Dakar, Dakar 3006, Senegal

Elhadji Souleymane Sarr, Department of Pathology, Hopital Principal de Dakar, Dakar 3006, Senegal

Papa Silman Diawara, Department of Biology, Hopital Principal de Dakar, Dakar 3006, Senegal

Cherif Mouhamed Dial, Department of Pathology, Hopital Général Idrissa Pouye, Dakar 3006, Senegal

Ababacar Mbengue, Department of Imaging, Hopital Principal de Dakar, Dakar 3006, Senegal

Corresponding author: Ibrahima Diallo, MD, Chief Doctor, Hepato-Gastroenterology, Hopital Principal de Dakar, 01 Avenue Nelson Mandela, Dakar 3006, Senegal. iduallo601@yahoo.fr

Abstract

BACKGROUND

Tuberculosis is endemic in Senegal. While its extra-pulmonary localization is rare, esophageal tuberculosis, particularly the isolated form, is exceptional. We report here a case of isolated esophageal tuberculosis in an immunocompetent patient.

CASE SUMMARY

A 58-year-old man underwent consultation for mechanical dysphagia that had developed over 3 mo with non-quantified weight loss, anorexia, and fever. Upper digestive endoscopy showed extensive ulcerated lesions, suggesting neoplasia. The diagnosis was confirmed by histopathology, which showed gigantocellular epithelioid granuloma surrounding a caseous necrosis. Thoracoabdominal computed tomography scan did not show another localization of the tuberculosis. The outcome was favorable with treatment.

CONCLUSION

Esophageal tuberculosis should be considered when dysphagia is associated with atypical ulcerated lesions of the esophageal mucosa, in an endemic area.

Key Words: Tuberculosis; Esophagus; Endoscopy; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Isolated esophageal tuberculosis is rare. Often discovered during the exploration of dysphagia, the endoscopic aspects are not specific, and can simulate several pathologies. Biopsies can help with diagnosis by showing the granuloma to histology or by allowing molecular biology examinations. In this manuscript, we report a case of isolated esophageal tuberculosis with vast ulcers of the esophagus, which evolved without sequelae after treatment.

Citation: Diallo I, Touré O, Sarr ES, Sow A, Ndiaye B, Diawara PS, Dial CM, Mbengue A, Fall F. Isolated esophageal tuberculosis: A case report. *World J Gastrointest Endosc* 2022; 14(9): 575-580

URL: <https://www.wjgnet.com/1948-5190/full/v14/i9/575.htm>

DOI: <https://dx.doi.org/10.4253/wjge.v14.i9.575>

INTRODUCTION

Tuberculosis is endemic in Senegal, where it constitutes a major public health problem. In 2020, 12808 new cases of tuberculosis were reported in Senegal, the majority of which were pulmonary (National Controlling Tuberculosis Program, data not published). Extrapulmonary forms of tuberculosis are frequent, whether or not they are associated with pulmonary involvement. In the digestive tract, the terminal ileum and the cecum are most often affected. Esophageal localization is rare, especially in its isolated form. We report herein a case of isolated esophageal tuberculosis in an immunocompetent patient who responded well to antibacillary treatment.

CASE PRESENTATION

Chief complaints

A 58-year-old patient was seen in our department for dysphagia that had developed over 3 mo.

History of present illness

The patient had dysphagia that had been evolving for 3 mo with non-quantified weight loss, nonselective anorexia, and nocturnal fever.

History of past illness

The patient had undergone appendectomy at 23-years-old.

Personal and family history

The patient's other personal and family histories were unremarkable.

Physical examination

The patient was in good general condition (World Health Organization performance status of 0), with a body mass index of 21.55 kg/m². Clinical examination was normal.

Laboratory examinations


Biological investigations (blood count, liver function tests, glycemia, renal function, and C-reactive protein) were normal. The viral serologies for hepatitis B, hepatitis C, and human immunodeficiency virus were negative.

Imaging examinations

The thoracoabdominal computed tomography (CT) scan did not show any mediastinal lymph nodes in contact with the esophagus or other foci of tuberculosis.

ENDOSCOPIC EXAMINATION

Upper gastrointestinal (GI) endoscopy showed a jagged appearance of the thoracic esophageal mucosa for about 12 cm, stopping 3 cm above the cardia, with large irregular ulcers and raised contours. Nodules were present both at the level of the ulcers and in the normal-appearing mucosa (Figure 1A). Chromoendoscopy with narrow-band imaging did not detect areas that might suggest dysplasia or carcinoma (Figure 1B).



DOI: 10.4253/wjge.v14.i9.575 Copyright ©The Author(s) 2022.

Figure 1 Upper gastrointestinal endoscopy. A: Esophageal ulcer; B: Esophageal ulcer with nodules.

ANATOMICAL PATHOLOGY

Esophageal biopsies revealed a deep loss of wall tissue, reaching the muscularis mucosa. The normal tissue was replaced by granulation tissue containing a tuberculoid granuloma with several follicles consisting of epithelioid and multinucleated Langerhans histiocytes, surrounding a caseous necrosis (Figure 2). Neither culture of tissue samples nor PCR test for *Mycobacterium tuberculosis* were performed. Sputum and gastric acid liquid after aspiration were negative for acid-fast bacilli (AFB).

FINAL DIAGNOSIS

Isolated esophageal tuberculosis.

TREATMENT

An antituberculosis treatment was initiated [rifampicin, isoniazid, ethambutol, and pyrazinamide (RHEZ) and administered for 2 mo, and with rifampicin and isoniazid (RH) for 4 mo]. The patient showed good tolerance.

OUTCOME AND FOLLOW-UP

The patient's outcome was favorable, with a clear improvement of dysphagia after 15 d of treatment, which disappeared after 5 wk. Upper digestive endoscopy after 4 mo of treatment showed a normal esophageal mucosa. Six months after stopping the treatment, the patient was well, had regained weight, and did not complain of dysphagia.

DISCUSSION


Described for the first time in 1837 by Denonvilliers during an autopsy, infectious esophagitis due to tuberculosis is rare, even in countries with high tuberculosis endemicity. The esophageal localization represents 0.2%-1% of tuberculosis cases of the GI tract[1,2]. This low incidence can be explained by several mechanisms that allow the esophagus to fight infection, in particular, peristaltic movements leading to emptying of the contents into the stomach, and the presence of mucus and saliva lining the mucosa and its squamous epithelium[1]. These mechanisms provide a barrier against primary contamination caused by the ingestion of food or saliva containing germs such as *M. tuberculosis*. However, secondary contamination by contact with neighboring organs, especially in cases of tuberculosis in paraesophageal lymph nodes, is possible[3]. Blood-borne contamination is rare.

The most common symptom during esophageal tuberculosis is dysphagia (90% of cases), which was the main sign in our patient. Odynophagia, pyrosis, and chest pain may also be present[4]. The occurrence of coughing at mealtime should raise suspicion of an esotracheal or esophageal-mediastinal fistula, which is present in 13%-50% of cases[5]. The presence of hematemesis can also provide further evidence of a fistula[6].



DOI: 10.4253/wjge.v14.i9.575 Copyright ©The Author(s) 2022.

Figure 2 Esophageal biopsies. Esophageal ulcer detected in narrow band imaging.



DOI: 10.4253/wjge.v14.i9.575 Copyright ©The Author(s) 2022.

Figure 3 Granuloma with caseous necrosis (hematoxylin-eosin: 10 ×).

The endoscopic appearance of esophageal tuberculosis is variable and nonspecific. In our patient, the lesion was located in the lower two-thirds of the esophagus and consisted of a large ulcer with raised contours, associated with micronodules. The esophagus can be affected throughout its length, although the lesion is most often located in the middle third[3,7,8], because of the extensive lymphoid tissue in this region. Endoscopy may show an ulcer of variable size, superficial with regular contours or irregular and infiltrative simulating neoplasia, or show a more or less ulcerated budding aspect of the mucosa[3, 9]. An extrinsic compression aspect with a mucosa of normal appearance can also be found[8]. Endoscopic ultrasound can be helpful for diagnosis, allow analysis of the thick esophageal wall, and guide biopsies[7]. It also allows for exploration of the mediastinum and performance of fine-needle biopsy of potentially involved lymph nodes[7]. Thoracic CT scan often shows a thickening of the esophageal wall and allows for searching of adjacent lymph nodes, pulmonary location, or esophageal or esophago-mediastinal fistulas.

Histology can help in the diagnosis of esophageal tuberculosis. Mucosal biopsies during upper GI endoscopy can show the presence of a tuberculous granuloma or AFB in about 50% of cases[10,11], but sometimes neither of these lesions is found[12]. In our patient, an epithelioid gigantocellular granuloma with caseous necrosis was present on histology (Figure 3), confirming the diagnosis of esophageal tuberculosis. To improve diagnostic success, deep biopsy samples should be taken from ulcerated areas, as granulomas are most often found in the submucosa[1,8,11]. If endoscopic biopsies are not contributive, deep esophageal biopsy or fine-needle aspiration of a satellite lymph node, guided by endoscopic ultrasound, make it possible to find an epithelioid granuloma on histology (reportedly in 94.7% to 100% of cases, with caseous necrosis and/or AFB present in 55% to 75% of those cases)[7,11]. Histological samples are also used for PCR or culturing methods to identify *M. tuberculosis*. If an epithelioid granuloma without caseous necrosis is present, a differential diagnosis with sarcoidosis, Crohn's disease, or a carcinoma must be considered.

The treatment of esophageal tuberculosis is essentially medical, according to the standard protocol (rifampicin, isoniazid, ethambutol, and pyrazinamide daily for 2 mo, followed by rifampicin and isoniazid daily for 4 mo) for at least 6 mo. However, the optimal duration is not clinically established. In the case of fistula, clips are the reference treatment for lesion closure[11,13]. The outcome during treatment for esophageal tuberculosis is favorable and without sequelae in almost all cases[3,7,8,11]. In

our patient, no sequelae were noted during the follow-up. Upper digestive endoscopy, 4 mo after the beginning of treatment, was normal. The patient had no complaints at 6 mo after the end of treatment.

CONCLUSION

Esophageal tuberculosis is a rare cause of infectious esophagitis, even in a country where tuberculosis is endemic. Nevertheless, esophageal tuberculosis should be considered when dysphagia is associated with atypical ulcerated lesions of the esophageal mucosa. The presence of gigantocellular epithelioid granulomas on esophageal biopsies confirms the diagnosis. The patient's outcome is generally favorable after antibiologic treatment, as illustrated by our observation.

FOOTNOTES

Author contributions: Diallo I performed the upper digestive endoscopy, followed up with the patient, and wrote the manuscript; Touré O, Sow A, and Ndiaye B contributed to collecting the patient's clinical data, and participated in the follow-up; Sarr ES and Dial CM conducted the anatomopathological examinations; Diawara PS conducted the biological tests; Mbengue A performed the radiological examinations; Fall F supervised the manuscript; all authors have read and approved the final manuscript.

Informed consent statement: Consent was obtained from the patient.

Conflict-of-interest statement: All authors report no relevant conflict of interest for this article.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Senegal

ORCID number: Ibrahima Diallo [0000-0001-5635-0665](https://orcid.org/0000-0001-5635-0665).

S-Editor: Wu YXJ

L-Editor: A

P-Editor: Wu YXJ

REFERENCES

- 1 **Diallo I**, Omar Soko T, Rajack Ndiaye A, Klotz F. Tuberculose abdominale. *EMC - Gastro-entérologie* 2019; **37**: 1-13 [DOI: [10.1016/S1155-1968\(19\)92375-3](https://doi.org/10.1016/S1155-1968(19)92375-3)]
- 2 **Debi U**, Ravisankar V, Prasad KK, Sinha SK, Sharma AK. Abdominal tuberculosis of the gastrointestinal tract: revisited. *World J Gastroenterol* 2014; **20**: 14831-14840 [PMID: [25356043](https://pubmed.ncbi.nlm.nih.gov/25356043/) DOI: [10.3748/wjg.v20.i40.14831](https://doi.org/10.3748/wjg.v20.i40.14831)]
- 3 **Zhu R**, Bai Y, Zhou Y, Fang X, Zhao K, Tuo B, Wu H. EUS in the diagnosis of pathologically undiagnosed esophageal tuberculosis. *BMC Gastroenterol* 2020; **20**: 291 [PMID: [32859167](https://pubmed.ncbi.nlm.nih.gov/32859167/) DOI: [10.1186/s12876-020-01432-7](https://doi.org/10.1186/s12876-020-01432-7)]
- 4 **Vahid B**, Huda N, Esmaili A. An unusual case of dysphagia and chest pain in a non-HIV patient: esophageal tuberculosis. *Am J Med* 2007; **120**: e1-e2 [PMID: [17398209](https://pubmed.ncbi.nlm.nih.gov/17398209/) DOI: [10.1016/j.amjmed.2005.12.026](https://doi.org/10.1016/j.amjmed.2005.12.026)]
- 5 **Nagi B**, Lal A, Kochhar R, Bhasin DK, Gulati M, Suri S, Singh K. Imaging of esophageal tuberculosis: a review of 23 cases. *Acta Radiol* 2003; **44**: 329-333 [PMID: [12752007](https://pubmed.ncbi.nlm.nih.gov/12752007/) DOI: [10.1034/j.1600-0455.2003.00069.x](https://doi.org/10.1034/j.1600-0455.2003.00069.x)]
- 6 **Jain SS**, Somani PO, Mahey RC, Shah DK, Contractor QQ, Rathi PM. Esophageal tuberculosis presenting with hematemesis. *World J Gastrointest Endosc* 2013; **5**: 581-583 [PMID: [24255751](https://pubmed.ncbi.nlm.nih.gov/24255751/) DOI: [10.4253/wjge.v5.i11.581](https://doi.org/10.4253/wjge.v5.i11.581)]
- 7 **Tang Y**, Shi W, Sun X, Xi W. Endoscopic ultrasound in diagnosis of esophageal tuberculosis: 10-year experience at a tertiary care center. *Dis Esophagus* 2017; **30**: 1-6 [PMID: [28575247](https://pubmed.ncbi.nlm.nih.gov/28575247/) DOI: [10.1093/dote/dox031](https://doi.org/10.1093/dote/dox031)]
- 8 **Xiong J**, Guo W, Guo Y, Gong L, Liu S. Clinical and endoscopic features of esophageal tuberculosis: a 20-year retrospective study. *Scand J Gastroenterol* 2020; **55**: 1200-1204 [PMID: [32881605](https://pubmed.ncbi.nlm.nih.gov/32881605/) DOI: [10.1080/00365521.2020.1813799](https://doi.org/10.1080/00365521.2020.1813799)]
- 9 **Seo JH**, Kim GH, Jhi JH, Park YJ, Jang YS, Lee BE, Song GA. Endosonographic features of esophageal tuberculosis presenting as a subepithelial lesion. *J Dig Dis* 2017; **18**: 185-188 [PMID: [28139030](https://pubmed.ncbi.nlm.nih.gov/28139030/) DOI: [10.1111/1751-2980.12454](https://doi.org/10.1111/1751-2980.12454)]
- 10 **Park JH**, Kim SU, Sohn JW, Chung IK, Jung MK, Jeon SW, Kim SK. Endoscopic findings and clinical features of esophageal tuberculosis. *Scand J Gastroenterol* 2010; **45**: 1269-1272 [PMID: [20568972](https://pubmed.ncbi.nlm.nih.gov/20568972/) DOI: [10.1111/j.1365-3113.2010.04401.x](https://doi.org/10.1111/j.1365-3113.2010.04401.x)]

[10.3109/00365521.2010.501524](https://doi.org/10.3109/00365521.2010.501524)]

- 11 **Dahale AS**, Kumar A, Srivastava S, Varakanahalli S, Sachdeva S, Puri AS. Esophageal tuberculosis: Uncommon of common. *JGH Open* 2018; **2**: 34-38 [PMID: [30483561](https://pubmed.ncbi.nlm.nih.gov/30483561/) DOI: [10.1002/jgh3.12043](https://doi.org/10.1002/jgh3.12043)]
- 12 **Rana SS**, Bhasin DK, Rao C, Srinivasan R, Singh K. Tuberculosis presenting as Dysphagia: clinical, endoscopic, radiological and endosonographic features. *Endosc Ultrasound* 2013; **2**: 92-95 [PMID: [24949371](https://pubmed.ncbi.nlm.nih.gov/24949371/) DOI: [10.4103/2303-9027.117693](https://doi.org/10.4103/2303-9027.117693)]
- 13 **Rana SS**, Mandavdhare H, Sharma V, Sharma R, Dhalaria L, Bhatia A, Gupta R, Dutta U. Successful closure of chronic, nonhealing tubercular esophagobronchial fistula with an over-the-scope clip. *J Dig Endosc* 2017; **8** (1) : 33-5. [DOI: [10.4103/0976-5042.202820](https://doi.org/10.4103/0976-5042.202820)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

