# World Journal of *Clinical Cases*

World J Clin Cases 2022 July 26; 10(21): 7187-7619





Published by Baishideng Publishing Group Inc

W J C C World Journal of Clinical Cases

# Contents

# Thrice Monthly Volume 10 Number 21 July 26, 2022

# **OPINION REVIEW**

7187 Effects of glucocorticoids on leukocytes: Genomic and non-genomic mechanisms Jia WY, Zhang JJ

## **MINIREVIEWS**

- 7195 Apheresis: A cell-based therapeutic tool for the inflammatory bowel disease Yasmin F, Najeeb H, Naeem U, Moeed A, Koritala T, Surani S
- 7209 Helicobacter pylori infection and small intestinal bacterial overgrowth-more than what meets the eye Dharan M, Wozny D
- 7215 Anatomy of the anterolateral ligament of the knee joint Park JG, Han SB, Rhim HC, Jeon OH, Jang KM

## **ORIGINAL ARTICLE**

#### **Clinical and Translational Research**

7224 Molecular mechanisms of Biyu decoction as treatment for psoriasis: A network pharmacology and molecular docking study

Wang Z, Zhang HM, Guo YR, Li LL

7242 Expression of hepatocyte nuclear factor 4 alpha, wingless-related integration site, and  $\beta$ -catenin in clinical gastric cancer

Hu Q, Li LL, Peng Z, Yi P

#### **Case Control Study**

Improved Pittsburgh Sleep Quality Index scores on first postoperative night achieved by propofol 7256 anesthesia in patients undergoing ambulatory gynecologic surgery

Hu CH, Chou WY

Efficacy of Guhong injection versus Butylphthalide injection for mild ischemic stroke: A multicenter 7265 controlled study

Zhang WW, Xin J, Zhang GY, Zhai QJ, Zhang HM, Wu CS

#### **Retrospective Study**

7275 Clinical values of Barcelona Clinic Liver Cancer subgroup and up-to-7 criteria in intermediate stage hepatocellular carcinoma with transcatheter arterial chemoembolization

Lee SW, Peng YC, Lien HC, Ko CW, Tung CF, Chang CS

Intervention effect of encouraging mental and programmed nursing of patients in interventional operating 7285 room on their compliance and bad moods

Chi RB, Cai YY, Mao HP



Conton	World Journal of Clinical Cases
Conten	Thrice Monthly Volume 10 Number 21 July 26, 2022
7293	Preoperative neoadjuvant chemotherapy in patients with breast cancer evaluated using strain ultrasonic elastography
	Pan HY, Zhang Q, Wu WJ, Li X
7302	Risk factors for delayed intracranial hemorrhage secondary to ventriculoperitoneal shunt: A retrospective study
	Chen JC, Duan SX, Xue ZB, Yang SY, Li Y, Lai RL, Tan DH
7314	Sequential treatment of severe pneumonia with respiratory failure and its influence on respiratory mechanical parameters and hemodynamics
	Niu BY, Wang G, Li B, Zhen GS, Weng YB
7324	Effects of alendronate sodium combined with InterTan on osteoporotic femoral intertrochanteric fractures and fracture recurrence
	Wang KM, Wei SP, Yin XY, Meng QJ, Kong YM
7333	Correlation of magnetic resonance imaging quantitative parameters and apparent diffusion coefficient value with pathological breast cancer
	Wang Z, Ren GY, Yin Q, Wang Q
7341	Risk factors for delirium after surgery for craniocerebral injury in the neurosurgical intensive care unit
	Chen RY, Zhong CH, Chen W, Lin M, Feng CF, Chen CN
	Observational Study
7348	Effect of osteoarthritic knee flexion deformity correction by total knee arthroplasty on sagittal spinopelvic alignment in Indian population
	Puthiyapura LK, Jain M, Tripathy SK, Puliappadamb HM
7356	Imaging characteristics of orbital peripheral nerve sheath tumors: Analysis of 34 cases
	Dai M, Wang T, Wang JM, Fang LP, Zhao Y, Thakur A, Wang D
	Randomized Controlled Trial
7365	Comparison of involved-field intensity-modulated radiotherapy combined with S-1 vs radiotherapy alone for elderly patients with esophageal cancer
	Liu LH, Yan MH, Di YP, Fu ZG, Zhang XD, Li HQ
	Randomized Clinical Trial
7376	Dexmededomidine in pediatric unilateral internal inguinal ring ligation
	Liu G, Zhang L, Wang HS, Lin Y, Jin HQ, Wang XD, Qiao WN, Zhang YT, Sun JQ, Liu ZN
	META-ANALYSIS
7386	Impact of cancer on mortality rates in patients with sepsis: A meta-analysis and meta-regression of current studies
	Xiang MJ, Chen GL



Contents

# Thrice Monthly Volume 10 Number 21 July 26, 2022

# **CASE REPORT**

7397	Updated clinical and glycomic features of mannosyl-oligosaccharide glucosidase deficiency: Two case reports
	Abuduxikuer K, Wang L, Zou L, Cao CY, Yu L, Guo HM, Liang XM, Wang JS, Chen L
7409	Solitary necrotic nodules of the liver with "ring"-like calcification: A case report
	Bao JP, Tian H, Wang HC, Wang CC, Li B
7415	Corticosteroid-induced bradycardia in multiple sclerosis and maturity-onset diabetes of the young due to hepatocyte nuclear factor 4-alpha mutation: A case report
	Sohn SY, Kim SY, Joo IS
7422	Essential thrombocythemia with non-ST-segment elevation myocardial infarction as the first manifestation: A case report
	Wang ZM, Chen WH, Wu YM, Wang LQ, Ye FL, Yin RL
7429	Extranasopharyngeal angiofibroma in children: A case report
	Yan YY, Lai C, Wu L, Fu Y
7438	Deep Sylvian fissure meningiomas: A case report
	Wang A, Zhang X, Sun KK, Li C, Song ZM, Sun T, Wang F
7445	Acute pulmonary embolism originating from upper limb venous thrombosis following breast cancer surgery: Two case reports
	Duan Y, Wang GL, Guo X, Yang LL, Tian FG
7451	Managing spondylitis tuberculosis in a patient with underlying diabetes and hypothyroidism: A case report
	Novita BD, Muliono AC, Wijaya S, Theodora I, Tjahjono Y, Supit VD, Willianto VM
7459	Ovarian mucinous tumor with mural nodules of anaplastic carcinoma: Three case reports
	Wang XJ, Wang CY, Xi YF, Bu P, Wang P
7467	Transcatheter arterial infusion chemotherapy and embolization for primary lacrimal sac squamous cell carcinoma: A case report
	Sun MH, Yi WD, Shen L, Zhou L, Lu JX
7474	Programmed cell death-1 inhibitor combination treatment for recurrent proficient mismatch repair/ miscrosatellite-stable type endometrial cancer: A case report
	Zhai CY, Yin LX, Han WD
7483	Novel compound heterozygous mutation of <i>SLC12A3</i> in Gitelman syndrome co-existent with hyperthyroidism: A case report and literature review
	Qin YZ, Liu YM, Wang Y, You C, Li LN, Zhou XY, Lv WM, Hong SH, Xiao LX
7495	Successful treatment of hyperglycemia with liraglutide in a hospitalized 27-year-old patient with schizophrenia: A case report

Zhang L, Yu WJ, Zhu H, Li HF, Qiao J



Conton	World Journal of Clinical Cases
Conten	Thrice Monthly Volume 10 Number 21 July 26, 2022
7502	Refractory lymphoma treated with chimeric antigen receptor T cells combined with programmed cell death-1 inhibitor: A case report
	Zhang CJ, Zhang JY, Li LJ, Xu NW
7509	Median arcuate ligament syndrome with retroperitoneal haemorrhage: A case report
	Lu XC, Pei JG, Xie GH, Li YY, Han HM
7517	Novel frameshift mutation in the <i>AHDC1</i> gene in a Chinese global developmental delay patient: A case report
	Lin SZ, Xie HY, Qu YL, Gao W, Wang WQ, Li JY, Feng XC, Jin CQ
7523	Selective nerve block for the treatment of neuralgia in Kummell's disease: A case report
	Zhang X, Li ZX, Yin LJ, Chen H
7531	Traditional Chinese medicine manipulative reduction combined with percutaneous vertebroplasty for treating type III Kummell's disease: A case report
	Hao SS, Zhang RJ, Dong SL, Li HK, Liu S, Li RF, Ren HH, Zhang LY
7539	Differential diagnosis and treatment of foot drop caused by an extraneural ganglion cyst above the knee: A case report
	Won KH, Kang EY
7545	Effect of hydrogen intervention on refractory wounds after radiotherapy: A case report
	Zhao PX, Luo RL, Dang Z, Wang YB, Zhang XJ, Liu ZY, Wen XH, Liu MY, Zhang MZ, Adzavon YM, Ma XM
7553	Chronic urticaria associated with lung adenocarcinoma – a paraneoplastic manifestation: A case report and literature review
	Jiménez LF, Castellón EA, Marenco JD, Mejía JM, Rojas CA, Jiménez FT, Coronell L, Osorio-Llanes E, Mendoza-Torres E
7565	Spinal giant cell-rich osteosarcoma-diagnostic dilemma and treatment strategy: A case report
	Tseng CS, Wong CE, Huang CC, Hsu HH, Lee JS, Lee PH
7571	Primary clear cell sarcoma of soft tissue in the posterior cervical spine invading the medulla oblongata: A case report
	Liu CC, Huang WP, Gao JB
7577	<i>Pseudomonas aeruginosa</i> -related effusive-constrictive pericarditis diagnosed with echocardiography: A case report
	Chen JL, Mei DE, Yu CG, Zhao ZY
7585	Maternal peripartum bacteremia caused by intrauterine infection with Comamonas kerstersii: A case report
	Qu H, Zhao YH, Zhu WM, Liu L, Zhu M
7592	Considerations of single-lung ventilation in neonatal thoracoscopic surgery with cardiac arrest caused by bilateral pneumothorax: A case report
	Zhang X, Song HC, Wang KL, Ren YY



World Journal of Clinical Cases Contents Thrice Monthly Volume 10 Number 21 July 26, 2022 7599 Rare primary rectal mucosa-associated lymphoid tissue lymphoma with curative resection by endoscopic submucosal dissection: A case report and review of literature Tao Y, Nan Q, Lei Z, Miao YL, Niu JK Differences in examination results of small anastomotic fistula after radical gastrectomy with afterward 7609 treatments: A case report Lu CY, Liu YL, Liu KJ, Xu S, Yao HL, Li L, Guo ZS

# **LETTER TO THE EDITOR**

7617 Baseline differences may impact on relationship between dietary tryptophan and risk of obesity and type 2 diabetes

Ren XH, Ye YW, He LP



# Contents

Thrice Monthly Volume 10 Number 21 July 26, 2022

# **ABOUT COVER**

Editorial Board Member of World Journal of Clinical Cases, Rajesh Kumar Rajnish, MBBS, MS, Assistant Professor, Department of Orthopaedics, All India Institute of Medical Sciences, Bilaspur, Bilaspur 174001, Himachal Pradesh, India. duktiraj@gmail.com

# **AIMS AND SCOPE**

The primary aim of World Journal of Clinical Cases (WJCC, World J Clin Cases) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

# **INDEXING/ABSTRACTING**

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

# **RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Ying-Yi Yuan, Production Department Director: Xiang Li, Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL World Journal of Clinical Cases	INSTRUCTIONS TO AUTHORS https://www.wignet.com/bpg/gerinfo/204
<b>ISSN</b> ISSN 2307-8960 (online)	GUIDELINES FOR ETHICS DOCUMENTS https://www.wignet.com/bpg/GerInfo/287
LAUNCH DATE April 16, 2013	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH https://www.wignet.com/bpg/gerinfo/240
FREQUENCY Thrice Monthly	PUBLICATION ETHICS https://www.wignet.com/bpg/GerInfo/288
EDITORS-IN-CHIEF	PUBLICATION MISCONDUCT
Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku	https://www.wjgnet.com/bpg/gerinfo/208
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE
https://www.wjgnet.com/2307-8960/editorialboard.htm	https://www.wjgnet.com/bpg/gerinfo/242
PUBLICATION DATE	STEPS FOR SUBMITTING MANUSCRIPTS
July 26, 2022	https://www.wjgnet.com/bpg/GerInfo/239
COPYRIGHT © 2022 Baishideng Publishing Group Inc	ONLINE SUBMISSION https://www.f6publishing.com

© 2022 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: bpgoffice@wjgnet.com https://www.wjgnet.com



W J C C World Journal of Clinical Cases

Submit a Manuscript: https://www.f6publishing.com

World J Clin Cases 2022 July 26; 10(21): 7242-7255

DOI: 10.12998/wjcc.v10.i21.7242

ISSN 2307-8960 (online)

ORIGINAL ARTICLE

# **Clinical and Translational Research**

# Expression of hepatocyte nuclear factor 4 alpha, wingless-related integration site, and $\beta$ -catenin in clinical gastric cancer

Qian Hu, Ling-Li Li, Ze Peng, Ping Yi

Specialty type: Oncology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

# Peer-review report's scientific quality classification

Grade A (Excellent): 0 Grade B (Very good): B, B Grade C (Good): C Grade D (Fair): D Grade E (Poor): 0

P-Reviewer: Martin-Villa JM, Spain; Rojas A, Chile; Shamseldeen AA, Egypt; Tanabe S, Japan A-Editor: Zhu JQ, China

Received: March 25, 2022 Peer-review started: March 25, 2022 First decision: April 13, 2022 Revised: April 17, 2022 Accepted: June 3, 2022 Article in press: June 3, 2022 Published online: July 26, 2022



Qian Hu, Ling-Li Li, Ze Peng, Ping Yi, Department of Integrated Traditional Chinese and Western Medicine, Institute of Integrated Traditional Chinese and Western Medicine, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430030, Hubei Province, China

Corresponding author: Ping Yi, MD, PhD, Doctor, Department of Integrated Traditional Chinese and Western Medicine, Institute of Integrated Traditional Chinese and Western Medicine, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, No. 1095 Jie Fang Avenue, Wuhan 430030, Hubei Province, China. pyi219@163.com

# Abstract

# BACKGROUND

Gastric cancer (GC) is the second most common cause of cancer-related deaths worldwide. Hepatocyte nuclear factor 4 alpha (HNF4 $\alpha$ ) that belongs to the nuclear hormone receptor superfamily, is overexpressed in GC tissues, and might be involved in the development of GC by regulating its downstream winglessrelated integration site (WNT)/ $\beta$ -catenin signaling.

# AIM

To clarify the expression of HNF4 $\alpha$ /WNT5a/ $\beta$ -catenin signaling proteins in clinical GC tissues.

# **METHODS**

We immunohistochemically stained pathological blocks of GC and matched paracancerous tissues. The intensity of HNF4 $\alpha$ , WNT5a and  $\beta$ -catenin staining in the tumor cells was determined according to cell rates and staining intensity. The correlations between GC and HNF4 $\alpha$ , WNT5a, and  $\beta$ -catenin expression using chisquare and paired chi-square tests. Relationships between double-positive HNF4α and WNT5a expression and types of gastric tumor tissues were assessed using regression analysis. Correlations between HNF4a and WNT5a expression at the RNA level in GC tissues found in the TCGA database were analyzed using Pearson correlation coefficients.

# RESULTS

We found more abundant HNF4a and WNT5a proteins in GC, especially in mucinous adenocarcinoma and mixed GC than in adjacent tissues (P < 0.001). Low and high levels of cytoplasmic  $\beta$ -catenin respectively expressed in GC and



adjacent tissues (P < 0.001) were not significantly associated with pathological parameters.

#### CONCLUSION

The expressions of HNF4 $\alpha$  and WNT5a could serve as early diagnostic biomarkers for GC.

Key Words:  $\beta$ -catenin; Biomarker; Gastric cancer; Hepatocyte nuclear factor 4 alpha; Wingless-related integration site

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Gastric cancer (GC) is the second most common cause of cancer-related deaths worldwide. A precise biopsy biomarker that can predict the progression of GC and change its prognosis is urgently needed. We previously found that hepatocyte nuclear factor 4 alpha (HNF4a)/wingless-related integration site (WNT) signaling is involved in the process of GC in preclinical models. Here, we obtained gastric tumor and para-cancerous tissue sections from Tongji Hospital and investigated the expression of HNF4 $\alpha$ , WNT5a, and  $\beta$ -catenin. The expression of HNF4 $\alpha$  and WNT5a could serve as an early diagnostic biomarker of GC.

Citation: Hu Q, Li LL, Peng Z, Yi P. Expression of hepatocyte nuclear factor 4 alpha, wingless-related integration site, and β-catenin in clinical gastric cancer. World J Clin Cases 2022; 10(21): 7242-7255 URL: https://www.wjgnet.com/2307-8960/full/v10/i21/7242.htm DOI: https://dx.doi.org/10.12998/wjcc.v10.i21.7242

# INTRODUCTION

Gastric cancer (GC) is the second most common cause of cancer-related deaths worldwide. Because early symptoms are not obvious, most GCs are diagnosed at a late stage when a radical cure is largely ineffective[1,2]. Thus, to diagnose the disease at an early stage when the treatment is likely to be more effective is critical. Several screening approaches have been proposed, such as serum pepsinogens, serum ghrelin and Gastrin-17. However, further investigation is needed to determine the overall effectiveness of these approaches[3]. Upper gastrointestinal endoscopy is the most sensitive and specific diagnostic screening method, and biopsy tissues can be morphologically examined [4,5]. However, endoscopic diagnosis of early GC is quite challenging because it requires advanced endoscopic techniques, and expert endoscopists who are familiar with current and emerging techniques[4]. Therefore, accurate biopsy biomarkers are needed that can predict the progression and prognosis of GC in para-cancerous or cancerous tissues. Identifying such markers would lead to the development of accurately targeted therapy for GC.

Hepatocyte nuclear factor 4 alpha (HNF4a) is an endoderm-specific zinc-finger transcription factor that has a highly conserved orphan receptor and belongs to the nuclear hormone receptor superfamily [6]. It is found in the liver, kidney, pancreas, stomach, small intestine, and colon[7] and is associated with several types of cancer. The abundance of HNF4 $\alpha$  varies in different types of cancer and even reverses. For example, HNF4 $\alpha$  is expressed at lower levels in colon carcinoma and hepatocellular carcinoma tissues than in adjacent normal tissues [6,8]. In contrast, it is overexpressed in gastric tumor tissues compared to adjacent normal tissues and associated with a poorer prognosis in patients with GC [9]. Moreover, protein-protein interaction networks have revealed that HNF4 $\alpha$  is the most significant node in GC and might be a useful diagnostic marker and therapeutic target [10]. However, a relationship between HNF4 $\alpha$  and the pathological classification or demographic characteristics of GC has not been identified.

Wingless-related integration site (WNT) signaling pathways are deregulated in cancers[11]. Nam et al [12] applied PATHOME, a novel algorithm that can sensitively detect expressed pathways in GC gene expression datasets. They found that WNT pathway are involved in the development of primary GC and that WNT5a is a potential target gene[12,13]. A meta-analysis similarly associated WNT5a expression in human GC with aggressiveness and poor prognosis[14]. Anti-WNT5a antibodies can suppress GC metastasis in vivo[15]. Moreover, WNT5a is closely associated with the epithelialmesenchymal transition (EMT) and participates in GC development by promoting the EMT[16,17].

Hepatocyte nuclear factor 4 alpha is a transcription factor that regulates the transcription of WNT5a by competing with  $\beta$ -catenin for binding to transcription factor 4 (TCF4) and inhibiting WNT/ $\beta$ -catenin signaling in hepatocellular carcinoma cells[6,12,18]. A double-negative feedback mechanism controls WNT/ $\beta$ -catenin signaling and HNF4 $\alpha$  expression during EMT regulation, which eventually affects hepatocellular carcinoma development[19]. We previously found that an HNF4 $\alpha$ /WNT5a/ $\beta$ -catenin



signaling axis in GC cell lines and animal models<sup>[20,21]</sup>, and affects the growth and development of GC [12]. However, the expression of the HNF4 $\alpha$ /WNT5a/ $\beta$ -catenin signaling pathway in clinical GC tissues remains unclear. Thus, we investigated the relationship between this pathway and GC in a preclinical model of GC tissues.

In this study, 158 cases of gastric tumoral tissues and 164 cases of adjacent para-cancerous tissues in Tongji hospital were selected. We immunohistochemically stained HNF4 $\alpha$ , WNT5a, and  $\beta$ -catenin expression in these tissues. And then we explored relationships between these molecules and the clinicopathological features of GC as well as the sociological features of patients. This is an extension of a series of studies.

# MATERIALS AND METHODS

#### Sample collection and characterization

Human GC tissues were obtained from patients who presented between 2016 and 2018 at Tongji Hospital of Tongji Medical College of Huazhong University of Science and Technology (Wuhan, China). According to the WHO histological classifications, GC comprises signet ring cell carcinoma, and tubular, as well as mucinous adenocarcinoma. We initially selected wax blocks of paraffin-embedded GC and matched para-cancerous tissue blocks (n = 164 each) to reduce the influence of pathological WHO classification of GC on the expression of HNF4 $\alpha$ , WNT5 $\alpha$ , and  $\beta$ -catenin in GC tissues. However, six small gastric tumor tissues of insufficient quality were excluded. Therefore, we analyzed 158 GC (59 signet-ring cell carcinoma, and 35 tubular, and 64 mucinous adenocarcinomas) and 164 para-cancerous tissues. The GC tissues comprised 97, 46, and 15 samples of intestinal, diffuse mixed types according to the Lauren classification. Pathologists from the Department of Pathology at Tongji Hospital determined the histological classification. The study was approved by the Ethics Committee of Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology (Approval no: TJ-IRB20190501).

#### Immunohistochemistry staining

The paraffin-embedded tissues were cut into 4-µm-thick sections and mounted on triplicate APEScoated slides (AR0001; Boster Biological Technology Co., Ltd., Wuhan, China). The tissues were deparaffinized in xylene (Hubei Jing-Hengye Technology Co., Ltd., Wuhan, China) and rehydrated in graded ethanol (National Medicine Group Chemical Reagent Co., Ltd.). Endogenous peroxidase activity was quenched with a 3% hydrogen peroxide in methanol (National Medicine Group Chemical Reagent Co., Ltd.) at room temperature for 30 min, then the tissues were rinsed in phosphate-buffered saline (PBS) at pH 6.0 after antigen retrieval in 10 mmol/L citrate buffer (pH 6.0) at 94°C for 8 and 10 min, the slides were immediately cooled for 20 min at room temperature. Nonspecific antigen binding sites were blocked by incubation with wash buffer containing 10% normal goat serum (YJ0130; Shanghai Yanjin Biology Science and Technology Co., Ltd., Shanghai, China) at 37°C for 30 min. The sections were then incubated overnight at 4°C with primary antibodies against HNF4α diluted 1:200 (#3113; Cell Signaling Technologies, Danvers, MA, United States), WNT5a diluted 1:80 (abs113167; Abgent, San Diego, CA, United States), and β-catenin diluted 1:100 (#8480; Cell Signaling Technologies). Positive tissues were stained brown with the peroxidase substrate 3-3'-diaminobenzidine tetrahydrochloride (DAB) (Hubei Biossci Biological Co., Ltd., Wuhan, China) and all tissues were lightly counterstained with hematoxylin. Specimens were examined under a light microscope (200× magnification).

#### Interpretation of immunohistochemical findings

The intensity of HNF4 $\alpha$ , WNT5a and  $\beta$ -catenin staining in the tumor cells was determined using a single-blind method. The samples were scored as 0, +1 or +2 according to cell rates, and from 0 to +3 according to staining intensity<sup>[22]</sup>. Cell positivity scores: 0, weakly positive (< 10% of tumor cells emitted faint signals); +1, moderately positive (10%-50% of tumor cells emitted clear signals); +2 strongly positive (> 50% of tumor cells emitted signals). Staining intensity of tumor cells was scored as 0 (negative), +1, +2 or +3 (weakly, moderately, and strongly positive, respectively). The product of the positive cell rate and staining intensity generated the immunohistochemical score, where 0-1 was negative (-), and 2-3, 4, and 6 were mildly (+), moderately positive (++), and strongly (+++) positive, respectively.

#### Statistical analysis

All data were analyzed using SPSS v. 20.0 (IBM Corp, Armonk, NY, United States). Relationships between the expression of HNF4 $\alpha$ , WNT5a, and  $\beta$ -catenin and the pathological features of GC as well as the sociological features of the patients were analyzed using chi-squared tests. Correlations among HNF4 $\alpha$ , WNT5a, and  $\beta$ -catenin expression were evaluated using chi-squared tests of paired comparisons. Relationships between double-positive HNF4 $\alpha$  and WNT5a expression and types of gastric tumor tissues were assessed using regression analysis. Correlations between  $HNF4\alpha$  and WNT5a



WJCC | https://www.wjgnet.com

expression at the RNA level in GC tissues found in the TCGA database were analyzed using Pearson correlation coefficients. Values with two-tailed P < 0.05 were considered statistically significant.

# RESULTS

#### Demographic characteristics of patients and pathological characteristics of GC sections

Among the 158 GC tissues from 101 men and 57 women, 97, 52 and 9 were poorly, moderately, and highly differentiated, respectively. The GC tissues were staged according to the depth of invasion (T), lymph node (N), and distant metastasis (M) (TNM classification). We staged 28, 38, 74, and 18 GC tissues as I, II, III, and IV, respectively (Table 1).

#### Expression of HNF4a in gastric tumoral tissues and para-cancerous tissues

Figure 1A and B shows representative HNF4 $\alpha$  expression in different grades of GC and para-cancerous tissues. The positive rates of HNF4 $\alpha$  in GC and para-cancerous tissues were 145 (91.7%) of 158 *vs.* 16 (9.8%) of 164 (P < 0.001 Table 2).

#### Expression of WNT5a in gastric tumoral tissues and para-cancerous tissues

As a ligand for the Frizzled family of atypical G protein-coupled receptors, WNT5a plays a critical role in postnatal cellular functions and development[14]. Both HNF4 $\alpha$  and WNT5a, which is a downstream signaling molecule and effector of HNF4 $\alpha$ , are involved in GC development. Therefore, we analyzed WNT5a expression in clinical samples[18]. Figure 2 shows WNT5a expression in different grades of gastric tumor and para-cancerous tissues, and cytoplasmic staining. The positive rate of WNT5a expression was 149 (94.3%) of 158 (Table 3) in GC, and significantly decreased in para-cancerous tissues (P < 0.001).

#### Relationships between HNF4a, WNT5a expression and different types of GC

The WHO histologically classifies GC as tubular, signet ring cell, and mucinous adenocarcinoma. Table 4 shows that HNF4 $\alpha$  expression was abundant in tubular and mucinous adenocarcinomas and relatively weak in signet ring cell carcinoma (P < 0.001). Conversely, WNT5a expression was more abundant in mucinous, than in tubular adenocarcinoma and signet ring cell carcinoma (P < 0.001). The results of regression analyses showed that tubular GC was more likely to moderately express positive WNT5a (Supplementary Table 1).

The Lauren classification also categorizes GC into intestinal, diffuse, and mixed types. Compared with intestinal and diffuse GC, both HNF4 $\alpha$  and WNT5a were strongly positive in mixed GC (Table 5, *P* < 0.001), but overall, both were abundantly expressed in mucinous and mixed GCs.

# Co-expression of HNF4α and WNT5a in gastric tumoral tissues and correlation with the clinicopathological characteristics

We examined pairwise co-expression associations between HNF4 $\alpha$  and WNT5a in clinical GC tissue specimens. The  $\chi^2$  test of paired comparisons found no positive correlations between HNF4 $\alpha$  and WNT5a expression in these tissues ( $\chi^2 = 1.5$ , P > 0.05; Table 6), although both were more abundantly expressed in GC, than para-cancerous tissues.

We analyzed the diagnostic accuracy, sensitivity, specificity, and positive and negative predictive rates of HNF4 $\alpha$  and WNT5a to further determine their potential roles in diagnosing GC. Table 7 shows that the diagnostic accuracy, sensitivity, specificity, and positive and negative predictive rates of HNF4 $\alpha$  were 91.0%, 91.8% 90.2%, 90.1% and 91.9%, respectively. In contrast, the diagnostic accuracy and specificity of WNT5a were relatively low at 85.7% and 77.4%, respectively, but the diagnostic sensitivity reached 94.3%. The positive and negative predictive rates of WNT5a expression were 80.1% and 93.4%, respectively. The expression of HNF4 $\alpha$  and WNT5a in GC tissues did not correlate with the patients' sex and age, or tumor differentiation, invasion depth, distant metastasis, and TNM stage (P > 0.05). In contrast, HNF4 $\alpha$  expression was associated with GC differentiation, and the positive rate was the highest among the moderately differentiated GC tissues (P < 0.05) (Supplementary Table 2).

#### β-catenin expression in gastric tumoral tissues and para-cancerous tissues

Excessive cytoplasmic  $\beta$ -catenin in the  $\beta$ -catenin-dependent WNT signaling pathway migrates to the nucleus where it acts as a transcription factor and activates a series of downstream signaling pathways. We analyzed whether the canonical  $\beta$ -catenin-dependent or the non-canonical  $\beta$ -catenin-independent WNT pathway participates in clinical gastric carcinogenesis. Figure 3 shows abundant membranous and cytoplasmic  $\beta$ -catenin expression in gastric tumoral and para-cancerous tissues, but the ratio of positive  $\beta$ -catenin expression was significantly lower in the latter 114 (72.2%) of 158 *vs.* 153 (93.3%) of 164 (P < 0.001; Table 8).

WJCC | https://www.wjgnet.com

Table 1 Demographic characteristics of patients and pathological characteristics of gastric cancer sections					
Groups	n				
Gender					
Male	101				
Female	57				
Age (yr)					
≤ 60	97				
> 60	61				
Differentiation					
Low	97				
Moderate	52				
High	9				
Invasion depth (T)					
T <sub>1</sub>	26				
T <sub>2</sub>	20				
T <sub>3</sub>	39				
T <sub>4</sub>	73				
Lymph node metastasis (N)					
N <sub>0</sub>	65				
N <sub>1</sub>	16				
N <sub>2</sub>	32				
N <sub>3</sub>	45				
Distant metastasis (M)					
$M_0$	126				
$M_1$	32				
TNM stage					
Ι	28				
П	38				
ш	74				
IV	18				

TNM: Tumor-node-metastasis.

We further analyzed correlations between  $\beta$ -catenin expression in the GC tissues and the clinical demographic features of the patients as well as the histopathological characteristics of GC. The expression of β-catenin was not significantly associated with sex, age, tumor differentiation, depth of invasion, distant metastasis, and TNM stage in these patients (P > 0.05; Supplementary Table 3).

# DISCUSSION

The nuclear transcription factor HNF4 $\alpha$  is expressed in the liver, kidney, and intestine[6,23]. It participates in glucose and lipid metabolism and insulin secretion, and is associated with metabolic diseases such as diabetes and obesity [24-26] as well as malignant tumor differentiation via gene regulation[27]. For example, Sugano *et al*[28] found that HNF4 $\alpha$  could serve as a marker for invasive mucinous adenocarcinoma of the lung, and Ma *et al*[9] found that HNF4 $\alpha$  is linked to GC caused by Helicobacter pylori. We previously showed that  $HNF4\alpha$  can regulate the proliferation, invasion, and metastasis of GC in vitro and in vivo by modulating the tumorigenic WNT signaling pathway[20]. The present study found that more nuclei stained with HNF4α in GC, than para-cancerous tissues had high



Baishidena® WJCC | https://www.wjgnet.com

Table 2 Expression of hepatocyte nuclear factor 4 alpha protein in gastric tumoral tissues and para-cancerous tissues							
Groups	n	HNF4 $\alpha$ expression				Duralua	
		Negative	Positive	Positive (%)	X	r value	
GC	158	13	145	91.7			
PC	164	148	16	9.8	216.5	0.000	

HNF4α: Hepatocyte nuclear factor 4 alpha; GC: Gastric cancer; PC: Para-cancerous.

#### Table 3 Expression of wingless-related integration site protein in gastric tumoral tissues and para-cancerous tissues

Groupo	n	WNT5a expression		Positivo (%)	v <sup>2</sup>	Byalua
Groups		Negative	Positive		X	rvalue
GC	158	9	149	94.3		
PC	164	127	37	22.6	169.8	0.000

WNT5A: Wingless-related integration site; GC: Gastric cancer; PC: Para-cancerous.

Table 4 Correlation of hepatocyte nuclear factor 4 alpha and wingless-related integration site expression with types of gastric carcinomas according to WHO classification

Groups	n	Grade of HNF4 $\alpha$ expression			Grade of WNT5a expression		
		+ (%)	++ (%)	+++ (%)	+ (%)	++ (%)	+++ (%)
Tubular	35	4 (11.4)	2 (5.7)	29 (82.7) <sup>b</sup>	2 (5.7)	11 (31.4)	22 (62.9)
Signet ring	59	5 (8.5)	11 (18.6)	43 (72.9)	4 (6.8)	17 (28.8)	38 (64.4)
Mucinous	64	4 (6.3)	7 (10.9)	53 (82.8) <sup>b</sup>	3 (4.7)	10 (15.6)	51 (79.7) <sup>d</sup>

 $^{b}P < 0.001$  hepatocyte nuclear factor 4 alpha ( $\chi^2$  = 83.909) vs signet ring;  $^{d}P < 0.001$  wingless-related integration site ( $\chi^2$  = 105.865) vs tubular and signet ring; HNF4a: Hepatocyte nuclear factor 4 alpha; WNT5A: Wingless-related integration site.

Table 5 Correlation of hepatocyte nuclear factor 4 alpha and wingless-related integration site expression with types of gastric carcinomas according to Lauren classification

Groups	n	Grade of HNF4α	expression	Grade of WNT5a expression			
		+ (%)	++ (%) ++	+ (%)	+ (%)	++ (%)	+++ (%)
Intestinal	97	11 (11.3)	15 (15.5)	71 (73.2)	7 (7.2)	23 (23.7)	67 (69.1)
Diffuse	46	2 (4.4)	5 (10.9)	39 (84.8)	2 (4.4)	14 (30.4)	30 (65.2)
Mixed	15	0 (0.00)	0 (0.00)	15 (100.0) <sup>b</sup>	0 (0.00)	1 (6.7)	14 (93.3) <sup>d</sup>

 $^{b}P < 0.001$  hepatocyte nuclear factor 4 alpha ( $\chi^2 = 79.404$ ) and  $^{d}P < 0.001$  wingless-related integration site ( $\chi^2 = 80.539$ ) vs intestinal and diffuse. HNF4 $\alpha$ : Hepatocyte nuclear factor 4 alpha; WNT5A: Wingless-related integration site.

> diagnostic accuracy, sensitivity, and specificity. Furthermore, HNF4a regulates organic acid metabolism, sustains oncogenic metabolism in GC, and promotes GC development by positively regulating the isocitrate dehydrogenase 1 (*IDH1*) gene[29]. Moreover, HNF4α enhances multidrug resistance by increasing GC cell apoptosis, thus suppressing therapeutic responses[30]. Therefore, HNF4 a can promote GC development by affecting the metabolism and multiple drug resistance of GC cells.

> Aberrant WNT signaling often leads to tumor development and WNT5a is a ligand for seven Frizzled family transmembrane receptors, most of which are associated with GC. We previously found significantly increased expression of WNT5a in SGC7901 and MGC803 GC cells and in mice with GC xenografts. Consistent with these results, WNT5a is abundantly expressed in GC tissues, weakly in para-cancerous tissues, and in the cytoplasm of clinical GC sections[12,31]. The selective effects of



Table 6 Positive correlation between hepatocyte nuclear factor 4 alpha and wingless-related integration site expressions in gastric cancer							
	HNF4α expression		Total		Duralua		
WIN I Sa	Negative	Positive	I otal	X	P value		
Negative	8	1	9	1.5	> 0.05		
Positive	5	144	149				
Total	13	145	158				

WNT5A: Wingless-related integration site; HNF4α: Hepatocyte nuclear factor 4 alpha.

Table 7 Evaluation of hepatocyte nuclear factor 4 alpha and wingless-related integration site in the diagnosis of gastric cancer								
	Diagnostic accurate rate	Sensitivity	Specificity	Positive predictive rate	Negative predictive rate			
HNF4α	91.0	91.8	90.2	90.1	91.9			
WNT5a	85.7	94.3	77.4	80.1	93.4			

HNF4α: Hepatocyte nuclear factor 4 alpha; WNT5a: Wingless-related integration site.

Table 8 Expression of β-catenin protein in gastric carcinoma and para-cancerous tissue								
Groups	n	β-catenin expression		Depitive (9/)		Dyelve		
Groups		Negative	Positive		X	r value		
GC	158	44	114	72.2				
PC	164	11	153	93.3	25.394	0.000		

GC: Gastric cancer: PC: Para-cancerous

WNT5a on cancer cells include the promoted invasion but not proliferation of KKLS GC cells, and the stimulation of both in A549 Lung cancer cells[32]. The progression of GC is promoted by WNT5a mainly by increasing the expression of metastatic proteins such as laminin gamma 2 and the EMT[16,33]. However, WNT5a induces the M2 polarization of tumor-associated macrophages through the CaKMII-ERK1/2-STAT3 pathway in CRC, thus promoting the proliferation, migration, and invasiveness of cancer cells[34]. Therefore, abundant WNT5a in gastric tumor tissues might contribute to GC metastasis and invasion by affecting EMT.

We previously found that the WNT5a/ $\beta$ -catenin signaling pathway is cell downstream of HNF4 $\alpha$  in GC cells, and the HNF4 $\alpha$ /WNT5a signaling affected GC development in GC cell lines and animal models [20,21]. The present study also found more abundant HNF4 $\alpha$  and WNT5a expression in GC, than that in adjacent tissues, especially in mucinous adenocarcinoma and mixed GC, and that both had high diagnostic accuracy. However, HNF4 $\alpha$  and WNT5a expression did not correlate in GC tissues. We similarly identified a weak correlation (0 < r = 0.11) between HNF4 $\alpha$  and WNT5a in GC tissues at RNA level in the Cancer Genome Atlas (TCGA, https://cancergenome.nih.gov/) (Supplementary Figure 1), whereas others have found that  $HNF4\alpha$  and WNT5a expression in GC positively correlates at the RNA and protein levels [12,18,20]. Other than the abundant HNF4 $\alpha$  expression in moderately differentiated GC, the present study found that neither HNF4 $\alpha$  nor WNT5a expression correlates with the demographic and pathological characteristics of GC of sex, age, tumor differentiation degree, invasion depth, distant metastasis and TNM stage.

The present study included tissues from patients who were treated between 2016 and 2018, and follow-up is currently underway. Therefore, we were unable to provide a complete (or 5-year) survival analysis of double positive or double negative HNF4 $\alpha$  and WNT5a in GC. However, we analyzed the progression-free (PFS), and overall (OS) survival of 447 patients with HNF4α and WNT5a double positive and negative GC in the TCGA database. We found that the although patients tended to survive longer if they were double negative than double positive, the differences in 5-year OS or PFS did not reach statistical significance (Supplementary Figures 2-5). A retrospective study found that a combination of traditional Chinese Medicine (TCM) and chemotherapy significantly prolongs survival and improves the quality of life among patients with GC compared with chemotherapy alone [35]. For



WJCC | https://www.wjgnet.com



Figure 1 Expression of hepatocyte nuclear factor 4 alpha in gastric cancer tissues and para-cancerous tissues. A, B: Representative figures of hepatocyte nuclear factor 4 alpha (HNF4a) expression (nuclear staining) at different grades of pathological scores in gastric cancer tissues (GC) (A) and paracancerous (PC) tissues (B); C: Numbers of samples expressing HNF4a in GC and PC groups. Specimens were examined under a light microscope (200 × magnification). HNF4: Hepatocyte nuclear factor 4 alpha.

example, berberine, an extract of Rhizoma Coptidis, improves the chemotherapy effect and inhibit GC development by regulating the HNF4 $\alpha$ /WNT5a signal axis[20].

The WNT/ $\beta$ -catenin pathway is crucial for tissue development and homeostasis in all animal, and its dysregulation is one of the most relevant events linked to cancer development and dissemination[36].

Baisbideng® WJCC https://www.wjgnet.com

Hu Q et al. HNF4 $\alpha$ /WNT5a/ $\beta$ -catenin in gastric cancer



Figure 2 Expression of wingless-related integration site in gastric cancer tissues and para-cancerous tissues. A, B: Representative figures of wingless-related integration site (WNT5a) expression (cytoplasmic staining) at different grades of pathological scores in gastric cancer tissues (GC) (A) and para-cancerous (PC) tissues (B); C: Numbers of samples expressing WNT5a in GC and PC groups. Specimens were examined under a light microscope (200 × magnification). WNT5a: Wingless-related integration site.

The representative WNT ligand WNT5a can activate both canonical  $\beta$ -catenin-dependent and noncanonical  $\beta$ -catenin-independent WNT pathways[37] that participate in cancer cell migration and invasion by mediating the EMT[23,32,37-43]. Inhibition of the WNT/ $\beta$ -catenin signaling pathway also suppresses GC metastasis[21,44]. We examined  $\beta$ -catenin expression in clinical gastric tumor sections to

Baisbideng® WJCC https://www.wjgnet.com



Figure 3 Expression of β-catenin in gastric cancer tissues and para-cancerous tissues. A, B: Representative figures of β-catenin expression (membranous and cytoplasmic staining) at different grades of pathological scores in gastric cancer (GC) (A) and para-cancerous (PC) tissues (B); C: Numbers of samples expressing β-catenin in GC and PC groups. Specimens were examined under a light microscope (200 × magnification).

> determine which of the WNT/ $\beta$ -catenin pathways are involved in the development of GC. We found more abundant  $\beta$ -catenin in the membrane and cytoplasm of para-cancerous, than GC tissues, and that this did not correlate with sex and age of patients, or the degree of differentiation, invasion depth, distant metastasis, or TNM stage of tumors. This implies that WNT5a activated the canonical WNT signaling pathway in GC, resulting in the reduction of cytoplasmic  $\beta$ -catenin. However, to confirm that

Saisbideng® WJCC | https://www.wjgnet.com

WNT5a regulates the canonical  $\beta$ -catenin-dependent signaling pathway in GC metastasis requires further investigation of  $\beta$ -catenin nuclear staining.

# CONCLUSION

The expression of HNF4 $\alpha$  and WNT5a was more abundant in GC, than para-cancerous tissues, whereas that of cytoplasmic  $\beta$ -catenin was lower. The diagnostic accuracy rate of HNF4 $\alpha$  and WNT5a for GC was 91.0% and 85.7%, respectively, indicating that both could serve as potential diagnostic tools for GC. To our knowledge, this is the first study to find that HNF4 $\alpha$ , WNT5a, and  $\beta$ -catenin proteins are expressed in GC. However, this study is limited by the retrospective design. Because all specimens were pathological paraffin blocks of clinical GC, we could not investigate expression of the HNF4 $\alpha$ /WNT5a/ $\beta$ -catenin signaling axis using western blotting or PCR. Therefore, clinical trials are warranted to verify our conclusion.

# **ARTICLE HIGHLIGHTS**

#### Research background

The authors have previously found in gastric cancer (GC) cell lines and animal models that the expression of hepatocyte nuclear factor 4 alpha (HNF4 $\alpha$ ) is significantly increased, and it promotes the invasion and metastasis of GC by regulating its downstream wingless-related integration site (WNT)/ $\beta$ -catenin signaling pathway, while negatively regulating the HNF4 $\alpha$ /WNT signaling pathway can inhibit the development of GC.

#### **Research motivation**

Extending from the preclinical model to clinical GC tissues, we will continue to explore the expression of HNF4 $\alpha$  and its downstream WNT/ $\beta$ -catenin signaling pathway in GC tissues, in the hope of providing new treatment ideas for inhibiting the development of GC.

#### **Research objectives**

To explore the expression of HNF4 $\alpha$ , WNT5a and  $\beta$ -catenin in clinical GC tissues and adjacent tissues, and whether the expression of these three molecules in GC tissues is related.

## **Research methods**

Screening GC tissues containing each subtype and the adjacent para-cancerous tissues; recording the demographic characteristics of the patients and the pathological type and stage of the GC tissues; and detecting the expressions of HNF4 $\alpha$ , WNT5a and  $\beta$ -catenin in the GC tissues and the para-cancerous tissues by immunohistochemistry. The expression levels of these molecules were observed under the microscope and the expression levels were recorded according to the corresponding scoring criteria. Finally, the statistical methods such as analysis of chi-square and paired chi-square were used to analyze the expression differences and correlation of HNF4 $\alpha$ , WNT5a and  $\beta$ -catenin in GC tissues and para-cancerous tissues.

#### Research results

Compared with the para-cancerous tissues, the expressions of HNF4 $\alpha$  (nuclear staining) and WNT5a (cytoplasmic staining) in GC tissues were significantly increased, while the expression of  $\beta$ -catenin (cytoplasmic staining) was significantly decreased. HNF4 $\alpha$  expression was abundant in tubular and mucinous adenocarcinomas and relatively weak in signet ring cell carcinoma; WNT5a expression was more abundant in mucinous, than in tubular adenocarcinoma and signet ring cell carcinoma; both HNF4 $\alpha$  and WNT5a were strongly positive in mixed GC, although no positive correlations between HNF4 $\alpha$  and WNT5a expression in GC.

#### **Research conclusions**

Comparing the expression of HNF4 $\alpha$ , WNT5a and  $\beta$ -catenin in GC and para-cancerous tissues, we can find the expressions of HNF4 $\alpha$  and WNT5a could serve as early diagnostic biomarkers for GC.

#### Research perspectives

It is hopeful that HNF4α and WNT5a can be included into the molecules for the diagnosis of GC.

Raisbideng® WJCC | https://www.wjgnet.com

# ACKNOWLEDGEMENTS

We would like to thank Yao-Bing Chen for his help, a pathologist of Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology.

# FOOTNOTES

Author contributions: Yi P designed the study; Hu Q, Li LL and Peng Z conducted the experiments; Hu Q wrote and revised the manuscript; Yi P checked the pathological results; and All authors approved the final version of the article.

Supported by National Natural Science Foundation of China, No. 81673757.

Institutional review board statement: The study was reviewed and approved by the Institutional Review Board at Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology.

Conflict-of-interest statement: All authors have nothing to disclose.

Data sharing statement: No additional date is available.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is noncommercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

#### Country/Territory of origin: China

**ORCID number:** Qian Hu 0000-0002-5783-7085; Ling-Li Li 0000-0003-4518-1418; Ze Peng 0000-0003-2175-4322; Ping Yi 0000-0001-8560-2024.

S-Editor: Ma Y I-Editor: A P-Editor: Ma YJ

## REFERENCES

- De Re V. Molecular Features Distinguish Gastric Cancer Subtypes. Int J Mol Sci 2018; 19 [PMID: 30314372 DOI: 1 10.3390/ijms19103121]
- Song Z, Wu Y, Yang J, Yang D, Fang X. Progress in the treatment of advanced gastric cancer. Tumour Biol 2017; 39: 1010428317714626 [PMID: 28671042 DOI: 10.1177/1010428317714626]
- 3 Karimi P, Islami F, Anandasabapathy S, Freedman ND, Kamangar F. Gastric cancer: descriptive epidemiology, risk factors, screening, and prevention. Cancer Epidemiol Biomarkers Prev 2014; 23: 700-713 [PMID: 24618998 DOI: 10.1158/1055-9965.EPI-13-1057]
- 4 Pasechnikov V, Chukov S, Fedorov E, Kikuste I, Leja M. Gastric cancer: prevention, screening and early diagnosis. World J Gastroenterol 2014; 20: 13842-13862 [PMID: 25320521 DOI: 10.3748/wjg.v20.i38.13842]
- 5 Sitarz R, Skierucha M, Mielko J, Offerhaus GJA, Maciejewski R, Polkowski WP. Gastric cancer: epidemiology, prevention, classification, and treatment. Cancer Manag Res 2018; 10: 239-248 [PMID: 29445300 DOI: 10.2147/CMAR.S149619
- Yao HS, Wang J, Zhang XP, Wang LZ, Wang Y, Li XX, Jin KZ, Hu ZQ, Wang WJ. Hepatocyte nuclear factor 4a 6 suppresses the aggravation of colon carcinoma. Mol Carcinog 2016; 55: 458-472 [PMID: 25808746 DOI: 10.1002/mc.22294]
- Duncan SA, Manova K, Chen WS, Hoodless P, Weinstein DC, Bachvarova RF, Darnell JE Jr. Expression of transcription factor HNF-4 in the extraembryonic endoderm, gut, and nephrogenic tissue of the developing mouse embryo: HNF-4 is a marker for primary endoderm in the implanting blastocyst. Proc Natl Acad Sci USA 1994; 91: 7598-7602 [PMID: 8052626 DOI: 10.1073/pnas.91.16.7598]
- Vuong LM, Chellappa K, Dhahbi JM, Deans JR, Fang B, Bolotin E, Titova NV, Hoverter NP, Spindler SR, Waterman ML, Sladek FM. Differential Effects of Hepatocyte Nuclear Factor 4a Isoforms on Tumor Growth and T-Cell Factor 4/AP-1 Interactions in Human Colorectal Cancer Cells. Mol Cell Biol 2015; 35: 3471-3490 [PMID: 26240283 DOI: 10.1128/MCB.00030-15
- 9 Ma L, Zeng J, Guo Q, Liang X, Shen L, Li S, Sun Y, Li W, Liu S, Yu H, Chen C, Jia J. Mutual amplification of HNF4α and IL-1R1 composes an inflammatory circuit in Helicobacter pylori associated gastric carcinogenesis. Oncotarget 2016; 7: 11349-11363 [PMID: 26870992 DOI: 10.18632/oncotarget.7239]
- 10 Saberi Anvar M, Minuchehr Z, Shahlaei M, Kheitan S. Gastric cancer biomarkers; A systems biology approach. Biochem Biophys Rep 2018; 13: 141-146 [PMID: 29556568 DOI: 10.1016/j.bbrep.2018.01.001]



- 11 Taciak B, Pruszynska I, Kiraga L, Bialasek M, Krol M. Wnt signaling pathway in development and cancer. J Physiol Pharmacol 2018; 69 [PMID: 29980141 DOI: 10.26402/jpp.2018.2.07]
- 12 Nam S, Chang HR, Kim KT, Kook MC, Hong D, Kwon CH, Jung HR, Park HS, Powis G, Liang H, Park T, Kim YH. PATHOME: an algorithm for accurately detecting differentially expressed subpathways. Oncogene 2014; 33: 4941-4951 [PMID: 24681952 DOI: 10.1038/onc.2014.80]
- 13 Saitoh T, Mine T, Katoh M. Frequent up-regulation of WNT5A mRNA in primary gastric cancer. Int J Mol Med 2002; 9: 515-519 [PMID: 11956659]
- Nam S, Chung JW, Yang JY. WNT5A Correlates with Clinicopathological Characteristics in Gastric Cancer: a Meta-14 Analysis. Cell Physiol Biochem 2017; 41: 33-40 [PMID: 28135710 DOI: 10.1159/000455934]
- Hanaki H, Yamamoto H, Sakane H, Matsumoto S, Ohdan H, Sato A, Kikuchi A. An anti-Wnt5a antibody suppresses 15 metastasis of gastric cancer cells in vivo by inhibiting receptor-mediated endocytosis. Mol Cancer Ther 2012; 11: 298-307 [PMID: 22101459 DOI: 10.1158/1535-7163.MCT-11-0682]
- Xue Y, Zhang L, Zhu Y, Ke X, Wang Q, Min H. Regulation of Proliferation and Epithelial-to-Mesenchymal Transition (EMT) of Gastric Cancer by ZEB1 via Modulating Wnt5a and Related Mechanisms. Med Sci Monit 2019; 25: 1663-1670 [PMID: 30829316 DOI: 10.12659/MSM.912338]
- Gao M, Liu L, Yang Y, Li M, Ma Q, Chang Z. LncRNA HCP5 Induces Gastric Cancer Cell Proliferation, Invasion, and 17 EMT Processes Through the miR-186-5p/WNT5A Axis Under Hypoxia. Front Cell Dev Biol 2021; 9: 663654 [PMID: 34178988 DOI: 10.3389/fcell.2021.663654]
- 18 Chang HR, Nam S, Kook MC, Kim KT, Liu X, Yao H, Jung HR, Lemos R Jr, Seo HH, Park HS, Gim Y, Hong D, Huh I, Kim YW, Tan D, Liu CG, Powis G, Park T, Liang H, Kim YH. HNF4α is a therapeutic target that links AMPK to WNT signalling in early-stage gastric cancer. Gut 2016; 65: 19-32 [PMID: 25410163 DOI: 10.1136/gutjnl-2014-307918]
- 19 Yang M, Li SN, Anjum KM, Gui LX, Zhu SS, Liu J, Chen JK, Liu QF, Ye GD, Wang WJ, Wu JF, Cai WY, Sun GB, Liu YJ, Liu RF, Zhang ZM, Li BA. A double-negative feedback loop between Wnt-β-catenin signaling and HNF4α regulates epithelial-mesenchymal transition in hepatocellular carcinoma. J Cell Sci 2013; 126: 5692-5703 [PMID: 24101726 DOI: 10.1242/jcs.135053]
- 20 Hu Q, Li L, Zou X, Xu L, Yi P. Berberine Attenuated Proliferation, Invasion and Migration by Targeting the AMPK/HNF4 a/WNT5A Pathway in Gastric Carcinoma. Front Pharmacol 2018; 9: 1150 [PMID: 30405404 DOI: 10.3389/fphar.2018.01150
- 21 Wang H, Dong S, Liu Y, Ma F, Fang J, Zhang W, Shao S, Shen H, Jin J. DAB2 suppresses gastric cancer migration by regulating the Wnt/β-catenin and Hippo-YAP signaling pathways. Transl Cancer Res 2020; 9: 1174-1184 [PMID: 35117462 DOI: 10.21037/tcr.2019.12.96]
- 22 Fusco N, Rocco EG, Del Conte C, Pellegrini C, Bulfamante G, Di Nuovo F, Romagnoli S, Bosari S. HER2 in gastric cancer: a digital image analysis in pre-neoplastic, primary and metastatic lesions. Mod Pathol 2013; 26: 816-824 [PMID: 23348899 DOI: 10.1038/modpathol.2012.228]
- 23 Kanzawa M, Semba S, Hara S, Itoh T, Yokozaki H. WNT5A is a key regulator of the epithelial-mesenchymal transition and cancer stem cell properties in human gastric carcinoma cells. Pathobiology 2013; 80: 235-244 [PMID: 23615002 DOI: 10.1159/000346843]
- Kulkarni RN, Kahn CR. Molecular biology. HNFs--linking the liver and pancreatic islets in diabetes. Science 2004; 303: 24 1311-1312 [PMID: 14988544 DOI: 10.1126/science.1095486]
- 25 Marcil V, Amre D, Seidman EG, Boudreau F, Gendron FP, Ménard D, Beaulieu JF, Sinnett D, Lambert M, Levy E. Hepatocyte nuclear factor 4 alpha polymorphisms and the metabolic syndrome in French-Canadian youth. PLoS One 2015; 10: e0117238 [PMID: 25671620 DOI: 10.1371/journal.pone.0117238]
- Naiki T, Nagaki M, Shidoji Y, Kojima H, Imose M, Kato T, Ohishi N, Yagi K, Moriwaki H. Analysis of gene expression 26 profile induced by hepatocyte nuclear factor 4alpha in hepatoma cells using an oligonucleotide microarray. J Biol Chem 2002; 277: 14011-14019 [PMID: 11834723 DOI: 10.1074/jbc.M105403200]
- 27 Chen Z, Gropler MC, Mitra MS, Finck BN. Complex interplay between the lipin 1 and the hepatocyte nuclear factor 4 α (HNF4α) pathways to regulate liver lipid metabolism. PLoS One 2012; 7: e51320 [PMID: 23236470 DOI: 10.1371/journal.pone.0051320]
- 28 Sugano M, Nagasaka T, Sasaki E, Murakami Y, Hosoda W, Hida T, Mitsudomi T, Yatabe Y. HNF4a as a marker for invasive mucinous adenocarcinoma of the lung. Am J Surg Pathol 2013; 37: 211-218 [PMID: 23108025 DOI: 10.1097/PAS.0b013e31826be303
- 29 Xu C, Ooi WF, Qamra A, Tan J, Chua BY, Ho SWT, Das K, Adam Isa ZF, Li Z, Yao X, Yan T, Xing M, Huang KK, Lin JS, Nandi T, Tay ST, Lee MH, Tan ALK, Ong X, Ashktorab H, Smoot D, Li S, Ng SC, Teh BT, Tan P. HNF4a pathway mapping identifies wild-type IDH1 as a targetable metabolic node in gastric cancer. Gut 2020; 69: 231-242 [PMID: 31068366 DOI: 10.1136/gutjnl-2018-318025]
- 30 Ma Y, Wei X, Wu Z. HNF-4α promotes multidrug resistance of gastric cancer cells through the modulation of cell apoptosis. Oncol Lett 2017; 14: 6477-6484 [PMID: 29344114 DOI: 10.3892/ol.2017.7095]
- Kurayoshi M, Oue N, Yamamoto H, Kishida M, Inoue A, Asahara T, Yasui W, Kikuchi A. Expression of Wnt-5a is 31 correlated with aggressiveness of gastric cancer by stimulating cell migration and invasion. Cancer Res 2006; 66: 10439-10448 [PMID: 17079465 DOI: 10.1158/0008-5472.CAN-06-2359]
- 32 Shojima K, Sato A, Hanaki H, Tsujimoto I, Nakamura M, Hattori K, Sato Y, Dohi K, Hirata M, Yamamoto H, Kikuchi A. Wht5a promotes cancer cell invasion and proliferation by receptor-mediated endocytosis-dependent and -independent mechanisms, respectively. Sci Rep 2015; 5: 8042 [PMID: 25622531 DOI: 10.1038/srep08042]
- Yamamoto H, Kitadai Y, Yamamoto H, Oue N, Ohdan H, Yasui W, Kikuchi A. Laminin gamma2 mediates Wnt5a-33 induced invasion of gastric cancer cells. Gastroenterology 2009; 137: 242-252, 252.e1 [PMID: 19582886 DOI: 10.1053/j.gastro.2009.02.003]
- 34 Liu Q, Yang C, Wang S, Shi D, Wei C, Song J, Lin X, Dou R, Bai J, Xiang Z, Huang S, Liu K, Xiong B. Wnt5a-induced M2 polarization of tumor-associated macrophages via IL-10 promotes colorectal cancer progression. Cell Commun Signal 2020; 18: 51 [PMID: 32228612 DOI: 10.1186/s12964-020-00557-2]



- 35 Liu X, Xiu LJ, Jiao JP, Zhao J, Zhao Y, Lu Y, Shi J, Li YJ, Ye M, Gu YF, Wang XW, Xu JY, Zhang CA, Liu YY, Luo Y, Yue XQ. Traditional Chinese medicine integrated with chemotherapy for stage IV non-surgical gastric cancer: a retrospective clinical analysis. J Integr Med 2017; 15: 469-475 [PMID: 29103417 DOI: 10.1016/S2095-4964(17)60377-7]
- 36 Koni M, Pinnarò V, Brizzi MF. The Wnt Signalling Pathway: A Tailored Target in Cancer. Int J Mol Sci 2020; 21 [PMID: 33080952 DOI: 10.3390/ijms21207697]
- 37 Gencer S, Sen G, Doğusoy G, Bellı AK, Paksoy M, Yazicioğlu MB. β-Catenin-independent noncanonical Wnt pathway might be induced in gastric cancers. Turk J Gastroenterol 2010; 21: 224-230 [PMID: 20931424 DOI: 10.4318/tjg.2010.0092]
- 38 Asem MS, Buechler S, Wates RB, Miller DL, Stack MS. Wnt5a Signaling in Cancer. Cancers (Basel) 2016; 8 [PMID: 27571105 DOI: 10.3390/cancers8090079]
- 39 Tian S, Hu J, Tao K, Wang J, Chu Y, Li J, Liu Z, Ding X, Xu L, Li Q, Cai M, Gao J, Shuai X, Wang G, Wang L, Wang Z. Secreted AGR2 promotes invasion of colorectal cancer cells via Wnt11-mediated non-canonical Wnt signaling. Exp Cell Res 2018; 364: 198-207 [PMID: 29427622 DOI: 10.1016/j.yexcr.2018.02.004]
- 40 Tseng JC, Lin CY, Su LC, Fu HH, Yang SD, Chuu CP. CAPE suppresses migration and invasion of prostate cancer cells via activation of non-canonical Wnt signaling. Oncotarget 2016; 7: 38010-38024 [PMID: 27191743 DOI: 10.18632/oncotarget.9380]
- 41 Tian S, Peng P, Li J, Deng H, Zhan N, Zeng Z, Dong W. SERPINH1 regulates EMT and gastric cancer metastasis via the Wnt/β-catenin signaling pathway. Aging (Albany NY) 2020; 12: 3574-3593 [PMID: 32091407 DOI: 10.18632/aging.102831]
- 42 Wu Q, Ma J, Wei J, Meng W, Wang Y, Shi M. IncRNA SNHG11 Promotes Gastric Cancer Progression by Activating the Wnt/β-Catenin Pathway and Oncogenic Autophagy. Mol Ther 2021; 29: 1258-1278 [PMID: 33068778 DOI: 10.1016/j.vmthe.2020.10.011
- 43 Guo Q, Xu J, Huang Z, Yao Q, Chen F, Liu H, Zhang Z, Lin J. ADMA mediates gastric cancer cell migration and invasion via Wnt/β-catenin signaling pathway. Clin Transl Oncol 2021; 23: 325-334 [PMID: 32607811 DOI: 10.1007/s12094-020-02422-7]
- 44 Ge Q, Hu Y, He J, Chen F, Wu L, Tu X, Qi Y, Zhang Z, Xue M, Chen S, Zhong J, Wang L. Zicl suppresses gastric cancer metastasis by regulating Wnt/β-catenin signaling and epithelial-mesenchymal transition. FASEB J 2020; 34: 2161-2172 [PMID: 31909528 DOI: 10.1096/fj.201901372RR]





# Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: bpgoffice@wjgnet.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

