

World Journal of *Clinical Cases*

World J Clin Cases 2022 August 16; 10(23): 8057-8431



Contents

Thrice Monthly Volume 10 Number 23 August 16, 2022

OPINION REVIEW

- 8057** Invasive intervention timing for infected necrotizing pancreatitis: Late invasive intervention is not late for collection

Xiao NJ, Cui TT, Liu F, Li W

- 8063** Clinical utility of left atrial strain in predicting atrial fibrillation recurrence after catheter ablation: An up-to-date review

Yu ZX, Yang W, Yin WS, Peng KX, Pan YL, Chen WW, Du BB, He YQ, Yang P

MINIREVIEWS

- 8076** Gut microbiota and COVID-19: An intriguing pediatric perspective

Valentino MS, Esposito C, Colosimo S, Caprio AM, Puzone S, Guarino S, Marzuillo P, Miraglia del Giudice E, Di Sessa A

- 8088** Beta receptor blocker therapy for the elderly in the COVID-19 era

Santillo E, Migale M

ORIGINAL ARTICLE

Retrospective Cohort Study

- 8097** Nonselective beta-blocker use is associated with increased hepatic encephalopathy-related readmissions in cirrhosis

Fallahzadeh MA, Asrani SK, Tapper EB, Saracino G, Rahimi RS

Retrospective Study

- 8107** Different squatting positions after total knee arthroplasty: A retrospective study

Li TJ, Sun JY, Du YQ, Shen JM, Zhang BH, Zhou YG

- 8115** Outcomes of seromuscular bladder augmentation compared with standard bladder augmentation in the treatment of children with neurogenic bladder

Sun XG, Li YX, Ji LF, Xu JL, Chen WX, Wang RY

- 8124** Distinctive clinical features of spontaneous pneumoperitoneum in neonates: A retrospective analysis

Kim SH, Cho YH, Kim HY

- 8133** Cognitive training for elderly patients with early Alzheimer's disease in the Qinghai-Tibet Plateau: A pilot study

Wang XH, Luo MQ

- 8141** Diagnostic value of elevated serum carbohydrate antigen 125 level in sarcoidosis

Zhang Q, Jing XY, Yang XY, Xu ZJ

- 8152** Evaluation of progressive early rehabilitation training mode in intensive care unit patients with mechanical ventilation

Qie XJ, Liu ZH, Guo LM

- 8161** Comparison of demographic features and laboratory parameters between COVID-19 deceased patients and surviving severe and critically ill cases

Wang L, Gao Y, Zhang ZJ, Pan CK, Wang Y, Zhu YC, Qi YP, Xie FJ, Du X, Li NN, Chen PF, Yue CS, Wu JH, Wang XT, Tang YJ, Lai QQ, Kang K

Clinical Trials Study

- 8170** Role of H₂receptor blocker famotidine over the clinical recovery of COVID-19 patients: A randomized controlled trial

Mohiuddin Chowdhury ATM, Kamal A, Abbas MKU, Karim MR, Ali MA, Talukder S, Hamidullah Mehedi H, Hassan H, Shahin AH, Li Y, He S

Observational Study

- 8186** Short-term prognostic factors for hepatitis B virus-related acute-on-chronic liver failure

Ye QX, Huang JF, Xu ZJ, Yan YY, Yan Y, Liu LG

- 8196** Three-dimensional psychological guidance combined with evidence-based health intervention in patients with liver abscess treated with ultrasound

Shan YN, Yu Y, Zhao YH, Tang LL, Chen XM

- 8205** Role of serum β 2-microglobulin, glycosylated hemoglobin, and vascular endothelial growth factor levels in diabetic nephropathy

Yang B, Zhao XH, Ma GB

SYSTEMATIC REVIEWS

- 8212** Gallbladder neuroendocrine carcinoma diagnosis, treatment and prognosis based on the SEER database: A literature review

Cai XC, Wu SD

CASE REPORT

- 8224** Sepsis complicated with secondary hemophagocytic syndrome induced by giant gouty tophi rupture: A case report

Lai B, Pang ZH

- 8232** Spontaneous remission of autoimmune pancreatitis: Four case reports

Zhang BB, Huo JW, Yang ZH, Wang ZC, Jin EH

- 8242** Epstein-Barr-virus-associated hepatitis with aplastic anemia: A case report

Zhang WJ, Wu LQ, Wang J, Lin SY, Wang B

- 8249** Aspiration as the first-choice procedure for airway management in an infant with large epiglottic cysts: A case report

Zheng JQ, Du L, Zhang WY

- 8255** Sequential multidisciplinary minimally invasive therapeutic strategy for heart failure caused by four diseases: A case report
Zhao CZ, Yan Y, Cui Y, Zhu N, Ding XY
- 8262** Primary ascending colon cancer accompanying skip metastases in left shoulder skin and left neck lymph node: A case report
Zhou JC, Wang JJ, Liu T, Tong Q, Fang YJ, Wu ZQ, Hong Q
- 8271** Clinical and genetic study of ataxia with vitamin E deficiency: A case report
Zhang LW, Liu B, Peng DT
- 8277** Complete resection of large-cell neuroendocrine and hepatocellular carcinoma of the liver: A case report
Noh BG, Seo HI, Park YM, Kim S, Hong SB, Lee SJ
- 8284** Immunotherapy combined with antiangiogenic agents in patients with advanced malignant pleural mesothelioma: A case report
Xuan TT, Li GY, Meng SB, Wang ZM, Qu LL
- 8291** Bladder malacoplakia: A case report
Wang HK, Hang G, Wang YY, Wen Q, Chen B
- 8298** Delayed inflammatory response evoked in nasal alloplastic implants after COVID-19 vaccination: A case report
Seo MG, Choi EK, Chung KJ
- 8304** Phosphoglyceride crystal deposition disease requiring differential diagnosis from malignant tumors and confirmed by Raman spectroscopy: A case report
Ohkura Y, Uruga H, Shiiba M, Ito S, Shimoyama H, Ishihara M, Ueno M, Udagawa H
- 8312** Vulvovaginal myeloid sarcoma with massive pelvic floor infiltration: A case report and review of literature
Wang JX, Zhang H, Ning G, Bao L
- 8323** Femoral neck stress fracture and medial tibial stress syndrome following high intensity interval training: A case report and review of literature
Tan DS, Cheung FM, Ng D, Cheung TLA
- 8330** Periosteal chondroma of the rib: A case report
Gao Y, Wang JG, Liu H, Gao CP
- 8336** Papillary thyroid carcinoma occurring with undifferentiated pleomorphic sarcoma: A case report
Lee YL, Cheng YQ, Zhu CF, Huo HZ
- 8344** Laparoscopic treatment of bilateral duplex kidney and ectopic ureter: A case report
Wang SB, Wan L, Wang Y, Yi ZJ, Xiao C, Cao JZ, Liu XY, Tang RP, Luo Y
- 8352** Incontinentia pigmenti with intracranial arachnoid cyst: A case report
Li WC, Li ML, Ding JW, Wang L, Wang SR, Wang YY, Xiao LF, Sun T

- 8360** Relapsing polychondritis causing breathlessness: Two case reports
Zhai SY, Zhang YH, Guo RY, Hao JW, Wen SX
- 8367** Endodontic management of a fused left maxillary second molar and two paramolars using cone beam computed tomography: A case report
Mei XH, Liu J, Wang W, Zhang QX, Hong T, Bai SZ, Cheng XG, Tian Y, Jiang WK
- 8375** Infant biliary cirrhosis secondary to a biliary inflammatory myofibroblastic tumor: A case report and review of literature
Huang Y, Shu SN, Zhou H, Liu LL, Fang F
- 8384** Metastatic low-grade endometrial stromal sarcoma with variable morphologies in the ovaries and mesentery: A case report
Yu HY, Jin YL
- 8392** Bronchogenic cysts with infection in the chest wall skin of a 64-year-old asymptomatic patient: A case report
Ma B, Fu KW, Xie XD, Cheng Y, Wang SQ
- 8400** Incidental accumulation of Technetium-99m pertechnetate in subacute cerebral infarction: A case report
Han YH, Jeong HJ, Kang HG, Lim ST
- 8406** Metal stent combined with ileus drainage tube for the treatment of delayed rectal perforation: A case report
Cheng SL, Xie L, Wu HW, Zhang XF, Lou LL, Shen HZ
- 8417** Using ketamine in a patient with a near-occlusion tracheal tumor undergoing tracheal resection and reconstruction: A case report
Xu XH, Gao H, Chen XM, Ma HB, Huang YG

LETTER TO THE EDITOR

- 8422** Reflections on the prevalence of human leukocyte antigen-B27 and human leukocyte antigen-B51 co-occurrence in patients with spondylarthritis
Gonçalves Júnior J, Sampaio-Barros PD, Shinjo SK
- 8425** Comment on "Disease exacerbation is common in inflammatory bowel disease patients treated with immune checkpoint inhibitors for malignancy"
Argyriou K, Kotsakis A
- 8428** Intranasal sufentanil combined with intranasal dexmedetomidine: A promising method for non-anesthesiologist sedation during endoscopic ultrasonography
Wang Y, Ge ZJ, Han C

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Peng Liang, MD, Associate Professor, Day Surgery Center, Department of Anesthesiology, West China Hospital of Sichuan University, Chengdu 610041, Sichuan Province, China. 39485572@qq.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Hua-Ge Yin; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

August 16, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Intranasal sufentanil combined with intranasal dexmedetomidine: A promising method for non-anesthesiologist sedation during endoscopic ultrasonography

Yong Wang, Zhi-Jun Ge, Chao Han

Specialty type: Medicine, research and experimental

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Amornytin S, Thailand; Raghuraman MS, India

Received: March 30, 2022

Peer-review started: March 30, 2022

First decision: June 19, 2022

Revised: June 24, 2022

Accepted: July 16, 2022

Article in press: July 16, 2022

Published online: August 16, 2022



Yong Wang, Chao Han, Department of Anesthesiology, The Affiliated Yixing Hospital of Jiangsu University, Yixing 214200, Jiangsu Province, China

Yong Wang, Zhi-Jun Ge, School of Medical, Jiangsu University, Zhenjiang 212013, Jiangsu Province, China

Corresponding author: Chao Han, MD, Assistant Professor, Department of Anesthesiology, The Affiliated Yixing Hospital of Jiangsu University, No. 75 Tongzhenguan Road, Yixing 214200, Jiangsu Province, China. staff940@yxph.com

Abstract

Intranasal sufentanil combined with intranasal dexmedetomidine exhibited an estimated sedation success probability as high as 94.9%, higher satisfaction scores, and only minor adverse events during endoscopic ultrasonography (EUS). This is a promising method for EUS sedation that does not require the presence of an anesthesiologist.

Key Words: Intranasal; Sufentanil; Dexmedetomidine; Sedation; Endoscopic Ultrasonography

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Endoscopic ultrasonography (EUS) requires moderate-to-deep sedation due to a prolonged procedure time and a larger and stiffer probe. Propofol-based sedation is the predominant method used in such cases for rapid onset and improved sedation with rapid full recovery. However, there are still restrictions regarding the administration of propofol in the absence of an anesthesiologist. The combination of intranasal sufentanil and intranasal dexmedetomidine exhibited an estimated sedation success probability, higher satisfaction scores, and minor adverse events, thus highlighting a promising method for EUS sedation in the absence of an anesthesiologist.

Citation: Wang Y, Ge ZJ, Han C. Intranasal sufentanil combined with intranasal dexmedetomidine: A promising method for non-anesthesiologist sedation during endoscopic ultrasonography. *World J Clin Cases* 2022; 10(23): 8428-8431

URL: <https://www.wjgnet.com/2307-8960/full/v10/i23/8428.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i23.8428>

TO THE EDITOR

We read with interest the paper published by Zou *et al*[1] exploring the ED95 of intranasal sufentanil (SUF) combined with intranasal dexmedetomidine (DEX) for moderate sedation during endoscopic ultrasonography (EUS). EUS has been widely used clinically due to its unique diagnostic value regarding lesions arising from the pancreas, upper gastrointestinal tract, as well as adjacent structures, such as the liver and lymph nodes[2-4]. However, EUS procedures take a long time and use a larger and stiffer probe compared to conventional endoscopes. Thus, it is important that we ensure that EUS tolerability is acceptable.

In patients undergoing complex endoscopic procedures, the current standard of practice is to administer sedative medication intravenously. Propofol-based sedation has become the first choice for endoscopic sedation over the last two decades because of its fast onset of action and short half-life[5,6]. Although there is compelling evidence to support the quality, cost effectiveness, and safety profile of the administration of propofol for EUS in the absence of an anesthesiologist[7,8], it is currently recommended that the administration and monitoring of propofol sedation for endoscopic procedures should be the responsibility of a dedicated and appropriately trained anesthetist only. A joint position statement endorsed by the British Society of Gastroenterology, the Joint Advisory Group, and the Royal College of Anesthetists highlights the role of anesthetist-led deep sedation practice with a focus on propofol sedation[9]. This is because propofol can produce serious and potentially fatal side effects such as hypotension, bradycardia, hypoventilation, hypoxemia, and even apnea. Therefore, propofol-independent sedation regimens have received increasing attention since such regimens could avoid the controversy related to non-anesthesiologist-administered propofol sedation[10,11].

Here, the paper proposed a promising and alternative regimen for non-anesthesiologist propofol administration during EUS, despite the authors' original intention not to do so. Intranasal administration offers a noninvasive, rapid, and efficient route for drug delivery with stable hemodynamics compared to the intravenous route due to a slower and more gradual onset. The authors made ingenious use of the pharmacokinetic characteristics of SUF and DEX. Intranasal 1 µg/kg DEX was administered 45 min before EUS and intranasal 0.3 µg/kg SUF was administered 20 min later, ensuring that the two drugs achieved peak effect during the procedure[12,13]. Sequential intranasal therapy exhibited an estimated sedation success probability that reached 94.9%, higher satisfaction scores, and minor adverse events. The emerging sedation regimen makes it possible to administer EUS sedation without an anesthesiologist and has high clinical significance. However, there are still several issues that need to be discussed with the authors before this technology is popularized. The modified observer's assessment of alert (MOAA/S) scale is generally classified as deep sedation (0 to 1), moderate sedation (2 to 3) and slight sedation (4 to 5)[14,15]. However, the authors defined a successful moderate sedation as a score ≤ 3 on the MOAA/S a scale; this would mean that deep sedation (MOAA/S: 0 to 1) into the category of moderate sedation. Although deep sedation does not mean oversedation, this might be more understandable if the authors could clarify the difference between moderate sedation and oversedation. In addition, the paper does not describe the MOAA/S scores of patients throughout their procedures. The authors acknowledged the presence of fluctuating levels of sedation although the use of rescue sedation was not reported. Insufficient or excessive sedation will inevitably lead to complications and may cause readers to worry about the safety of this technique.

Another issue worth noting is that the authors did not describe the time elapsed from the onset of sedation to discharge, and according to the pharmacokinetics of SUF and DEX, the maintenance time for both drugs is approximately 2 h[12,13]. Conversely, the average procedure time in this study was only 30 min, indicating that patients remained under sedation for 1 h after the completion of EUS. This might delay recovery when compared with the established protocol involving propofol sedation. We hope that the authors will provide relevant data to answer these concerns.

In our opinion, the most significant problem with this novel technology is the feasibility of clinical promotion. The sedation preparation prior to EUS took 45 min in this study. Transnasal medication also requires medical monitoring, thus increasing the occupation of medical resources and causing delays in the procedure, especially in outpatients. A separate induction room and complete service flow are prerequisites for the application of this new sedation regimen. Furthermore, as the article mentioned, the study was performed in patients with a normal body mass index (BMI). This would exclude the occurrence of serious respiratory depression in those with a high BMI.

To conclude, we believe that the authors provided a promising regimen for non-anesthesiologist sedation during EUS even if sedation levels fluctuated throughout the study period. Nevertheless, further studies are now needed to confirm the safety profile of this technique in different populations.

FOOTNOTES

Author contributions: Wang Y and Ge ZJ designed and performed the research; Han C wrote and revised the letter.

Conflict-of-interest statement: All authors have no conflicts of interest to declare.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Chao Han 0000-0002-6869-0701.

Corresponding Author's Membership in Professional Societies: Jiangsu Medical Association, M010002136M.

S-Editor: Wang LL

L-Editor: Filipodia

P-Editor: Wang LL

REFERENCES

- 1 Zou Y, Li N, Shao LJ, Liu FK, Xue FS, Tao X. Determination of the ED₉₅ of intranasal sufentanil combined with intranasal dexmedetomidine for moderate sedation during endoscopic ultrasonography. *World J Clin Cases* 2022; **10**: 2773-2782 [PMID: 35434098 DOI: 10.12998/wjcc.v10.i9.2773]
- 2 Su Q, Peng J, Chen X, Xiao Z, Liu R, Wang F. Role of endoscopic ultrasonography for differential diagnosis of upper gastrointestinal submucosal lesions. *BMC Gastroenterol* 2021; **21**: 365 [PMID: 34620107 DOI: 10.1186/s12876-021-01945-9]
- 3 Tanaka H, Matsusaki S. The Utility of Endoscopic-Ultrasonography-Guided Tissue Acquisition for Solid Pancreatic Lesions. *Diagnostics (Basel)* 2022; **12** [PMID: 35328306 DOI: 10.3390/diagnostics12030753]
- 4 De Lisi S, Giovannini M. Endoscopic ultrasonography: Transition towards the future of gastro-intestinal diseases. *World J Gastroenterol* 2016; **22**: 1779-1786 [PMID: 26855537 DOI: 10.3748/wjg.v22.i5.1779]
- 5 Qin Y, Chen S, Zhang Y, Liu W, Lin Y, Chi X, Chen X, Yu Z, Su D. A Bibliometric Analysis of Endoscopic Sedation Research: 2001-2020. *Front Med (Lausanne)* 2021; **8**: 775495 [PMID: 35047526 DOI: 10.3389/fmed.2021.775495]
- 6 Teh JL, Shabbir A, Yuen S, So JB. Recent advances in diagnostic upper endoscopy. *World J Gastroenterol* 2020; **26**: 433-447 [PMID: 32063692 DOI: 10.3748/wjg.v26.i4.433]
- 7 Fatima H, DeWitt J, LeBlanc J, Sherman S, McGreevy K, Imperiale TF. Nurse-administered propofol sedation for upper endoscopic ultrasonography. *Am J Gastroenterol* 2008; **103**: 1649-1656 [PMID: 18557709 DOI: 10.1111/j.1572-0241.2008.01906.x]
- 8 Redondo-Cerezo E, Sánchez-Robaina A, Martínez Cara JG, Ojeda-Hinojosa M, Matas-Cobos A, Sánchez Capilla AD, López de Hierro Ruiz M, Pleguezuelo-Díaz J, de Teresa J. Gastroenterologist-guided sedation with propofol for endoscopic ultrasonography in average-risk and high-risk patients: a prospective series. *Eur J Gastroenterol Hepatol* 2012; **24**: 506-512 [PMID: 22330236 DOI: 10.1097/MEG.0b013e328350fcbf]
- 9 Sidhu R, Turnbull D, Newton M, Thomas-Gibson S, Sanders DS, Hebbard S, Haidry RJ, Smith G, Webster G. Deep sedation and anaesthesia in complex gastrointestinal endoscopy: a joint position statement endorsed by the British Society of Gastroenterology (BSG), Joint Advisory Group (JAG) and Royal College of Anaesthetists (RCoA). *Frontline Gastroenterol* 2019; **10**: 141-147 [PMID: 31205654 DOI: 10.1136/flgastro-2018-101145]
- 10 Guimaraes ES, Campbell EJ, Richter JM. The safety of nurse-administered procedural sedation compared to anesthesia care in a historical cohort of advanced endoscopy patients. *Anesth Analg* 2014; **119**: 349-356 [PMID: 24859079 DOI: 10.1213/ANE.0000000000000258]
- 11 Perel A. Non-anesthesiologists should not be allowed to administer propofol for procedural sedation: a Consensus Statement of 21 European National Societies of Anaesthesia. *Eur J Anaesthesiol* 2011; **28**: 580-584 [PMID: 21705907 DOI: 10.1097/EJA.0b013e328348a977]
- 12 Weerink MAS, Struys MMRF, Hannivoort LN, Barends CRM, Absalom AR, Colin P. Clinical Pharmacokinetics and Pharmacodynamics of Dexmedetomidine. *Clin Pharmacokinet* 2017; **56**: 893-913 [PMID: 28105598 DOI: 10.1007/s40262-017-0507-7]
- 13 Helmers JH, Noorduyn H, Van Peer A, Van Leeuwen L, Zuurmond WW. Comparison of intravenous and intranasal sufentanil absorption and sedation. *Can J Anaesth* 1989; **36**: 494-497 [PMID: 2529048 DOI: 10.1007/BF03005373]
- 14 Gan TJ, Berry BD, Ekman EF, Muckerman RC 2nd, Shore N, Hardi R. Safety evaluation of fospropofol for sedation

- during minor surgical procedures. *J Clin Anesth* 2010; **22**: 260-267 [PMID: [20522356](#) DOI: [10.1016/j.jclinane.2009.08.007](#)]
- 15 **Pastis NJ**, Hill NT, Yarmus LB, Schippers F, Imre M, Sohngen W, Randall O, Callahan SP, Silvestri GA. Correlation of Vital Signs and Depth of Sedation by Modified Observer's Assessment of Alertness and Sedation (MOAA/S) Scale in Bronchoscopy. *J Bronchology Interv Pulmonol* 2022; **29**: 54-61 [PMID: [34238838](#) DOI: [10.1097/LBR.0000000000000784](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

