

# World Journal of *Clinical Cases*

*World J Clin Cases* 2022 September 6; 10(25): 8808-9179



**MINIREVIEWS**

- 8808** Ear, nose, and throat manifestations of COVID-19 and its vaccines  
*Al-Ani RM*
- 8816** Potential influences of religiosity and religious coping strategies on people with diabetes  
*Onyishi CN, Eseadi C, Itechukwu LC, Okoro KN, Okolie CN, Egbule E, Asogwa E*

**ORIGINAL ARTICLE****Case Control Study**

- 8827** Effectiveness of six-step complex decongestive therapy for treating upper limb lymphedema after breast cancer surgery  
*Zhang HZ, Zhong QL, Zhang HT, Luo QH, Tang HL, Zhang LJ*

**Retrospective Study**

- 8837** Hospital admissions from alcohol-related acute pancreatitis during the COVID-19 pandemic: A single-centre study  
*Mak WK, Di Mauro D, Pearce E, Karran L, Myintmo A, Duckworth J, Orabi A, Lane R, Holloway S, Manzelli A, Mossadegh S*

- 8844** Indocyanine green plasma clearance rate and <sup>99m</sup>Tc-galactosyl human serum albumin single-photon emission computed tomography evaluated preoperative remnant liver  
*Iwaki K, Kaihara S, Kita R, Kitamura K, Hashida H, Uryuhara K*

- 8854** Arthroscopy with subscapularis upper one-third tenodesis for treatment of recurrent anterior shoulder instability independent of glenoid bone loss  
*An BJ, Wang FL, Wang YT, Zhao Z, Wang MX, Xing GY*

- 8863** Evaluation of the prognostic nutritional index for the prognosis of Chinese patients with high/extremely high-risk prostate cancer after radical prostatectomy  
*Yang F, Pan M, Nie J, Xiao F, Zhang Y*

**Observational Study**

- 8872** Chlorine poisoning caused by improper mixing of household disinfectants during the COVID-19 pandemic: Case series  
*Lin GD, Wu JY, Peng XB, Lu XX, Liu ZY, Pan ZG, Qiu ZW, Dong JG*

- 8880** Mental health of the Slovak population during COVID-19 pandemic: A cross-sectional survey  
*Kralova M, Brazinova A, Sivcova V, Izakova L*

**Prospective Study**

- 8893** Arthroscopic anatomical reconstruction of lateral collateral ligaments with ligament advanced reinforcement system artificial ligament for chronic ankle instability  
*Wang Y, Zhu JX*

**SYSTEMATIC REVIEWS**

- 8906** How to select the quantitative magnetic resonance technique for subjects with fatty liver: A systematic review  
*Li YW, Jiao Y, Chen N, Gao Q, Chen YK, Zhang YF, Wen QP, Zhang ZM*
- 8922** Lymphocytic choriomeningitis virus: An under-recognized congenital teratogen  
*Ferenc T, Vujica M, Mrzljak A, Vilibic-Cavlek T*

**CASE REPORT**

- 8932** Alagille syndrome associated with total anomalous pulmonary venous connection and severe xanthomas: A case report  
*Zeng HS, Zhang ZH, Hu Y, Zheng GL, Wang J, Zhang JW, Guo YX*
- 8939** Colo-colonic intussusception with post-polypectomy electrocoagulation syndrome: A case report  
*Moon JY, Lee MR, Yim SK, Ha GW*
- 8945** Portal vein gas combined with pneumatosis intestinalis and emphysematous cystitis: A case report and literature review  
*Hu SF, Liu HB, Hao YY*
- 8954** Quadricuspid aortic valve and right ventricular type of myocardial bridging in an asymptomatic middle-aged woman: A case report  
*Sopek Merkaš I, Lakušić N, Paar MH*
- 8962** Treatment of gastric carcinoma with lymphoid stroma by immunotherapy: A case report  
*Cui YJ, Ren YY, Zhang HZ*
- 8968** Gallstone associated celiac trunk thromboembolisms complicated with splenic infarction: A case report  
*Wu CY, Su CC, Huang HH, Wang YT, Wang CC*
- 8974** Extracorporeal membrane oxygenation for lung cancer-related life-threatening hypoxia: A case report  
*Yoo SS, Lee SY, Choi SH*
- 8980** Multi-disciplinary treatment of maxillofacial skeletal deformities by orthognathic surgery combined with periodontal phenotype modification: A case report  
*Liu JY, Li GF, Tang Y, Yan FH, Tan BC*
- 8990** X-linked recessive Kallmann syndrome: A case report  
*Zhang P, Fu JY*
- 8998** Delayed complications of intradural cement leakage after percutaneous vertebroplasty: A case report  
*Ma QH, Liu GP, Sun Q, Li JG*

- 9004** Coexistent Kaposi sarcoma and post-transplant lymphoproliferative disorder in the same lymph nodes after pediatric liver transplantation: A case report  
*Zhang SH, Chen GY, Zhu ZJ, Wei L, Liu Y, Liu JY*
- 9012** Misdiagnosis of pancreatic metastasis from renal cell carcinoma: A case report  
*Liang XK, Li LJ, He YM, Xu ZF*
- 9020** Discoid medial meniscus of both knees: A case report  
*Zheng ZR, Ma H, Yang F, Yuan L, Wang GD, Zhao XW, Ma LF*
- 9028** Simultaneous laparoscopic and arthroscopic excision of a huge juxta-articular ganglionic cyst compressing the sciatic nerve: A case report  
*Choi WK, Oh JS, Yoon SJ*
- 9036** One-stage revision arthroplasty in a patient with ochronotic arthropathy accompanied by joint infection: A case report  
*Wang XC, Zhang XM, Cai WL, Li Z, Ma C, Liu YH, He QL, Yan TS, Cao XW*
- 9044** Bladder paraganglioma after kidney transplantation: A case report  
*Wang L, Zhang YN, Chen GY*
- 9050** Total spinal anesthesia caused by lidocaine during unilateral percutaneous vertebroplasty performed under local anesthesia: A case report  
*Wang YF, Bian ZY, Li XX, Hu YX, Jiang L*
- 9057** Ruptured splenic artery aneurysms in pregnancy and usefulness of endovascular treatment in selective patients: A case report and review of literature  
*Lee SH, Yang S, Park I, Im YC, Kim GY*
- 9064** Gastrointestinal metastasis secondary to invasive lobular carcinoma of the breast: A case report  
*Li LX, Zhang D, Ma F*
- 9071** Post-bulbar duodenal ulcer with anterior perforation with kissing ulcer and duodenocaval fistula: A case report and review of literature  
*Alzerwi N*
- 9078** Modified orthodontic treatment of substitution of canines by first premolars: A case report  
*Li FF, Li M, Li M, Yang X*
- 9087** Renal cell carcinoma presented with a rare case of icteric Stauffer syndrome: A case report  
*Popov DR, Antonov KA, Atanasova EG, Pentchev CP, Milatchkov LM, Petkova MD, Neykov KG, Nikolov RK*
- 9096** Successful resection of a huge retroperitoneal venous hemangioma: A case report  
*Qin Y, Qiao P, Guan X, Zeng S, Hu XP, Wang B*
- 9104** Malignant transformation of biliary adenofibroma combined with benign lymphadenopathy mimicking advanced liver carcinoma: A case report  
*Wang SC, Chen YY, Cheng F, Wang HY, Wu FS, Teng LS*

- 9112 Congenital hepatic cyst: Eleven case reports  
*Du CX, Lu CG, Li W, Tang WB*
- 9121 Endovascular treatment of a ruptured pseudoaneurysm of the internal carotid artery in a patient with nasopharyngeal cancer: A case report  
*Park JS, Jang HG*
- 9127 Varicella-zoster virus meningitis after spinal anesthesia: A case report  
*Lee YW, Yoo B, Lim YH*
- 9132 Chondrosarcoma of the toe: A case report and literature review  
*Zhou LB, Zhang HC, Dong ZG, Wang CC*
- 9142 Tamsulosin-induced life-threatening hypotension in a patient with spinal cord injury: A case report  
*Lee JY, Lee HS, Park SB, Lee KH*
- 9148 CCNO mutation as a cause of primary ciliary dyskinesia: A case report  
*Zhang YY, Lou Y, Yan H, Tang H*
- 9156 Repeated bacteremia and hepatic cyst infection lasting 3 years following pancreatoduodenectomy: A case report  
*Zhang K, Zhang HL, Guo JQ, Tu CY, Lv XL, Zhu JD*
- 9162 Idiopathic cholesterol crystal embolism with atheroembolic renal disease and blue toes syndrome: A case report  
*Cheng DJ, Li L, Zheng XY, Tang SF*
- 9168 Systemic lupus erythematosus with visceral varicella: A case report  
*Zhao J, Tian M*

**LETTER TO THE EDITOR**

- 9176 Imaging of fibroadenoma: Be careful with imaging follow-up  
*Ece B, Aydin S*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Mohsen Khosravi, MD, Assistant Professor, Department of Psychiatry and Clinical Psychology, Zahedan University of Medical Sciences, Zahedan 9819713955, Iran. m.khosravi@zaums.ac.ir

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases (WJCC, World J Clin Cases)* is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

*WJCC* mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The *WJCC* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for *WJCC* as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The *WJCC*'s CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Xu Guo*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

September 6, 2022

**COPYRIGHT**

© 2022 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

# Post-bulbar duodenal ulcer with anterior perforation with kissing ulcer and duodenocaval fistula: A case report and review of literature

Nasser Alzerwi

**Specialty type:** Gastroenterology and hepatology

**Provenance and peer review:** Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): A  
Grade B (Very good): B, B  
Grade C (Good): 0  
Grade D (Fair): 0  
Grade E (Poor): 0

**P-Reviewer:** Dragonieri S, Italy; Harouachi A, Morocco; Salimi M, Iran

**Received:** April 8, 2022

**Peer-review started:** April 8, 2022

**First decision:** May 12, 2022

**Revised:** May 27, 2022

**Accepted:** July 25, 2022

**Article in press:** July 25, 2022

**Published online:** September 6, 2022



**Nasser Alzerwi**, Department of Surgery, College of Medicine, Majmaah University, Riyadh 66, Riyadh, Saudi Arabia

**Corresponding author:** Nasser Alzerwi, MS, Associate Professor, Department of Surgery, College of Medicine, Majmaah University, Ministry of Education, P. O. Box 66, Al-Majmaah City, Riyadh Region 11952, Saudi Arabia. [n.alzerwi@mu.edu.sa](mailto:n.alzerwi@mu.edu.sa)

## Abstract

### BACKGROUND

A post-bulbar duodenal ulcer (PBDU) is an ulcer in the duodenum that is distal to the duodenal bulb. PBDU may coexist with a synchronous posterior ulcer in rare occurrences, resulting in a kissing ulcer (KU). Duodenocaval fistula (DCF) is another uncommon but potentially fatal complication related to PBDU. There is limited knowledge of the scenarios in which PBDU is complicated by KU and DCF simultaneously.

### CASE SUMMARY

A 22-year-old man was admitted to the emergency department with abdominal pain, stiffness, and vomiting. The X-ray showed pneumoperitoneum, suggesting a perforated viscus. Laparotomy revealed a KU with anterior perforation and a DCF. After Kocherization, venorrhaphy was used to control caval bleeding. Due to the critical condition of the patient, only primary duodenorrhaphy with gastrojejunostomy was performed as a damage control strategy. However, later, the patient developed obstructive jaundice and leakage, and two additional jejunal perforations were detected. Due to the poor condition of the duodenum and the involvement of the ampulla in the posterior ulcer, neither primary repair nor pancreatic-free duodenectomy and ampulloplasty/ampullary reimplantation were considered viable; therefore, an emergency pancreaticoduodenectomy was performed, along with resection and anastomosis of the two jejunal perforations. The patient had a smooth recovery after surgery and was discharged after 27 d.

### CONCLUSION

The timely diagnosis of PBDU and radical surgery can aid in the smooth recovery of patients, even in the most complex cases.

**Key Words:** Duodenal ulcer; Duodenocaval fistula; Kissing ulcer; Emergency Whipple's

surgery; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** A post-bulbar duodenal ulcer, in combination with a kissing ulcer (KU) and a duodenocaval fistula (DCF), is a severe complication with a high mortality rate. In the present case, the patient had a KU with anterior perforation and a DCF. After Kocherization, venorrhaphy was used to control caval bleeding. The patient, however, later developed obstructive jaundice and leakage. Due to the poor condition of the duodenum and the involvement of the ampulla in the posterior ulcer an emergency pancreaticoduodenectomy was performed, along with resection and anastomosis of the two jejunal perforations. The patient had a smooth recovery after surgery.

**Citation:** Alzerwi N. Post-bulbar duodenal ulcer with anterior perforation with kissing ulcer and duodenocaval fistula: A case report and review of literature. *World J Clin Cases* 2022; 10(25): 9071-9077

**URL:** <https://www.wjgnet.com/2307-8960/full/v10/i25/9071.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v10.i25.9071>

## INTRODUCTION

According to current estimates, post-bulbar duodenal ulcers (PBDU) account for 5%-10% of all duodenal ulcers[1]. Most of these ulcers develop in the second part of the duodenum, with the majority occurring on the posteromedial wall of the duodenum[2]. The overall incidence of duodenal ulcers is 9.33%[3]. According to autopsy studies, 5%-20% of peptic ulcers are post-bulbar ulcers, whereas 5% of duodenal ulcers are situated distal to the bulb[4]. The clinical presentation and specific diagnostic criteria for PBDU have not yet been fully established, making diagnosis and treatment difficult[5]. Abdominal pain is the most common peptic ulcer symptom; however, melaena and hematemesis are also common[6].

PBDU, in a rare condition, can accompany a synchronous posterior ulcer and form a 'kissing' ulcer (KU)[7-10]. In the context of kissing duodenal ulcer, some may choose a vagotomy and pyloroplasty over a proximal gastric vagotomy. However, plication therapy for perforated duodenal ulcers results in postoperative bleeding problems due to KU. Bleeding in the upper gastrointestinal tract makes duodenal bulb ulcers worse and less responsive to treatment[11]. Any instance of symptomless hematemesis should be thoroughly explored for post-bulbar ulcers. Furthermore, since these abnormalities may overlap with presentations commonly seen in Zollinger Ellison syndrome (ZES), a precise diagnosis and treatment strategy become critical[12].

Another severe and rare complication of PBDU is the duodenocaval fistula (DCF). DCF is an uncommon but severe form of digestive fistula. It develops due to problems with the duodenum-inferior vena cava junction. Nontraumatic DCF is sporadic and may be caused by various factors such as penetrating duodenal peptic ulcers, foreign bodies, malignancies, right nephrectomy, and radiation therapy to the upper abdomen. DCF is often distinguished by gastrointestinal bleeding; however, it can also be accompanied by fever and infection. DCF should be approached cautiously, and decisions should be made quickly since it is associated with a high mortality rate before surgical intervention[13].

Several cases on the treatment of PBDU and when KU or DCF complicates PBDU have been documented in the literature. However, PBDU in conjunction with KU and DCF is not adequately described in the literature. This study presents the successful management of a rare case of PBDU complicated by KU and DCF, providing a detailed overview of diagnostic problems and surgical complications. A brief account of the previous research published in this field is also included.

## CASE PRESENTATION

### Chief complaints

Stomach discomfort and vomiting for two days.

### History of present illness

On January 24, 2019, a 22-year-old man came to the emergency department with stomach discomfort and vomiting for two days. He appeared to be stressed due to acute dehydration.

**History of past illness**

The patient was taking non-steroidal anti-inflammatory drugs with no history of peptic ulcer disease. He was hypotensive and tachycardic, with blood pressure (BP) of 100/70 mmHg and a pulse rate of 110 beats per minute.

**Personal and family history**

There was no family history relevant to this case.

**Physical examination**

Physical examination indicated board-like stiffness of the abdomen, and auscultation revealed a negative bowel sound.

**Laboratory examinations**

See [Table 1](#) for lab results at admission.

**Imaging examinations**

An erect chest radiograph revealed air under the diaphragm.

---

**FINAL DIAGNOSIS**

---

Following resuscitation, a nasogastric tube (NG) and a foley catheter were placed, and the patient became anuric. He underwent an exploratory laparotomy, which revealed a KU, a post bulbar ulcer in the second part of the duodenum with a severely deformed and fibrotic duodenum, and an anterior ulcer that was perforated and obstructed by clots ([Figure 1](#), [Video 1](#)).

---

**TREATMENT**

---

Due to the penetration of the posterior ulcer into the inferior vena cava (DCF), which was in contact with the ampulla, the duodenum was filled with venous blood. Duodenotomy was performed and pressure was applied to achieve temporary control of the source until the blood transfusion. Duodenal Kocherization was performed and after proximal and distal control of the inferior vena cava with sponge-on-sticks, caval hole venorrhaphy was conducted. The patient received the equivalent of 2 total blood volumes of packed red blood cell, platelets, and fresh-frozen plasma. Due to the critical condition of the patient, a damage control approach was used. Gastrojejunostomy was conducted after primary repair (transverse duodenorrhaphy/duodenoplasty of the duodenotomy that included both ulcers) as a bypass procedure due to narrowing of the duodenal. A drain was left in the subhepatic space, and the patient developed obstructive jaundice and biliary leakage from the drain while in the intensive care unit (ICU). He was brought for re-exploration, which revealed two additional perforations in the jejunum. Due to the poor condition of the duodenum and the involvement of the ampulla in the posterior ulcer, neither primary repair nor pancreas sparing duodenectomy and ampullectomy/ampullary reimplantation were viable options. An emergency Whipple procedure (pancreaticoduodenectomy) was performed, along with resection and anastomosis of the jejunal perforation.

---

**OUTCOME AND FOLLOW-UP**

---

The patient recovered quickly from the surgery. However, he experienced two episodes of aspiration pneumonia, for which he was intubated and ventilated in the ICU. The dehiscence of the wound developed as a result of renal failure. On the eighth postoperative day, a computed tomography (CT) scan revealed no collection or leakage. Every other day, the drain effluent was sent for analysis of amylase and bilirubin, and white cell count and only turbid fluid was found, without amylase leakage or pus. On the 12<sup>th</sup> postoperative day, the patient was extubated and kept in the ICU with 2 Liters of O<sub>2</sub> through a nasal cannula saturating 98%, with vitals of BP: 137/70 mmHg, pulse: 104 beats per minute, and temperature: 37.8 °C. The last culture from the chest revealed Klebsiella pneumoniae, and Vancomycin-resistant enterococci. Other laboratory findings are given in [Table 1](#).

On day 13 after the operation, the patient was fed through NG and tolerated well glasgow coma scale of 13/15. Evaluation of anastomotic leakage (enteric, biliary, or pancreatic) by abdominal CT revealed no leak or collection. Tigecycline, Colistin, and Imipenem were prescribed to the patient. After 27 d, the patient was discharged with a good clinical condition.

**Table 1 Laboratory parameters of the patient at admission**

Parameter	Value
Hite blood count	19
Hemoglobin	9.2
Creatinine	230
Urea	15
Sodium	146
Albumin	31
Alanine transaminase	29
Aspartate transferase	42
Total bilirubin	30



DOI: 10.12998/wjcc.v10.i25.9071 Copyright ©The Author(s) 2022.

**Figure 1 Exploratory laparotomy revealed post-bulbar duodenal ulcer.** It can be seen that the anterior ulcer perforation is plugged with a clot.

## DISCUSSION

Early diagnosis and successful management of PBDU present several challenges. The risk of mortality and morbidity increases many folds when KU and DCF complicate the condition; however, little is understood about the clinical signs, diagnosis, and management when all these complexities (PBDU, KU, and DCF) arise simultaneously. This study unveiled numerous critical aspects of this complex condition by offering a comprehensive discussion on the successful treatment of a patient with PBDU, KU, and DCF. This case presented several diagnostic and clinical challenges, and the patient was subjected to surgical treatment with caval venorrhaphy, pancreaticoduodenectomy, segmental enterectomy, and jejunojejunostomy, which resolved KU and DCF and jejunal perforations.

Perforations in the duodenum can be free or limited. The term "free perforation" refers to the time when the intestinal material seeps into the abdominal cavity, resulting in diffuse peritonitis. Limited perforation occurs when an ulcer produces a full-thickness hole, but the open leaking is blocked by surrounding organs such as the pancreas. Although patients with duodenal perforation require surgical treatment in most cases, in patients with perivaterian injuries, conservative management can also produce successful outcomes. In acute duodenal perforations, Whipple surgery is highly complicated [14,15]; therefore, in our case, radical surgery differed initially. However, when pancreatic-free duodenectomy and ampuloplasty/ampullary reimplantation became unfeasible, emergent Whipple surgery was successfully performed, reflecting the feasibility of radical approaches in the treatment of such complex cases.

The most common indication for surgery is bleeding, and surgical options include stomach resection +/- vagotomy. The mortality rate is 2.6%, except in fistulous cases, which have a mortality rate of 7.7%. Of note, when complicated with KS, the mortality rate in bleeders is reported to be as high as 50% [16]. Therefore, in events of gastrointestinal blood loss in a perforated duodenal ulcer, an intraoperative search for a posterior KS should be considered. If a KU was found, an acid-reducing operation and suture ligation are viable approaches [7].

Compared to bulbar and gastric ulcers, the frequency of bleeding in PBDU is approximately twice as high. PBDU appears later in life and affects men more than women [17]; however, unlike bulbar duodenal ulcers, patients with PBDU tend to show severe acidity, a persistent pattern of acid secretion,

and different clinical and radiographic findings[18]. Postoperative mortality from PBDU is greater than bulbar duodenal ulcer[3]. In a study from India that provided a detailed description of the aspects associated with duodenal ulcer, post-bulbar ulcer *vs* bulbar ulcer was found to be 1:1.5, with deformed bulbs observed in half of the cases[17]. Smokers accounted for 42% of patients with duodenal ulcers, while tobacco chewers accounted for 15% and alcoholism for 18%. Consumption of tea, rice, and spices has also been associated with increased acid secretion and the development of duodenal ulcers. Most importantly, *H. Pylori* was estimated to infect 80% of the population. It may be noted that, unlike duodenal bulb ulcer, in certain cases, PBDU did not heal with *Helicobacter pylori* eradication therapy, suggesting that post-bulbar ulcer etiologically differs from bulbar ulcer[19]. Notably, *H. pylori* was also implicated in the development of KU in the duodenal bulb[9].

In 1989, the first recorded instance of a duodenal ulcer associated with pentagastrin-fast achlorhydria was described[20]; wherein a 55-year-old male was diagnosed with a post bulbar duodenal ulcer, hemorrhage, and fasting hypergastrinemia, and ulcer healing was reported in eight weeks, after antrectomy, vagotomy, gastrojejunostomy, and a course of sucralfate medication. In another study, 12 patients were reported to have PBDU[21]. In ten patients, a truncal vagotomy and outlet surgery were performed and the anterior duodenotomy extended to the stenotic region. A Jaboulay gastroduodenostomy was performed in one patient, while a pyloroplasty duodenoplasty was performed in another. There were no deaths among these individuals during the six-year follow-up period. However, three individuals had recurrences of peptic ulcers, which could be related to undetected post-bulbar stenosis after surgery. This showed that to detect the ulcer, the duodenum must be checked by intraluminal palpation in all patients undergoing surgery for peptic ulcer disease.

Notably, in 121 individuals with PBDU, 72.7% had duodenal stenosis, 41.3% had penetration, and 5.8% had choledochoduodenal fistulas[22]. In approximately 40% of the patients, the progress of the ulcer was exacerbated by bleeding and in 8.3% by perforation. In 34 patients, a selective proximal vagotomy was performed, and in 28 cases, a selective proximal vagotomy and a draining operation were performed. In 41 patients, proximal vagotomy and duodenoplasty were performed. The use of selective proximal vagotomy and duodenoplasty produced better results. There were no deaths or serious complications. In contrast, a comprehensive case review of 1087 patients with PBDU showed complications in 1014 of them; importantly, penetration occurred in 707 individuals in conjunction with hemorrhage from stenosis[23]. Organ saving techniques combined with vagotomy have been shown to offer benefits over resection techniques[24]. After surgery, therapeutic outcomes can vary depending on the patient's predisposition to the post-bulbar lesion[16].

Our patient also had DCF, a type of intestinal fistula that is a fatal clinical entity with a high death rate prior to definitive therapy[13]. If DCF is clinically suspected, the first-line study should be CT and magnetic resonance imaging, with a thorough evaluation of images of inferior vena cava (IVC) and surrounding structures. The prognosis is decided by early diagnosis and surgery before bleeding starts. DCF treatment includes fistula closure, as well as duodenal and IVC repair[25]. This argument is based on a postmortem examination of a 54-year-old man, which found that the cause of death was upper gastrointestinal hemorrhage, resulting in a considerable volume of blood in the intestinal lumen due to gastric ulcer rupture in the IVC. In our case, DCF management involved surgical intervention, which was decided after a meticulous assessment of the extent of damage to the duodenum and IVC. Barloon *et al*[26] reported a morbidity rate of 61% after radical surgery. To prevent fistula recurrences, surgery-butressed repair, such as a jejunal patch or an epiploic flap, is generally preferred; however, other approaches, such as truncal vagotomy, antrectomy, and/or duodenal exclusion, can be utilized in combination, particularly in the case of a peptic ulcer. Conservative surgical therapy was also successfully applied to a 73-year-old man with DCF, occult intestinal bleeding, and sepsis[27]. In this case, the IVC and duodenum were sutured after sharp dissection, duodenal mobilization, and digital control of hemorrhage. The duodenal exclusion was performed using gastroenterostomy and truncal vagotomy of antral stapling and interposition of the epiploic patch to prevent recurrence of the fistula [27].

The presence of a gastric ulcer with a fistula in the IVC and food embolization in the lung has also been documented, indicating the severity of the condition[28]. The first case of DCF caused by a large descending duodenal peptic ulcer was described in 1990[29]. DCF hemorrhage was contained by direct compression above and below the fistula in another example involving a 49-year-old male, and the IVC defect was repaired with 5/0 prolene[30]. In 1996, a case of polymicrobial fungemia and fatal gastro-intestinal hemorrhage associated with DCF caused by a peptic ulcer was reported. This raised the possibility of candidal endocarditis[31]. The first case of DCF with peptic ulcer showed complications due to embolization of the intestinal contents in the lung, with numerous intravascular mucin[32]. In 2005, a case of a 44-year-old patient with DCF was also reported. The patient had no history of peptic ulcer disease. Septic shock preceded hemorrhagic shock as a clinical characteristic, but only after laparotomy was the diagnosis established[33]. Finally, in our case, the patient developed jaundice, which is rare, with a prevalence of 0.14% [34]. The development of jaundice is due to the close relationship of the common bile duct with the second part of the duodenum. It may be noted that though our case provides vital information on the management of PBDU, KU, and DCF, more studies are needed to standardize the diagnostic and therapeutic approaches.

## CONCLUSION

PBDUs generally involve a surgical emergency and are difficult to diagnose and manage. This rare case highlights that PBDU should be the first of surgical intent and a radical procedure should be considered if viable. In our case, only primary repair of the two ulcers was performed along with a proximal drainage procedure (gastrojejunostomy); however, when additional perforations were detected and conservative measurements were not viable, emergent Whipple surgery was successfully performed. The duodenocaval fistula, in particular, is difficult to diagnose in this situation and is linked with a significant mortality rate before conclusive therapeutic efforts. Our case demonstrates that a careful diagnosis and timely treatment will be helpful for patient recovery and a good prognosis.

## ACKNOWLEDGEMENTS

Author thanks the patient and his family for valuable support.

## FOOTNOTES

**Author contributions:** Alzerwi N conducted case analysis, literature review, and manuscript writing.

**Informed consent statement:** A written informed consent was obtained from the patient for publication of this case report.

**Conflict-of-interest statement:** All the authors have nothing to disclose.

**CARE Checklist (2016) statement:** The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** Saudi Arabia

**ORCID number:** Nasser alzerwi [0000-0002-4877-2152](https://orcid.org/0000-0002-4877-2152).

**S-Editor:** Wang DM

**L-Editor:** A

**P-Editor:** Wang DM

## REFERENCES

- 1 **Matsuhashi T**, Fukuda S, Abe Y, Mikami T, Tatsuta T, Hikichi T, Nakamura J, Onozato Y, Hatta W, Masamune A, Ohyauchi M, Ito H, Hanabata N, Araki Y, Yanagita T, Imamura H, Tsuji T, Sugawara K, Horikawa Y, Ohara S, Kondo Y, Dohmen T, Iijima K. Nature and treatment outcomes of bleeding post-bulbar duodenal ulcers. *Dig Endosc* 2022; **34**: 984-993 [PMID: [34609030](https://pubmed.ncbi.nlm.nih.gov/34609030/) DOI: [10.1111/den.14160](https://doi.org/10.1111/den.14160)]
- 2 **Britt LG**, Wolf RY. Postbulbar ulcer with retrocecal abscess. A case report. *Arch Surg* 1966; **92**: 98-100 [PMID: [5901269](https://pubmed.ncbi.nlm.nih.gov/5901269/) DOI: [10.1001/archsurg.1966.01320190100023](https://doi.org/10.1001/archsurg.1966.01320190100023)]
- 3 **Ghelase F**, Georgescu I, Ghelase S, Mărgăritescu D, Sandu E, Cioară F, Cioară D, Bratiloveanu T. [Duodenal post-bulbar ulcers. Diagnostic and therapeutic problems]. *Chirurgia (Bucur)* 2006; **101**: 47-53 [PMID: [16623377](https://pubmed.ncbi.nlm.nih.gov/16623377/)]
- 4 **Kaufman SA**, LEVENE G. Postbulbar duodenal ulcer. *Radiology* 1957; **69**: 848-852 [PMID: [13494714](https://pubmed.ncbi.nlm.nih.gov/13494714/) DOI: [10.1148/69.6.848](https://doi.org/10.1148/69.6.848)]
- 5 **Ramsdell JA**, Bartholomew LG, Cain JC, Davis GD. Postbulbar duodenal ulcer. *Ann Intern Med* 1957; **47**: 700-710 [PMID: [13470696](https://pubmed.ncbi.nlm.nih.gov/13470696/) DOI: [10.7326/0003-4819-47-4-700](https://doi.org/10.7326/0003-4819-47-4-700)]
- 6 **Samuel E**. Post-bulbar duodenal ulcers. *S Afr Med J* 1952; **26**: 121-125 [PMID: [14913269](https://pubmed.ncbi.nlm.nih.gov/14913269/)]
- 7 **Stabile BE**, Hardy HJ, Passaro E Jr. 'Kissing' duodenal ulcers. *Arch Surg* 1979; **114**: 1153-1156 [PMID: [485825](https://pubmed.ncbi.nlm.nih.gov/485825/) DOI: [10.1001/archsurg.1979.01370340059010](https://doi.org/10.1001/archsurg.1979.01370340059010)]
- 8 **Feliciano DV**. Do perforated duodenal ulcers need an acid-decreasing surgical procedure now that omeprazole is available? *Surg Clin North Am* 1992; **72**: 369-380 [PMID: [1549799](https://pubmed.ncbi.nlm.nih.gov/1549799/) DOI: [10.1016/s0039-6109\(16\)45684-7](https://doi.org/10.1016/s0039-6109(16)45684-7)]
- 9 **Avramenko AA**. The case of formation of «kissing» ulcers of duodenal bulb of the patient with chronic nonatrophic gastritis on the background of the eating of a vegetable salad and physical exertion. *Wiad Lek* 2019; **72**: 946-949 [PMID: [31811111](https://pubmed.ncbi.nlm.nih.gov/31811111/)]

- 31175802]
- 10 **Avramenko AA**. Influence of peculiarities of anatomical structure (SHAPE) of duodenal bulb on the mechanism of formation of "kissing" ulcers of patients with chronic non-atrophic gastritis. *Wiad Lek* 2020; **73**: 2568-2571 [PMID: 33577469]
  - 11 **Hunt PS**, Clarke G. Perforation in patients with bleeding ulcer. *Aust N Z J Surg* 1991; **61**: 183-185 [PMID: 2003835 DOI: 10.1111/j.1445-2197.1991.tb07589.x]
  - 12 **Rossi RE**, Elvevi A, Citterio D, Coppa J, Invernizzi P, Mazzaferro V, Massironi S. Gastrinoma and Zollinger Ellison syndrome: A roadmap for the management between new and old therapies. *World J Gastroenterol* 2021; **27**: 5890-5907 [PMID: 34629807 DOI: 10.3748/wjg.v27.i35.5890]
  - 13 **Rheudasil JM**, Chuang VP, Amerson JR. Duodenocaval fistula: case report and literature review. *Am Surg* 1988; **54**: 169-171 [PMID: 3348551]
  - 14 **Standop J**, Glowka T, Schmitz V, Schaefer N, Hirner A, Kalff JC. Emergency Kausch-Whipple procedure: indications and experiences. *Pancreas* 2010; **39**: 156-159 [PMID: 20182309 DOI: 10.1097/MPA.0b013e3181bb98d2]
  - 15 **Varshney VK**, Nayar R, Sreesanth KS, Soni S, Varshney B. Emergency Pancreatoduodenectomy for Ampullary Cancer Post-Iatrogenic Duodenal Perforation: No Option but to Strike. *Cureus* 2020; **12**: e11384 [PMID: 33312785 DOI: 10.7759/cureus.11384]
  - 16 **Mateş IN**, Constantinoiu S. [Postbulbar ulcer. An eclectic surgical approach]. *Chirurgia (Bucur)* 1997; **92**: 47-57 [PMID: 9296747]
  - 17 **Rao SS**, Murthy KV. Post-bulbar and coexisting ulceration: unique features of peptic ulcer in Hyderabad. *Gut* 1993; **34**: 1327-1330 [PMID: 8244096 DOI: 10.1136/gut.34.10.1327]
  - 18 **Kaminishi M**, Shimazu R, Kuramoto S, Sadatsuki H, Johjima Y, Oohara T. A clinicopathological study of post-bulbar duodenal ulcer--analysis of 6 cases and comparison to usual chronic duodenal ulcer. *Jpn J Surg* 1987; **17**: 140-145 [PMID: 3626205 DOI: 10.1007/BF02470655]
  - 19 **Yamane T**, Uchiyama K, Ishii T, Omura M, Fujise K, Tajiri H. Two cases of refractory post-bulbar duodenal ulcer. *Intern Med* 2007; **46**: 1413-1417 [PMID: 17827841 DOI: 10.2169/internalmedicine.46.0113]
  - 20 **Goldschmiedt M**, Peterson WL, Vuitch F, Feldman M. Postbulbar duodenal ulcer in a patient with pentagastrin-fast achlorhydria. *Gastroenterology* 1989; **97**: 771-774 [PMID: 2753337 DOI: 10.1016/0016-5085(89)90652-5]
  - 21 **Hines JR**, Geurkink RE, Gordon RT. Postbulbar peptic ulceration of the duodenum. *Surg Gynecol Obstet* 1976; **142**: 13-15 [PMID: 1244687]
  - 22 **Posmelov VS**, Kubyshkin VA, Kozlov IA. [Surgical treatment of postbulbar duodenal ulcers]. *Khirurgiia (Mosk)* 1997; **5**: 4-9 [PMID: 9297023]
  - 23 **Lobzhanidze GV**. Clinical features and surgical treatment of postbulbar ulcers of the duodenum. *Khirurgiia (Mosk)* 1989; **10**: 29-34 [PMID: 2593568]
  - 24 **Kurbanov KM**. Surgical treatment of postbulbar ulcers, complicated by penetration into pancreatic head. *Khirurgiia (Mosk)* 1999; **2**: 8-10 [PMID: 10081244]
  - 25 **Guo Y**, Zhang YQ, Lin W. Radiological diagnosis of duodenocaval fistula: a case report and literature review. *World J Gastroenterol* 2010; **16**: 2314-2316 [PMID: 20458773 DOI: 10.3748/wjg.v16.i18.2314]
  - 26 **Barloon TJ**, Lu CH, Honda H, Walker WP, Murray J. Primary adenocarcinoma of the duodenal bulb: radiographic and pathologic findings in two cases. *Gastrointest Radiol* 1989; **14**: 223-225 [PMID: 2731695 DOI: 10.1007/BF01889202]
  - 27 **Guillem PG**, Binot D, Dupuy-Cuny J, Laberrenne JE, Lesage J, Triboulet JP, Chambon JP. Duodenocaval fistula: a life-threatening condition of various origins. *J Vasc Surg* 2001; **33**: 643-645 [PMID: 11241139 DOI: 10.1067/mva.2001.111741]
  - 28 **Palmer RC**. Vena caval penetration by gastric ulcer: massive hemorrhage and embolization of gastric contents to lungs. *J Clin Gastroenterol* 1989; **11**: 455-457 [PMID: 2760436 DOI: 10.1097/00004836-198908000-00022]
  - 29 **deSa LA**, Roddie ME, Williamson RC. Fatal duodenocaval fistula resulting from a giant peptic ulcer. Case report. *Acta Chir Scand* 1990; **156**: 647-650 [PMID: 2264448]
  - 30 **Geraghty JG**, Coveney E, Kennedy TE, O'Dwyer PJ, Murphy JJ. Duodenocaval fistula in peptic ulceration. *Gut* 1991; **32**: 452-453 [PMID: 2026346 DOI: 10.1136/gut.32.4.452]
  - 31 **Lesho EP**, LeBrun C, Landry FJ, Tsuchida A, Cooper RH. Fatal duodenocaval fistula caused by peptic ulcer. *South Med J* 1996; **89**: 925-926 [PMID: 8790321 DOI: 10.1097/00007611-199609000-00017]
  - 32 **Cyr PV**, Shuhaibar H, Kay JM. Spontaneous duodenal-caval fistula with embolization of intestinal contents. *Hum Pathol* 1998; **29**: 1165-1166 [PMID: 9781659 DOI: 10.1016/s0046-8177(98)90431-2]
  - 33 **Ousmane ML**, Herbecq P, Nyunga M, Vignozzi G, Lemaire C. [Duodenocaval fistula: an unusual form of peptic ulcer revelation]. *Ann Fr Anesth Reanim* 2005; **24**: 1294-1296 [PMID: 16183244 DOI: 10.1016/j.annfar.2005.04.030]
  - 34 **Hansen EF**, Jensen HQ. [Postbulbar duodenal ulcer. A rare cause of icterus]. *Ugeskr Laeger* 1996; **158**: 3947-3948 [PMID: 8701513]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA  
**Telephone:** +1-925-3991568  
**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
**Help Desk:** <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>

