

# World Journal of *Clinical Cases*

*World J Clin Cases* 2022 November 6; 10(31): 11214-11664



## Contents

Thrice Monthly Volume 10 Number 31 November 6, 2022

### REVIEW

- 11214** Diabetes and skin cancers: Risk factors, molecular mechanisms and impact on prognosis  
*Dobrică EC, Banciu ML, Kipkorir V, Khazeei Tabari MA, Cox MJ, Simhachalam Kutikuppala LV, Găman MA*
- 11226** Endocrine disruptor chemicals as obesogen and diabetogen: Clinical and mechanistic evidence  
*Kurşunoğlu NE, Sarer Yurekli BP*
- 11240** Intestinal microbiota in the treatment of metabolically associated fatty liver disease  
*Wang JS, Liu JC*

### MINIREVIEWS

- 11252** Lactation mastitis: Promising alternative indicators for early diagnosis  
*Huang Q, Zheng XM, Zhang ML, Ning P, Wu MJ*
- 11260** Clinical challenges of glycemic control in the intensive care unit: A narrative review  
*Sreedharan R, Martini A, Das G, Aftab N, Khanna S, Ruetzler K*
- 11273** Concise review on short bowel syndrome: Etiology, pathophysiology, and management  
*Lakkasani S, Seth D, Khokhar I, Touza M, Dacosta TJ*
- 11283** Role of nickel-regulated small RNA in modulation of *Helicobacter pylori* virulence factors  
*Freire de Melo F, Marques HS, Fellipe Bueno Lemos F, Silva Luz M, Rocha Pinheiro SL, de Carvalho LS, Souza CL, Oliveira MV*
- 11292** Surgical intervention for acute pancreatitis in the COVID-19 era  
*Su YJ, Chen TH*

### ORIGINAL ARTICLE

#### Clinical and Translational Research

- 11299** Screening of traditional Chinese medicine monomers as ribonucleotide reductase M2 inhibitors for tumor treatment  
*Qin YY, Feng S, Zhang XD, Peng B*

#### Case Control Study

- 11313** Covered transjugular intrahepatic portosystemic stent-shunt *vs* large volume paracentesis in patients with cirrhosis: A real-world propensity score-matched study  
*Dhaliwal A, Merhzad H, Karkhanis S, Tripathi D*

**Retrospective Cohort Study**

- 11325** Endoscopic submucosal tunnel dissection for early esophageal squamous cell carcinoma in patients with cirrhosis: A propensity score analysis  
*Zhu LL, Liu LX, Wu JC, Gan T, Yang JL*

**Retrospective Study**

- 11338** Nomogram for predicting overall survival in Chinese triple-negative breast cancer patients after surgery  
*Lin WX, Xie YN, Chen YK, Cai JH, Zou J, Zheng JH, Liu YY, Li ZY, Chen YX*
- 11349** Early patellar tendon rupture after total knee arthroplasty: A direct repair method  
*Li TJ, Sun JY, Du YQ, Shen JM, Zhang BH, Zhou YG*
- 11358** Coxsackievirus A6 was the most common enterovirus serotype causing hand, foot, and mouth disease in Shiyang City, central China  
*Li JF, Zhang CJ, Li YW, Li C, Zhang SC, Wang SS, Jiang Y, Luo XB, Liao XJ, Wu SX, Lin L*
- 11371** Dynamic changes of estimated glomerular filtration rate are conversely related to triglyceride in non-overweight patients  
*Liu SQ, Zhang XJ, Xue Y, Huang R, Wang J, Wu C, He YS, Pan YR, Liu LG*
- 11381** C-reactive protein as a non-linear predictor of prolonged length of intensive care unit stay after gastrointestinal cancer surgery  
*Yan YM, Gao J, Jin PL, Lu JJ, Yu ZH, Hu Y*

**Clinical Trials Study**

- 11391** Dan Bai Xiao Formula combined with glucocorticoids and cyclophosphamide for pediatric lupus nephritis: A pilot prospective study  
*Cao TT, Chen L, Zhen XF, Zhao GJ, Zhang HF, Hu Y*

**Observational Study**

- 11403** Relationship between lipids and sleep apnea: Mendelian randomization analysis  
*Zhang LP, Zhang XX*
- 11411** Efficacy and safety profile of two-dose SARS-CoV-2 vaccines in cancer patients: An observational study in China  
*Cai SW, Chen JY, Wan R, Pan DJ, Yang WL, Zhou RG*

**Prospective Study**

- 11419** Pressure changes in tapered and cylindrical shaped cuff after extension of head and neck: A randomized controlled trial  
*Seol G, Jin J, Oh J, Byun SH, Jeon Y*

**Randomized Controlled Trial**

- 11427** Effect of intradermal needle therapy at combined acupoints on patients' gastrointestinal function following surgery for gastrointestinal tumors  
*Guo M, Wang M, Chen LL, Wei FJ, Li JE, Lu QX, Zhang L, Yang HX*

## SYSTEMATIC REVIEWS

- 11442** Video-assisted bystander cardiopulmonary resuscitation improves the quality of chest compressions during simulated cardiac arrests: A systemic review and meta-analysis

*Pan DF, Li ZJ, Ji XZ, Yang LT, Liang PF*

## META-ANALYSIS

- 11454** Efficacy of the femoral neck system in femoral neck fracture treatment in adults: A systematic review and meta-analysis

*Wu ZF, Luo ZH, Hu LC, Luo YW*

- 11466** Prevalence of polymyxin-induced nephrotoxicity and its predictors in critically ill adult patients: A meta-analysis

*Wang JL, Xiang BX, Song XL, Que RM, Zuo XC, Xie YL*

## CASE REPORT

- 11486** Novel compound heterozygous variants in the LHX3 gene caused combined pituitary hormone deficiency: A case report

*Lin SZ, Ma QJ, Pang QM, Chen QD, Wang WQ, Li JY, Zhang SL*

- 11493** Fatal bleeding due to an aorto-esophageal fistula: A case report and literature review

*Ćeranić D, Nikolić S, Lučev J, Slanić A, Bujas T, Ocepek A, Skok P*

- 11500** Tolvaptan ameliorated kidney function for one elderly autosomal dominant polycystic kidney disease patient: A case report

*Zhou L, Tian Y, Ma L, Li WG*

- 11508** Extensive right coronary artery thrombosis in a patient with COVID-19: A case report

*Dall'Orto CC, Lopes RPF, Cancela MT, de Sales Padilha C, Pinto Filho GV, da Silva MR*

- 11517** Yokoyama procedure for a woman with heavy eye syndrome who underwent multiple recession-resection operations: A case report

*Yao Z, Jiang WL, Yang X*

- 11523** Rectal cancer combined with abdominal tuberculosis: A case report

*Liu PG, Chen XF, Feng PF*

- 11529** Malignant obstruction in the ileocecal region treated by self-expandable stent placement under the fluoroscopic guidance: A case report

*Wu Y, Li X, Xiong F, Bao WD, Dai YZ, Yue LJ, Liu Y*

- 11536** Granulocytic sarcoma with long spinal cord compression: A case report

*Shao YD, Wang XH, Sun L, Cui XG*

- 11542** Aortic dissection with epileptic seizure: A case report

*Zheng B, Huang XQ, Chen Z, Wang J, Gu GF, Luo XJ*



- 11549** Multiple bilateral and symmetric C1-2 ganglioneuromas: A case report  
*Wang S, Ma JX, Zheng L, Sun ST, Xiang LB, Chen Y*
- 11555** Acute myocardial infarction due to Kounis syndrome: A case report  
*Xu GZ, Wang G*
- 11561** Surgical excision of a large retroperitoneal lymphangioma: A case report  
*Park JH, Lee D, Maeng YH, Chang WB*
- 11567** Mass-like extragonadal endometriosis associated malignant transformation in the pelvis: A rare case report  
*Chen P, Deng Y, Wang QQ, Xu HW*
- 11574** Gastric ulcer treated using an elastic traction ring combined with clip: A case report  
*Pang F, Song YJ, Sikong YH, Zhang AJ, Zuo XL, Li RY*
- 11579** Novel liver vein deprivation technique that promotes increased residual liver volume (with video): A case report  
*Wu G, Jiang JP, Cheng DH, Yang C, Liao DX, Liao YB, Lau WY, Zhang Y*
- 11585** Linear porokeratosis of the foot with dermoscopic manifestations: A case report  
*Yang J, Du YQ, Fang XY, Li B, Xi ZQ, Feng WL*
- 11590** Primary hepatic angiosarcoma: A case report  
*Wang J, Sun LT*
- 11597** Hemorrhagic shock due to ruptured lower limb vascular malformation in a neurofibromatosis type 1 patient: A case report  
*Shen LP, Jin G, Zhu RT, Jiang HT*
- 11607** Gastric linitis plastica with autoimmune pancreatitis diagnosed by an endoscopic ultrasonography-guided fine-needle biopsy: A case report  
*Sato R, Matsumoto K, Kanzaki H, Matsumi A, Miyamoto K, Morimoto K, Terasawa H, Fujii Y, Yamazaki T, Uchida D, Tsutsumi K, Horiguchi S, Kato H*
- 11617** Favorable response of primary pulmonary lymphoepithelioma-like carcinoma to sintilimab combined with chemotherapy: A case report  
*Zeng SY, Yuan J, Lv M*
- 11625** Benign paroxysmal positional vertigo with congenital nystagmus: A case report  
*Li GF, Wang YT, Lu XG, Liu M, Liu CB, Wang CH*
- 11630** Secondary craniofacial necrotizing fasciitis from a distant septic emboli: A case report  
*Lee DW, Kwak SH, Choi HJ*
- 11638** Pancreatic paraganglioma with multiple lymph node metastases found by spectral computed tomography: A case report and review of the literature  
*Li T, Yi RQ, Xie G, Wang DN, Ren YT, Li K*

- 11646** Apnea caused by retrobulbar anesthesia: A case report  
*Wang YL, Lan GR, Zou X, Wang EQ, Dai RP, Chen YX*
- 11652** Unexplained septic shock after colonoscopy with polyethylene glycol preparation in a young adult: A case report  
*Song JJ, Wu CJ, Dong YY, Ma C, Gu Q*
- 11658** Metachronous isolated penile metastasis from sigmoid colon adenocarcinoma: A case report  
*Yin GL, Zhu JB, Fu CL, Ding RL, Zhang JM, Lin Q*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Muhammad Hamdan Gul, MD, Assistant Professor, Department of Internal Medicine, University of Kentucky, Chicago, IL 60657, United States.  
hamdan3802@hotmail.com

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Xu Guo*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

**EDITORIAL BOARD MEMBERS**

<https://www.wjnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

November 6, 2022

**COPYRIGHT**

© 2022 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

# Favorable response of primary pulmonary lymphoepithelioma-like carcinoma to sintilimab combined with chemotherapy: A case report

Shu-Ying Zeng, Jin Yuan, Min Lv

**Specialty type:** Oncology

**Provenance and peer review:**

Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): A, A

Grade B (Very good): 0

Grade C (Good): 0

Grade D (Fair): D

Grade E (Poor): 0

**P-Reviewer:** Mazilu L, Romania;  
Rizzo A, Italy

**Received:** July 14, 2022

**Peer-review started:** July 14, 2022

**First decision:** August 1, 2022

**Revised:** August 19, 2022

**Accepted:** September 21, 2022

**Article in press:** September 21, 2022

**Published online:** November 6, 2022



**Shu-Ying Zeng, Jin Yuan, Min Lv**, Department of Oncology, The First People's Hospital of Longquanyi District Chengdu, Chengdu 652532, Sichuan Province, China

**Corresponding author:** Shu-Ying Zeng, CCST, Doctor, MSc, PhD, Doctor, Department of Oncology, The First People's Hospital of Longquanyi District Chengdu, No. 39 Mudan Street, Jinjiang District, Chengdu 652532, Sichuan Province, China. [18982193611@163.com](mailto:18982193611@163.com)

## Abstract

### BACKGROUND

There is no established treatment for primary pulmonary lymphoepithelioma-like carcinoma (LELC) until now.

### CASE SUMMARY

In this study, the patient responded well to sintilimab combined with paclitaxel and carboplatin, showing no obvious side effects. Meantime, the values of carbohydrate antigen 15-3 (CA15-3) and carbohydrate antigen 72-4 (CA72-4) gradually returned to normal.

### CONCLUSION

Immunotherapy combined with chemotherapy in advanced-stage LELC may be more effective than immunotherapy or chemotherapy alone. CA15-3 and CA72-4 are biomarkers for evaluating therapeutic effects for LELC.

**Key Words:** Lymphoepithelioma-like carcinoma; Non-small cell lung cancer; Immunotherapy; Sintilimab; Programmed death-1; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Herein, we present the case of a lymphoepithelioma-like carcinoma (LELC) patient who was successfully treated with sintilimab combined with paclitaxel and carboplatin. Recent literature on LELC is also reviewed and discussed for future avenues for LELC research.

**Citation:** Zeng SY, Yuan J, Lv M. Favorable response of primary pulmonary lymphoepithelioma-like carcinoma to sintilimab combined with chemotherapy: A case report. *World J Clin Cases* 2022; 10(31): 11617-11624

**URL:** <https://www.wjgnet.com/2307-8960/full/v10/i31/11617.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v10.i31.11617>

## INTRODUCTION

Lymphoepithelioma-like carcinoma (LELC) is a rare subtype of NSCLC and first identified as a unique Epstein-Barr virus (EBV)-associated tumor by Bégin *et al*[1]. EBV infection is generally acknowledged as the most important etiology for LELC. Multi-dimensional comparison has revealed that pulmonary LELC resembles nasopharyngeal carcinoma (NPC) but is clearly different from other lung cancers[2]. Most LELC cases have been documented in Southeast Asia, including Hong Kong, Taiwan, Guangdong, and Singapore, with sporadic cases having been seen in the West[3]. Currently, specific treatment guidelines for LELC have not been established[3]. In recent years, dramatic progress has been made in immunotherapy for cancer.

Herein, we present the case of a LELC patient who was successfully treated with sintilimab combined with paclitaxel and carboplatin. Recent literature on LELC is also reviewed and discussed.


## CASE PRESENTATION

### Chief complaints

A 61-year-old nonsmoker Chinese woman was admitted to our hospital on November 19, 2018. She had no symptoms, but her chest computed tomography (CT) during her annual healthy check-up showed a mass in the right lung lower lobe. Contrast-enhanced CT of the chest, brain, whole abdomen, and nasopharynx and a whole body bone scan were performed on November 20, 2018. The contrast-enhanced CT of the chest revealed a 2.8 cm × 2.9 cm mass in the right lower lobe, showing superficial lobules with unclear boundaries; the adjacent oblique fissure was pulled (Figure 1A and B). The contrast-enhanced CT of the brain, whole abdomen, and nasopharynx and whole body bone scan revealed no abnormality (Figure 1C and D). At the same time, the serum tumor markers carbohydrate antigen 125 (CA-125) and neural specificity enolase (NSE) were higher than normal (53.46 U/mL *vs* < 35 U/mL, and 23.96 ng/mL *vs* < 15 ng/mL, respectively); other tumor markers were negative. Physical examination showed an eastern cooperative oncology group performance status of 0. Surgical contraindications were excluded; video-assisted thoracoscopic surgery, right lower lobe lobotomy, and mediastinal lymph node dissection were performed under general anesthesia on November 22, 2018. Pathologic examination indicated a stage IIA (pT1aN1M0) disease based on the tumor size and interlobular lymph node involvement. Pathology of surgical specimen of the right lower lobe suggested LELC based on hematoxylin-eosin (HE) and immunohistochemical (IHC) staining; IHC staining demonstrated CK5/6 (+), P40 (+), PCK (+), TTF-1 (-), CK7 (-), NapsinA (-), CgA (-), Syn (-), Ki67 (+ 40%), EBER1/2-ISH (+), programmed death-1 (PD-L1) (+, 70%, SP142, ZSGB-BIO) (Figure 2A and B), and mismatch repair (MMR) proteins (integrity) (Figure 2C-F). No genetic mutations of KRAS, BRAF, EGFR, and ERBB2 or rearrangements in ALK, ROS1, and RET were seen after next-generation sequencing of the right lower lobe specimen was performed.

This patient did not undergo postoperative adjuvant chemotherapy. After being lost to follow-up for 1 year, she presented to the hospital in November 2019 with a lump of 4.3 cm × 5.2 cm in size on her right chest and pain. Contrast-enhanced CT indicated bone destruction of the right 4<sup>th</sup> rib with a soft tissue mass and cortical thickening of the adjacent 5<sup>th</sup> rib (Figure 3A and B). A whole-body bone scan suggested increased bone metabolism in the right 3<sup>rd</sup>-5<sup>th</sup> coastal axillary segments and possible bone metastasis (Figure 3C and D). At the same time, the serum tumor markers carbohydrate antigen 15-3 (CA15-3) and carbohydrate antigen 72-4 (CA72-4) were higher than normal (36 U/mL *vs* < 28 U/mL, and 14 ng/mL *vs* < 3.3 ng/mL, respectively). After multidisciplinary consultation and obtaining the patient's informed consent, she was treated with sintilimab (Tyvyt®, 200 mg ivgtt d0), paclitaxel (210 mg ivgtt d1), and carboplatin (550 mg ivgtt d1) every 3 wk. After two cycles, her lump on the right chest and pain resolved, and she did not experience any toxicity. Interestingly, the mass size was significantly reduced, and the patient achieved partial remission (PR) (according to RECIST 1.1). We checked contrast-enhanced CT again after six cycles of sintilimab combined with chemotherapy, which still revealed PR. Subsequently, the chemotherapy was stopped on March 17, 2020 and sintilimab maintenance treatment of 200 mg ivgtt (given every 3 wk) was initiated.

So far, the patient has finished six cycles of sintilimab combined with chemotherapy and ten cycles of sintilimab maintenance treatment (Figure 4). The values of CA15-3 and CA72-4 gradually returned to normal (10.4 U/mL *vs* < 28 U/mL, and 2.1 ng/mL *vs* < 3.3 ng/mL, respectively). The patient was in a continuous PR state. No obvious side effects were observed during the treatment process.



DOI: 10.12998/wjcc.v10.i31.11617 Copyright ©The Author(s) 2022.

**Figure 1** Imaging examinations before surgery. A and B: Contrast-enhanced computed tomography of the chest showing a 2.8 cm × 2.9 cm mass in the right lower lobe; C and D: Whole body bone scan showed no abnormality.

### **History of present illness**

A 61-year-old nonsmoker Chinese woman was admitted to our hospital on November 19, 2018, and contrast-enhanced CT of the chest revealed a 2.8 cm × 2.9 cm mass in the right lower lobe, showing superficial lobules with unclear boundaries. She underwent video-assisted thoracoscopic surgery, right lower lobe lobotomy, and mediastinal lymph node dissection.

### **History of past illness**

The patient had a history of hypertension and diabetes.

### **Personal and family history**

This patient was born in Chengdu and lived here. She denied having a family history of hereditary disease.

### **Physical examination**

The patient's temperature was 36.7 °C, and her blood pressure was 120/80 mmHg, heart rate was 70 bpm, and oxygen saturation in room air was 99%. Physical examination showed that the breath sound of both lungs was normal. No other positive signs were found in the physical examination.


### **Laboratory examinations**

Laboratory examinations revealed that the patient's CA15-3 was 36 U/mL, CA72-4 was 4 ng/mL, CA-125 was 53.46 U/mL, and NSE was 23.96 ng/mL.

### **Imaging examinations**

Contrast-enhanced CT indicated bone destruction of the right 4<sup>th</sup> rib with a soft tissue mass and cortical thickening of the adjacent 5<sup>th</sup> rib. A whole-body bone scan suggested increased bone metabolism in the right 3<sup>rd</sup>-5<sup>th</sup> coastal axillary segments and possible bone metastasis.





DOI: 10.12998/wjcc.v10.i31.11617 Copyright ©The Author(s) 2022.

**Figure 2 Immunohistochemistry.** A: Immunohistochemical (IHC) staining for programmed death-1 (PD-1) revealed positive neoplastic cells with a membranous pattern ( $\times 200$ , scale bar: 100  $\mu\text{m}$ ); B: IHC staining for PD-1 revealed negative neoplastic cells ( $\times 200$ , scale bar: 100  $\mu\text{m}$ ); C-F: IHC staining for mismatch repair (MMR) proteins revealed integrity of the MMR system ( $\times 200$ , scale bar: 100  $\mu\text{m}$ ).

## FINAL DIAGNOSIS

The patient was diagnosed as having primary pulmonary LELC.

## TREATMENT

The patient was treated with sintilimab (Tyvyt®, 200mg ivgtt d0), paclitaxel (210 mg ivgtt d1), and carboplatin (550 mg ivgtt d1) every 3 wk. Subsequently, the chemotherapy was stopped and sintilimab maintenance treatment of 200 mg ivgtt (given every 3 wk) was initiated.


## OUTCOME AND FOLLOW-UP

The patient was in a continuous PR state. No obvious side effects were observed during the treatment process. The values of CA15-3 and CA72-4 gradually returned to normal. She has been followed for 1 year.

## DISCUSSION

As far as we know, this is the first report of a patient with LELC who positively responded to sintilimab combined with chemotherapy. The patient achieved a partial response to our therapeutic regimen after two cycles. She completed six cycles of sintilimab combined with chemotherapy and ten cycles of sintilimab maintenance treatment. Her tumor burden significantly decreased, without obvious side effects. The timeline summarizing the main events of this case report is showed in Figure 5.

In order to investigate the molecular mechanism underlying the development of LELC, several studies have examined the mutation status of LELC oncogenic driver and tumor suppressor genes, such as EGFR, KRAS, BRAF, ALK, ROS1, and TP53. Yin *et al*[4] found EGFR mutation in five patients (5/175, 2.9%) and ALK alteration in three patients (3/140, 2.1%). The subtype might not be sensitive to conventional targeted therapy for NSCLC. Several studies revealed that PD-L1 expression is prevalent in pulmonary LELC. Chang *et al*[5] observed detectable PD-L1 expression in 75.8% (50/66) of pulmonary LELCs. There are high PD-L1 expression and infrequent driver mutations in LELCs compared with conventional NSCLCs. Wu *et al*[6] reported that among PD-L1-positive cases, 91.5% of primary



DOI: 10.12998/wjcc.v10.i31.11617 Copyright ©The Author(s) 2022.

**Figure 3 Imaging examinations after recurrence.** A and B: Contrast-enhanced computed tomography of the chest before therapy. The right 4<sup>th</sup> rib with a soft tissue mass measuring 41 mm × 35 mm, and cortical thickening of the adjacent 5<sup>th</sup> rib were detected; C and D: Whole-body bone scan revealed that bone metabolism was increased in the 3<sup>rd</sup>-5<sup>th</sup> costal axillary segments on the right side.


pulmonary LELC patients (54/59) had a tumor proportion score (TPS)  $\geq 5\%$ , while more than half of PLELC patients (36/59, 61.0 %) with a TPS  $\geq 50\%$  showed high levels of PD-L1 expression. Patients with positive PD-L1 expression had longer PFS and OS than those with negative PD-L1 expression. Xie *et al* [7] found that 69% (20/29) of patients had positive PD-L1 expression, whereas three patients benefited from a PD-1 inhibitor. In contrary, Paz-Ares *et al* [8] showed that the positive rate of PD-L1 in LELC patients was 74.3%, and overexpression of PD-L1 was associated with impaired DFS.

The treatment for PLELC mainly depends on tumor stage. Surgery is the mainstay of treatment in early-stage tumors, while a combination of chemotherapy and radiotherapy is used for advanced patients [3]. The most commonly used chemotherapy include cisplatin or carboplatin combined with 5-fluorouracil, paclitaxel, docetaxel, or gemcitabine [9]. High PD-L1 level and lack of canonical druggable driver mutations raise the potential of checkpoint immunotherapy for PLELC [10]. Our patient revealed the characteristics of the EBER1/2-ISH (+) and PD-L1 (+, 70%), and the lack of any canonical druggable driver mutations, which supported the feasibility of checkpoint immunotherapy.


The immune checkpoint PD-1 blockade is a recent addition to the treatment approach against different types of advanced cancers [8]. Two anti-PD-1 antibodies, nivolumab and pembrolizumab, have shown survival benefits over standard second-line chemotherapy in treating NSCLC [11]. Sintilimab, as a monotherapy, owns a safety profile in adults with relapsed or refractory classical Hodgkin's lymphoma participating in ORIENT-1 [9], which is similar to the safety profiles of nivolumab and pembrolizumab. Gao *et al* [10] discovered that neoadjuvant sintilimab was well tolerated by patients with NSCLC, and a 40.5% major pathologic response rate is encouraging. Thus far, immunotherapy has rarely been adopted for patients with LELC. At present, there are few reports on immunotherapy for LELC. Kim *et al* [12] first reported treating a patient with LELC using nivolumab who underwent chemotherapy; side effects, including tachypnea and cough, occurred 10 d after the first treatment. Moreover, Tang *et al* [11] reported a patient who was given 3 mg/kg nivolumab every 2 wk after first-line chemotherapy; after two cycles, the patient achieved PR. CT revealed that the disease had progressed 5 mo later in our patient, after which nedaplatin and paclitaxel were added to the nivolumab, resulting in a reduction of the tumor.

Our patient achieved PR after two cycles of sintilimab, paclitaxel, and carboplatin every 3 wk. By September 2020, the patient successfully completed six cycles of sintilimab combined with chemotherapy and ten cycles of sintilimab maintenance treatment in a continuous PR state without obvious side effects during the treatment process. To date, five patients with primary advanced LELCs





**Figure 4 Imaging examination.** A and B: Computed tomography (CT) after two cycles of immunotherapy combined with chemotherapy; C and D: CT scan performed after six cycles of immunotherapy combined with chemotherapy; E and F: CT revealed a continuous PR state after 6 mo of immunotherapy combined with chemotherapy.



**Figure 5 Timeline summarizing main events of this case report.**

have been reported, including the one reported in this study, who have received immunotherapy combined with chemotherapy or targeted therapy (Table 1). It is possible that immunotherapy combined with chemotherapy in advanced primary pulmonary LELC may be more effective than immunotherapy or chemotherapy alone. Thus, immunotherapy combined with chemotherapy should be considered as the first line of treatment for patients with advanced primary pulmonary LELC.

Tang *et al* [11] found that NSE is an effective biomarker for immunotherapy in LELC. In our patients, we found that after effective treatment, CA15-3 and CA72-4 returned to normal. NSE remained normal throughout the relapse. Thus, CA15-3 and CA72-4 are efficient biomarkers for detecting LELC.

Although immune checkpoint inhibitor-based therapy has shown great potential in solid tumors, the shortage of validated biomarkers is still an important issue since only a part of patients could benefit from immunotherapy. In addition, some potential biomarkers should be further explored, such as PD-L1 level, tumor mutational burden, microsatellite instability status, and gut microbiota. Furthermore, the efficacy of immunotherapy in clinical trials may differ in terms of drugs, patients, designs, study phases, and so on.

## CONCLUSION

Our patient achieved PR after immunotherapy combined with chemotherapy, which suggested that the

**Table 1 Case reports and retrospective series of lymphoepithelioma-like carcinoma management with programmed cell death protein 1/programmed death-1 inhibition in combination with chemotherapy or targeted therapy**

Case	Reference	EBV	PD-L1 expression	Previous treatment history	Treatment	Onset time	Best response	PFS (M)
1	Zhou <i>et al</i> [13]	NA	60%	1 <sup>st</sup> -line chemotherapy	2 <sup>nd</sup> -line nivolumab, 3 <sup>rd</sup> -line nivolumab + anlotinib + gemcitabine	NA	SD	2, 4
2	Zhou <i>et al</i> [13]	NA	30%	NA	Nivolumab + gemcitabine	NA	SD	NA
3	Xie <i>et al</i> [7]	NA	15%	2 <sup>nd</sup> -line chemotherapy	Camrelizumab + apatinib	NA	SD	7 +
4	Tang <i>et al</i> [11]	NA	10%	1 <sup>st</sup> -line chemotherapy	2 <sup>nd</sup> -line nivolumab, 3 <sup>rd</sup> -line Nivolumab + nedaplatin + paclitaxel	Two cycles Four cycles	PR SD	5 NA
5	Zeng <i>et al</i> (our case)	+	70%	Surgery	Sintilimab + paclitaxel + carboplatin	Two cycles	PR	NA

EBV: Epstein-Barr virus; PD-L1: Programmed death-1; PFS: Progression-free survival; OS: Overall survival; PR: Partial response; SD: Stable disease; NA: Not available.

combined treatment regimen may be more effective than immunotherapy or chemotherapy alone. Meanwhile, both CA15-3 and CA72-4 are biomarkers for evaluating therapeutic effects for LELC.

## ACKNOWLEDGEMENTS

The authors wish to thank the patient for allowing the publication of her case details.

## FOOTNOTES

**Author contributions:** Zeng SY contributed to protocol/project development, data collection or management, data analysis, and manuscript writing/editing; Yuan J contributed to data collection and manuscript editing; Lv M contributed to protocol/project development and manuscript editing.

**Informed consent statement:** Informed written consent was obtained from the patient for the publication of this report and any accompanying images.

**Conflict-of-interest statement:** The authors declare that they have no conflict of interest to disclose.

**CARE Checklist (2016) statement:** The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** China

**ORCID number:** Shu-Ying Zeng 0000-0002-8298-9320.

**S-Editor:** Chen YL

**L-Editor:** Wang TQ

**P-Editor:** Chen YL

## REFERENCES

- 1 **Bégin LR**, Eskandari J, Joncas J, Panasci L. Epstein-Barr virus related lymphoepithelioma-like carcinoma of lung. *J Surg Oncol* 1987; **36**: 280-283 [PMID: [2826922](#) DOI: [10.1002/jso.2930360413](#)]
- 2 **Hong S**, Liu D, Luo S, Fang W, Zhan J, Fu S, Zhang Y, Wu X, Zhou H, Chen X, Chen G, Zhang Z, Zheng Q, Li X, Chen J, Liu X, Lei M, Ye C, Wang J, Yang H, Xu X, Zhu S, Yang Y, Zhao Y, Zhou N, Zhao H, Huang Y, Zhang L, Wu K. The genomic landscape of Epstein-Barr virus-associated pulmonary lymphoepithelioma-like carcinoma. *Nat Commun* 2019; **10**: 3108 [PMID: [31311932](#) DOI: [10.1038/s41467-019-10902-w](#)]
- 3 **Qin Y**, Gao G, Xie X, Zhu Z, Guan W, Lin X, Xie Z, Ming O, Chen R, Zhong N, Li S, Zhou C. Clinical Features and Prognosis of Pulmonary Lymphoepithelioma-like Carcinoma: Summary of Eighty-five Cases. *Clin Lung Cancer* 2019; **20**: e329-e337 [PMID: [30683631](#) DOI: [10.1016/j.clcc.2018.12.014](#)]
- 4 **Yin K**, Feng HB, Li LL, Chen Y, Xie Z, Lv ZY, Guo WB, Lu DX, Yang XN, Yan WQ, Wu YL, Zhang XC. Low frequency of mutation of epidermal growth factor receptor (EGFR) and arrangement of anaplastic lymphoma kinase (ALK) in primary pulmonary lymphoepithelioma-like carcinoma. *Thorac Cancer* 2020; **11**: 346-352 [PMID: [31794146](#) DOI: [10.1111/1759-7714.13271](#)]
- 5 **Chang YL**, Yang CY, Lin MW, Wu CT, Yang PC. PD-L1 is highly expressed in lung lymphoepithelioma-like carcinoma: A potential rationale for immunotherapy. *Lung Cancer* 2015; **88**: 254-259 [PMID: [25862146](#) DOI: [10.1016/j.lungcan.2015.03.017](#)]
- 6 **Wu Q**, Wang W, Zhou P, Fu Y, Zhang Y, Shao YW, Jiang L. Primary pulmonary lymphoepithelioma-like carcinoma is characterized by high PD-L1 expression, but low tumor mutation burden. *Pathol Res Pract* 2020; **216**: 153043 [PMID: [32703503](#) DOI: [10.1016/j.prp.2020.153043](#)]
- 7 **Xie Z**, Liu L, Lin X, Xie X, Gu Y, Liu M, Zhang J, Ouyang M, Lizaso A, Zhang H, Feng W, Li B, Han-Zhang H, Chen S, Li S, Zhong N, Liu H, Zhou C, Qin Y. A multicenter analysis of genomic profiles and PD-L1 expression of primary lymphoepithelioma-like carcinoma of the lung. *Mod Pathol* 2020; **33**: 626-638 [PMID: [31659278](#) DOI: [10.1038/s41379-019-0391-9](#)]
- 8 **Paz-Ares L**, Luft A, Vicente D, Tafreshi A, Gümüş M, Mazières J, Hermes B, Çay Şenler F, Csösz T, Fülöp A, Rodríguez-Cid J, Wilson J, Sugawara S, Kato T, Lee KH, Cheng Y, Novello S, Halmos B, Li X, Lubiniecki GM, Piperdi B, Kowalski DM; KEYNOTE-407 Investigators. Pembrolizumab plus Chemotherapy for Squamous Non-Small-Cell Lung Cancer. *N Engl J Med* 2018; **379**: 2040-2051 [PMID: [30280635](#) DOI: [10.1056/NEJMoa1810865](#)]
- 9 **Shi Y**, Su H, Song Y, Jiang W, Sun X, Qian W, Zhang W, Gao Y, Jin Z, Zhou J, Jin C, Zou L, Qiu L, Li W, Yang J, Hou M, Zeng S, Zhang Q, Hu J, Zhou H, Xiong Y, Liu P. Safety and activity of sintilimab in patients with relapsed or refractory classical Hodgkin lymphoma (ORIENT-1): a multicentre, single-arm, phase 2 trial. *Lancet Haematol* 2019; **6**: e12-e19 [PMID: [30612710](#) DOI: [10.1016/S2352-3026\(18\)30192-3](#)]
- 10 **Gao S**, Li N, Gao S, Xue Q, Ying J, Wang S, Tao X, Zhao J, Mao Y, Wang B, Shao K, Lei W, Wang D, Lv F, Zhao L, Zhang F, Zhao Z, Su K, Tan F, Gao Y, Sun N, Wu D, Yu Y, Ling Y, Wang Z, Duan C, Tang W, Zhang L, He S, Wu N, Wang J, He J. Neoadjuvant PD-1 inhibitor (Sintilimab) in NSCLC. *J Thorac Oncol* 2020; **15**: 816-826 [PMID: [32036071](#) DOI: [10.1016/j.jtho.2020.01.017](#)]
- 11 **Tang Z**, Fang R, Tong G, Liu P, Ou Z, Tang Y. Overcoming resistance to anti-PD-1 immunotherapy in lymphoepithelioma-like carcinoma: A case report and review of the literature. *Lung Cancer* 2020; **146**: 335-340 [PMID: [32623076](#) DOI: [10.1016/j.lungcan.2020.06.027](#)]
- 12 **Kim C**, Rajan A, DeBrito PA, Giaccone G. Metastatic lymphoepithelioma-like carcinoma of the lung treated with nivolumab: a case report and focused review of literature. *Transl Lung Cancer Res* 2016; **5**: 720-726 [PMID: [28149767](#) DOI: [10.21037/tlcr.2016.11.06](#)]
- 13 **Zhou N**, Lin Y, Peng X, Wang Y. Thorough survey and analysis of pulmonary lymphoepithelioma-like carcinoma in Macau and multimodality treatment for advanced disease. *Lung Cancer* 2019; **138**: 116-123 [PMID: [31683094](#) DOI: [10.1016/j.lungcan.2019.10.004](#)]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

