

# World Journal of *Gastrointestinal Surgery*

*World J Gastrointest Surg* 2022 November 27; 14(11): 1179-1319



**REVIEW**

- 1179 Diagnosis, severity stratification and management of adult acute pancreatitis—current evidence and controversies  
*Chan KS, Shelat VG*

**MINIREVIEWS**

- 1198 Development and future perspectives of natural orifice specimen extraction surgery for gastric cancer  
*Zhang ZC, Luo QF, Wang WS, Chen JH, Wang CY, Ma D*

**ORIGINAL ARTICLE****Retrospective Cohort Study**

- 1204 Clinical value of extended lymphadenectomy in radical surgery for pancreatic head carcinoma at different T stages  
*Lyu SC, Wang HX, Liu ZP, Wang J, Huang JC, He Q, Lang R*
- 1219 Comparison of clinicopathological characteristics between resected ampullary carcinoma and carcinoma of the second portion of the duodenum  
*Nishio K, Kimura K, Murata A, Ohira G, Shinkawa H, Kodai S, Amano R, Tanaka S, Shimizu S, Takemura S, Kanazawa A, Kubo S, Ishizawa T*

**Retrospective Study**

- 1230 Metastatic lymph nodes and prognosis assessed by the number of retrieved lymph nodes in gastric cancer  
*Wang H, Yin X, Lou SH, Fang TY, Han BL, Gao JL, Wang YF, Zhang DX, Wang XB, Lu ZF, Wu JP, Zhang JQ, Wang YM, Zhang Y, Xue YW*
- 1250 Comprehensive abdominal composition evaluation of rectal cancer patients with anastomotic leakage compared with body mass index-matched controls  
*Shao SL, Li YK, Qin JC, Liu L*
- 1260 Recombinant human thrombopoietin treatment in patients with chronic liver disease-related thrombocytopenia undergoing invasive procedures: A retrospective study  
*Ding JN, Feng TT, Sun W, Cai XY, Zhang Y, Zhao WF*

**Observational Study**

- 1272 Assessment of tumor markers CA 19-9, CEA, CA 125, and CA 242 for the early diagnosis and prognosis prediction of gallbladder cancer  
*Sinha SR, Prakash P, Singh RK, Sinha DK*
- 1285 Disturbed passage of jejunal limb near esophageal hiatus after overlapped esophagojejunostomy following laparoscopic total gastrectomy  
*Noshiro H, Okuyama K, Yoda Y*

- 1297 Development of a warning score for early detection of colorectal anastomotic leakage: Hype or hope?

*Rama NJG, Lourenço Ó, Motta Lima PC, Guarino MPS, Parente D, Castro R, Bento A, Rocha A, Castro-Poças F, Pimentel J*

**CASE REPORT**

- 1310 Preoperative blood circulation modification prior to pancreaticoduodenectomy in patients with celiac trunk occlusion: Two case reports

*Colella M, Mishima K, Wakabayashi T, Fujiyama Y, Al-Omari MA, Wakabayashi G*

**ABOUT COVER**

Editorial Board Member of *World Journal of Gastrointestinal Surgery*, Ketan Vagholkar, MBBS, MS, DNB, FRCS (Gen Surg), FACS, Full Professor, Department of Surgery, D.Y.Patil University School of Medicine, Navi Mumbai 400706, Maharashtra, India. ketan.vagholkar@dypatil.edu

**AIMS AND SCOPE**

The primary aim of *World Journal of Gastrointestinal Surgery* (*WJGS, World J Gastrointest Surg*) is to provide scholars and readers from various fields of gastrointestinal surgery with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

*WJGS* mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal surgery and covering a wide range of topics including biliary tract surgical procedures, biliopancreatic diversion, colectomy, esophagectomy, esophagostomy, pancreas transplantation, and pancreatectomy, etc.

**INDEXING/ABSTRACTING**

The *WJGS* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for *WJGS* as 2.505; IF without journal self cites: 2.473; 5-year IF: 3.099; Journal Citation Indicator: 0.49; Ranking: 104 among 211 journals in surgery; Quartile category: Q2; Ranking: 81 among 93 journals in gastroenterology and hepatology; and Quartile category: Q4.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Rui-Rui Wu, Production Department Director: Xiang Li; Editorial Office Director: Jia-Ru Fan.

**NAME OF JOURNAL**

*World Journal of Gastrointestinal Surgery*

**ISSN**

ISSN 1948-9366 (online)

**LAUNCH DATE**

November 30, 2009

**FREQUENCY**

Monthly

**EDITORS-IN-CHIEF**

Peter Schemmer

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/1948-9366/editorialboard.htm>

**PUBLICATION DATE**

November 27, 2022

**COPYRIGHT**

© 2022 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

## Retrospective Study

# Comprehensive abdominal composition evaluation of rectal cancer patients with anastomotic leakage compared with body mass index-matched controls

Sheng-Li Shao, Yang-Kun Li, Ji-Chao Qin, Lu Liu

**Specialty type:** Gastroenterology and hepatology**Provenance and peer review:** Unsolicited article; Externally peer reviewed.**Peer-review model:** Single blind**Peer-review report's scientific quality classification**Grade A (Excellent): 0  
Grade B (Very good): B  
Grade C (Good): C  
Grade D (Fair): 0  
Grade E (Poor): 0**P-Reviewer:** Rama N, Portugal;  
Urlin VM, Romania**Received:** August 2, 2022**Peer-review started:** August 2, 2022**First decision:** September 4, 2022**Revised:** September 27, 2022**Accepted:** November 4, 2022**Article in press:** November 4, 2022**Published online:** November 27, 2022**Sheng-Li Shao, Yang-Kun Li, Ji-Chao Qin, Lu Liu**, Department of Surgery, Huazhong University of Science and Technology, Wuhan 430030, Hubei Province, China**Corresponding author:** Lu Liu, MD, Associate Chief Physician, Department of Surgery, Huazhong University of Science and Technology, No. 1095 Jiefang Avenue, Wuhan 430030, Hubei Province, China. [halesan@163.com](mailto:halesan@163.com)

## Abstract

### BACKGROUND

Anastomotic leakage (AL) is a fatal complication in patients with rectal cancer after undergoing anterior resection. However, the role of abdominal composition in the development of AL has not been studied.

### AIM

To investigate the relationship between abdominal composition and AL in rectal cancer patients after undergoing anterior resection.

### METHODS

A retrospective case-matched cohort study was conducted. Complete data for 78 patients with AL were acquired and this cohort was defined as the AL group. The controls were matched for the same sex and body mass index ( $\pm 1 \text{ kg/m}^2$ ). Parameters related to abdominal composition including visceral fat area (VFA), subcutaneous fat area (SFA), subcutaneous fat thickness (SFT), skeletal muscle area (SMA), skeletal muscle index (SMI), abdominal circumference (AC), anterior to posterior diameter of abdominal cavity (APD), and transverse diameter of abdominal cavity (TD) were evaluated based on computed tomography (CT) images using the following Hounsfield Unit (HU) thresholds: SFA: -190 to -30, SMA: -29 to 150, and VFA: -150 to -20. The significance of abdominal composition-related parameters was quantified using feature importance analysis; an artificial intelligence method was used to evaluate the contribution of each included variable.

### RESULTS

Two thousand two hundred and thirty-eight rectal cancer patients who underwent anterior resection from 2010 to 2020 in a large academic hospital were investigated. Finally, 156 cases were enrolled in the study. Patients in the AL

group showed longer operative time ( $225.03 \pm 55.29$  vs  $207.17 \pm 40.80$ ,  $P = 0.023$ ), lower levels of preoperative hemoglobin ( $123.32 \pm 21.17$  vs  $132.60 \pm 6.31$ ,  $P = 0.003$ ) and albumin ( $38.34 \pm 4.01$  vs  $40.52 \pm 3.97$ ,  $P = 0.001$ ), larger tumor size ( $4.07 \pm 1.36$  vs  $2.76 \pm 1.28$ ,  $P < 0.001$ ), and later cancer stage ( $P < 0.001$ ) compared to the controls. Patients who developed AL exhibited a larger VFA ( $125.68 \pm 73.59$  vs  $97.03 \pm 57.66$ ,  $P = 0.008$ ) and a smaller APD ( $77.30 \pm 23.23$  vs  $92.09 \pm 26.40$ ,  $P < 0.001$ ) and TD ( $22.90 \pm 2.23$  vs  $24.21 \pm 2.90$ ,  $P = 0.002$ ) compared to their matched controls. Feature importance analysis revealed that TD, APD, and VFA were the three most important abdominal composition-related features.

### CONCLUSION

AL patients have a higher visceral fat content and a narrower abdominal structure compared to matched controls.

**Key Words:** Anastomotic leakage; Abdominal composition; Rectal cancer; Body mass index-matched; Anterior to posterior diameter; Transverse diameter

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** We investigated the association between abdominal composition and anastomotic leakage in rectal cancer patients who underwent anterior resection in a large academic hospital from 2010 to 2020. The data revealed that patients who developed anastomotic leakage had a higher visceral fat content and a narrower abdominal structure, despite body mass index matching.

**Citation:** Shao SL, Li YK, Qin JC, Liu L. Comprehensive abdominal composition evaluation of rectal cancer patients with anastomotic leakage compared with body mass index-matched controls. *World J Gastrointest Surg* 2022; 14(11): 1250-1259

**URL:** <https://www.wjgnet.com/1948-9366/full/v14/i11/1250.htm>

**DOI:** <https://dx.doi.org/10.4240/wjgs.v14.i11.1250>

## INTRODUCTION

Compelling evidence demonstrates that total mesorectal resection (TME) successfully reduces the local recurrence rate of rectal cancer and is the gold standard for managing mid- and low-lying rectal cancer [1-3]. However, the morbidity of anastomotic leakage (AL), a worrisome complication of TME, is on the rise [4]. Once AL develops, it often requires reintervention and can lead to perioperative death and adverse oncology outcomes [5-7]. Early identification of patients at high risk of AL is critical to AL prevention and reduction of the reoperation rate, and will guide intraoperative decisions (for instance on whether to choose a diverting ileostomy or not) and improve perioperative management.

Numerous studies have explored the risk factors associated with AL in rectal cancer patients who underwent anterior resection [8,9]. However, there is no effective approach for predicting AL, implying that potential predictors should be identified. Recent studies show that some abdominal composition related factors are key contributors to AL in patients with colorectal cancer after undergoing surgery [10]. Theoretically, a less visceral fat content and a bigger abdominal volume are more favorable for surgeons to perform anterior resection procedure and thus leads to less technical difficulty, shorter operation time, and lower probability of AL [11]. Computed tomography (CT) images have been employed to assess the possible effects of abdominal composition related parameters, including visceral fat area (VFA) and skeletal muscle index (SMI), on patient surgical outcome [10,12-15]. Large VFA, for instance, is potentially effective in predicting AL in patients with colorectal cancer who received anterior resection despite reports to the contrary [9]. Additionally, SMI, measured by a CT scan of the lower margin of the third lumbar spine, is a reliable indicator of the systemic nutritional status and is associated with perioperative complications [16]. Additional indicators, including abdominal circumference (AC), anterior to posterior diameter of abdominal cavity (APD), and transverse diameter of abdominal cavity (TD), are suggested to exert potential effects on perioperative complications but their roles in AL is unknown.

Considering the impact of abdominal composition on the surgeons and patients, it was hypothesized that the abdominal composition of rectal cancer patients who developed AL after anterior resection may be different from that of individuals with similar body mass index (BMI) who did not develop AL. Here, we compared the abdominal composition between AL patients and sex- and BMI-matched controls.

## MATERIALS AND METHODS

### Patients

A total of 2238 medical records of rectal cancer patients who underwent anterior resection at our center from January 1, 2010 to January 1, 2020 were reviewed. Of note, 173 patients were excluded due to non-primary rectal adenocarcinoma ( $n = 32$ ) and missing clinical data ( $n = 141$ ). All patients underwent a 90-d follow-up. Of the 2065 subjects, 107 (5.18%) developed clinical AL (*i.e.*, grades B and C). Among the AL patients, 29 were excluded for missing CT images, and the remaining 78 were included in the final analysis and defined as the AL group. The control group was matched 1:1 for the same sex and BMI ( $\pm 1$  kg/m<sup>2</sup>) from patients who did not develop AL. A flowchart of this study is shown in [Figure 1](#).

### Definition and variables

In this study, rectal cancer was defined as a tumor located between the dentate line and sacral promontory. AL refers to clinical AL, including grade B and grade C, defined as disruption and defect in intestinal wall integrity at the anastomosis site, making the internal and external compartments communicate with each other[17]. AL diagnosis is contingent on the fecal fluid from pelvic draining or water-soluble contrast agent enema and extra-rectal imaging. Alternatively, when AL was suspected, perianastomotic abscess or effusion detected by CT was examined to diagnose AL. Because water-soluble contrast agent enema is not performed routinely at our center, AL of grade A was not included. The clinical variables gender, age, height, weight, BMI, ASA score, previous abdominal history, hypertension, diabetes, cigarette smoking, alcohol use, tumorous obstruction, preoperative cleansing enema, preoperative antibiotic use, distance between tumor and anal margin, neoadjuvant, preoperative hemoglobin, preoperative albumin, type of operation, tumor size, clinical tumor stage, operation time, number of linear stapler firings, indwelling pelvic drainage tube, indwelling trans-anal tube, and stoma were also considered. Abdominal composition-related parameters assessed included BMI, AC, subcutaneous fat area (SFA), subcutaneous fat thickness (SFT), skeletal muscle area (SMA), SMI, VFA, APD, and TD.

### Assessment of abdominal composition associated parameters

Data of BMI and AC were acquired from medical records, whereas other indicators were examined at the lower margin of the third lumbar (L3) plane of the unenhanced CT image using Slice-O-Matic software (version 5.0; Tomovision, Montreal, Canada). CT images were saved in DICOM (Medical Digital Imaging and Communication) format and retrieved from the institutional database. SFA, SMA, and VFA were measured by setting Hounsfield Unit (HU) thresholds (SFA: -190 to -30, SMA: -29 to 150 and VFA: -150 to -20)[18]. SFT refers to the vertical distance from the linear alba to the skin. SMI was calculated as SMA/height<sup>2</sup> (cm<sup>2</sup>/m<sup>2</sup>)[19,20]. APD refers to the vertical distance from the linear alba to the anterior edge of the L3 spine. TD refers to the transverse diameter of the abdominal cavity through the anterior edge of the L3 spine.

### Statistical analysis

Continuous variables are presented as the mean and standard deviation (SD), whereas categorical variables are presented as numerical values (percentages). Student's *t*-test and chi-square test were used to compare continuous variables and categorical variables, respectively. A *P* value of < 0.05 denoted statistical significance. All statistical analyses were performed using IBM SPSS 24.0 (SPSS for Windows, IBM Corporation, Armonk, NY, United States).

### Feature importance analysis

Feature importance analysis is an artificial intelligence method used for examining the importance of each included feature. This approach is based on some ensemble learning algorithms, such as random forest and XGboost. In this study, we used the random forest analysis to calculate the importance of each abdominal composition related parameter. Random forest is an ensemble classifier based on a combination of multiple decision trees which are generated through sampling from the original data set and the final predictions are voted by integrating all the trees. Mean decrease accuracy was calculated by randomly permuting a variable to reassess the predictions. If a variable is important, the mean decrease accuracy will show a large change. Therefore, the random forest algorithm could compute the importance of each included variable. This procedure was conducted using Scikit-learn package (version 0.24.1) in Python 3.8.5.

## RESULTS

### Demographic and clinical characteristics


A total of 156 patients were included in the final analysis. [Table 1](#) shows the comparison of the clinical characteristics between the AL group and the control group. Compared to the controls, the patients in

**Table 1 Comparison of demographic and clinicopathologic characteristics in patients with anastomotic leakage and controls, *n* (%)**

Variable	Controls ( <i>n</i> = 78)	AL patients ( <i>n</i> = 78)	<i>P</i> value
Male sex	57 (71.3)	57 (71.3)	1.000
Age, mean (SD), yr	58.23 (9.46)	56.82 (10.54)	0.380
Height, mean (SD), cm	166.23 (7.92)	166.87 (7.34)	0.601
Weight, mean (SD), kg	63.41 (11.62)	65.40 (11.39)	0.282
Operative time, mean (SD), min	207.17 (40.80)	225.03 (55.29)	0.023
Laparoscopic surgery	77 (98.7)	76 (97.4)	1.000
Location of tumor, mean (SD), cm	7.86 (3.39)	8.22 (3.59)	0.507
Intraperitoneal chemotherapy	50 (64.1)	54 (69.2)	0.497
Cleansing enema	57 (73.1)	60 (76.9)	0.579
Indwelling trans-anal tube	73 (93.6)	68 (87.2)	0.174
Indwelling drainage tube	72 (92.3)	74 (94.9)	0.746
Tumorous obstruction	1 (1.3)	6 (7.7)	0.053
Cigarette smoking	24 (30.8)	35 (44.9)	0.098
Alcohol use	14 (17.9)	21 (26.9)	0.249
Hypertension	20 (25.6)	19 (24.4)	1.000
Diabetes	10 (12.8)	11 (14.1)	1.000
Previous abdominal surgery	11 (14.1)	5 (6.4)	0.186
Preoperative antibiotics	75 (76.2)	72 (92.3)	0.303
Hemoglobin, mean (SD), g/L	132.60 (16.31)	123.32 (21.17)	0.003
Albumin, mean (SD), g/L	40.52 (3.97)	38.34 (4.01)	0.001
Neoadjuvant therapy	1 (1.3)	3 (3.8)	0.620
Tumor size, mean (SD), cm	2.76 (1.28)	4.07 (1.36)	< 0.001
ASA score			0.049
1	17 (21.86)	9 (11.5)	
2	56 (71.8)	56 (71.8)	
3	5 (6.4)	13 (16.7)	
Stage			< 0.001
1	67 (85.9)	19 (24.4)	
2	5 (6.4)	33 (42.3)	
3	6 (7.7)	26 (33.3)	
Number of linear stapler firings			0.393
1	38 (48.7)	37 (47.4)	
2	39 (50.0)	37 (47.4)	
3	1 (1.3)	4 (5.1)	
Stoma	20 (25.6)	18 (23.1)	0.852

SD: Standard deviation; ASA: American Society of Anesthesiologists; AL: Anastomotic leakage.

the AL group had longer operative time ( $225.03 \pm 55.29$  vs  $207.17 \pm 40.80$ ,  $P = 0.023$ ). Patients in the AL group exhibited lower levels of preoperative hemoglobin ( $123.32$  vs  $132.60$ ,  $P = 0.003$ ) and albumin ( $38.34$  vs  $40.52$ ,  $P = 0.001$ ), larger tumor size ( $4.07$  vs  $2.76$ ,  $P < 0.001$ ), and later cancer stage ( $P < 0.001$ ) compared to the controls. The ASA score had a marginal effect ( $P = 0.049$ ). No statistical difference was found between the AL group and the control group for other features.



DOI: 10.4240/wjgs.v14.i11.1250 Copyright ©The Author(s) 2022.

Figure 1 Flow chart of patient inclusion and exclusion.

### Comparison of abdominal composition related parameters

Table 2 shows the difference in abdominal composition related parameters between the AL group and the control group. Patients in the AL group had a larger VFA (125.68 *vs* 97.03,  $P = 0.008$ ), a smaller APD (77.30 *vs* 92.09,  $P < 0.001$ ), and a smaller TD (22.90 *vs* 24.21,  $P = 0.002$ ) compared to those in the control group. These results are intriguing and suggest a potential contribution of a narrower abdominal cavity to AL development. Differences in other indicators were not statistically significant. A radar plot demonstrated the comparison of these indicators between the AL group and the control group (Figure 2).

### Feature importance analysis

Although determination of statistical significance of abdominal composition-related indicators can be used to prove correlations, it is not sufficient. Feature importance analysis was conducted to quantify the contribution of each abdominal composition related indicator in AL development. Results demonstrated that TD, APD, and VFA were the three most important features (Figure 3). Additionally, we performed univariate and multivariate logistic regression analyses to investigate whether the VFA, APD, and TD were independent risk factors for AL. The data indicated that the VFA, APD, and TD were independent risk factors ( $P < 0.05$ ) (Supplementary Table 1).

## DISCUSSION


The mechanism underlying AL occurrence involves several factors. The present work compared the clinical characteristics and abdominal composition in rectal cancer patients who received anterior resection and developed AL to controls who were matched for sex and BMI. This study was conducted in a large academic hospital in which more than 4000 gastrointestinal operations were performed annually. Analysis revealed a 5.18% incidence of clinical AL, which concur with previous reports[21-23]. In this study cohort, when comparison was conducted in clinical characteristics, lower levels of preoperative hemoglobin and albumin, longer operative time, larger tumor size, and later cancer stage were associated with AL. In addition, when comparing abdominal composition related parameters, it is interesting to find that a higher visceral fat content and a narrower abdominal structure were associated with AL. This work provides evidence that the occurrence of AL is not only associated with patient related factors, but also with the underlying factors that may affect surgical technique.

Related studies have demonstrated that BMI, an easily available and most commonly used index of obesity, is a risk factor for AL in rectal cancer patients who received anterior resection. However, other studies have reported contrary reports[24,25]. Considering that BMI cannot distinguish between the content and distribution of fat and skeletal muscle, it is imperative to explore whether fat and skeletal muscle content or distribution potentially impacts the development of AL. Verduin *et al*[9] investigated the role of VFA on AL in 2370 colon cancer patients and the results implicated VFA as an independent risk factor for AL in the elective colon resection patients (odds ratio = 1.026,  $P = 0.035$ ). Elsewhere, a study employed CT images to quantify the fat distribution and proposed the association of high adipose

**Table 2 Comparison of abdominal parameters in patients with anastomotic leakage and controls**

Variables	Controls (n = 78)	AL patients (n = 78)	P value
BMI (SD), kg/m <sup>2</sup>	23.05 (3.05)	23.17 (2.88)	0.797
AC, mean (SD), cm	87.00 (10.94)	89.71 (14.20)	0.120
SFA, mean (SD), cm <sup>2</sup>	108.72 (54.12)	113.72 (55.87)	0.571
SFT, mean (SD), mm	18.68 (8.20)	18.03 (7.31)	0.601
SMA, mean (SD), cm <sup>2</sup>	127.89 (29.57)	132.06 (33.40)	0.410
SMI, mean (SD), cm <sup>2</sup> /m <sup>2</sup>	46.00 (8.81)	47.10 (10.57)	0.482
VFA, mean (SD), cm <sup>2</sup>	97.03 (57.66)	125.68 (73.59)	0.008
APD, mean (SD), mm	92.09 (26.40)	77.30 (23.23)	< 0.001
TD, mean (SD), cm	24.21 (2.90)	22.90 (2.23)	0.002

BMI: Body mass index; SD: Standard deviation; AC: Abdominal circumference; SFA: Subcutaneous fat area; SFT: Subcutaneous fat thickness; SMA: Skeletal muscle area; SMI: Skeletal muscle index (SMA/height<sup>2</sup>); VFA: Visceral fat area; APD: Anterior to posterior diameter of abdominal cavity; TD: Transverse diameter of abdominal cavity; AL: Anastomotic leakage.




DOI: 10.4240/wjgs.v14.i11.1250 Copyright ©The Author(s) 2022.

**Figure 2 Radar plot for comparison of abdominal composition related parameters between the anastomotic leakage group and the control group.** AC: Abdominal circumference; SFA: Subcutaneous fat area; SFT: Subcutaneous fat thickness; SMA: Skeletal muscle area; SMI: Skeletal muscle index (SMA/height<sup>2</sup>); VFA: Visceral fat area; APD: Anterior to posterior diameter of abdominal cavity; TD: Transverse diameter of abdominal cavity; AL: Anastomotic leakage.

tissue with higher risk AL in rectal cancer patients[26]. However, whether VFA and other abdominal composition parameters potentially influence the occurrence of AL in patients with a similar BMI remains to be further evaluated. In addition, owing to the narrow pelvic structure, the male sex is widely accepted as an independent risk factor for AL in rectal cancer patients who received anterior resection, and some evidence has demonstrated the role of pelvic related parameters on AL[27]. Theoretically, a narrow pelvic structure is associated with the increased difficulty of the operation and prolonged operation time. All these features may increase the risk of AL. However, whether a narrow abdominal structure plays a similar role in AL occurrence is not known.

By comparing the differences in abdominal composition between AL and non-AL patients through sex and BMI matching, we found a higher VFA (125.68 *vs* 97.03,  $P = 0.008$ ) and smaller narrow abdominal cavity structure (APD, 77.30 *vs* 92.09,  $P < 0.001$ ; TD, 22.90 *vs* 24.21,  $P = 0.002$ ) in AL patients than in the controls. The differences in skeletal muscle-related parameters, including SMA and SMI,



**Figure 3 Importance of each feature in the development of anastomotic leakage.** AC: Abdominal circumference; SFA: Subcutaneous fat area; SFT: Subcutaneous fat thickness; SMA: Skeletal muscle area; SMI: Skeletal muscle index (SMA/height<sup>2</sup>); VFA: Visceral fat area; APD: Anterior to posterior diameter of abdominal cavity; TD: Transverse diameter of abdominal cavity; AL: Anastomotic leakage.

were not significant, which may be ascribed to the unbalanced matching of other variables between the AL patients and controls, because various variables are associated with muscle content and density. This study provides support to the hypothesis that even with a similar BMI, AL patients are characterized by a higher VFA and a narrower abdominal structure.

This study has several limitations. First, as a single-center case-matched study, selection bias cannot be completely ignored. Second, although standard and strict screening and matching criteria were employed, the large initial sample size and the small sample size for analysis may imply that the research results need to be further validated on a larger cohort. Third, some variables impacting abdominal composition were not collected, including whether subjects are athletes, metabolic syndrome, *etc.* Lastly, this study was performed based on abdominal CT images, and as such, some indicators such as muscle density and intermuscular fat could not be evaluated in detail. Given the retrospective nature of this study and the small sample size, future longitudinal investigations with large samples are advocated to provide reliable data to determine causality for the correlation of abdominal components and AL.

## CONCLUSION

The present analysis demonstrates the difference in abdominal components between AL patients and controls matched for sex and BMI. The contribution of each indicator to the development of AL was demonstrated. Intriguingly, in addition to the differences in VFA, the negative effects of APD and TD on AL were observed. This study adds considerable value to the field of AL preoperative risk assessment in rectal cancer patients. VFA, APD, and TD are potential indicators for predicting the risk of AL and can guide surgical decision-making (for example, performing a temporary ileostomy for high-risk patients).

## ARTICLE HIGHLIGHTS

### Research background

Compelling evidence demonstrates the relationship of abdominal composition and postoperative complications. Anastomotic leakage (AL) is a fatal complication in patients with rectal cancer who have

received anterior resection. However, the roles of abdominal composition on AL have not been studied.

### **Research motivation**

To study the characteristics of abdominal components in patients who received rectal cancer surgery and developed AL.

### **Research objectives**

To add risk factors for AL prediction in rectal cancer patients undergoing anterior resection for guiding surgical decision-making, *e.g.*, performing a temporary ileostomy or not.

### **Research methods**

A retrospective case-matched cohort study was conducted. The abdominal composition was quantified based on computed tomography images by setting Hounsfield Unit thresholds. The abdominal composition related parameters were compared and the importance of these indicators was quantified using feature importance analysis.

### **Research results**

A total of 156 cases were included in this study. Comparing the abdominal composition related parameters demonstrated that patients who developed AL exhibited a larger visceral fat area (VFA,  $125.68 \pm 73.59$  vs  $97.03 \pm 57.66$ ,  $P = 0.008$ ) and a smaller anterior to posterior diameter of abdominal cavity (APD,  $77.30 \pm 23.23$  vs  $92.09 \pm 26.40$ ,  $P < 0.001$ ) and transverse diameter of abdominal cavity (TD,  $22.90 \pm 2.23$  vs  $24.21 \pm 2.90$ ,  $P = 0.002$ ). Feature importance analysis revealed TD, APD, and VFA to be the three most important abdominal composition related parameters.

### **Research conclusions**

Rectal cancer patients who have a higher visceral fat content and a narrower abdominal structure might be at a higher risk of developing AL.

### **Research perspectives**

A narrow abdominal structure is associated with the increased difficulty of the operation and prolonged operation time. In addition, the association of abdominal composition related parameters and postoperative complications was reported. But, whether abdominal composition is associated with AL is not known.

---

## **FOOTNOTES**

**Author contributions:** Liu L and Shao SL contributed to conceptualization and design of the study; Shao SL and Li YK contributed to acquisition of the data; Li YK, Shao SL, and Qin JC contributed to methodology; Liu L, Li YK, and Shao SL contributed to formal analysis; Liu L contributed to software; Qin JC contributed to supervision; Shao SL and Liu L contributed to manuscript writing, review, and editing; all authors contributed to final approval of the version of the manuscript.

**Institutional review board statement:** The study was reviewed and approved by the Local Ethical Committee of Tongji Hospital of Huazhong University of Science and Technology (Approval No. TJ-IRB20210719).

**Informed consent statement:** The patients' consent was waived due to the retrospective nature of the study.

**Conflict-of-interest statement:** The authors declare no conflict of interest for this article.

**Data sharing statement:** Data used in this study is available from the corresponding author on reasonable request.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** China

**ORCID number:** Sheng-Li Shao 0000-0001-8786-0051; Yang-Kun Li 0000-0001-9122-7136; Ji-Chao Qin 0000-0002-2961-7624; Lu Liu 0000-0002-1547-4431.

**S-Editor:** Chen YL

**L-Editor:** Wang TQ

P-Editor: Chen YL

## REFERENCES

- 1 **Kitz J**, Fokas E, Beissbarth T, Ströbel P, Wittekind C, Hartmann A, Rüschoff J, Papadopoulos T, Rösler E, Ortloff-Kittredge P, Kania U, Schlitt H, Link KH, Bechstein W, Raab HR, Staib L, Germer CT, Liersch T, Sauer R, Rödel C, Ghadimi M, Hohenberger W; German Rectal Cancer Study Group. Association of Plane of Total Mesorectal Excision With Prognosis of Rectal Cancer: Secondary Analysis of the CAO/ARO/AIO-04 Phase 3 Randomized Clinical Trial. *JAMA Surg* 2018; **153**: e181607 [PMID: 29874375 DOI: 10.1001/jamasurg.2018.1607]
- 2 **Leonard D**, Penninckx F, Fieuw S, Jouret-Mourin A, Sempoux C, Jehaes C, Van Eycken E; PROCARE, a multidisciplinary Belgian Project on Cancer of the Rectum. Factors predicting the quality of total mesorectal excision for rectal cancer. *Ann Surg* 2010; **252**: 982-988 [PMID: 21107108 DOI: 10.1097/SLA.0b013e3181efc142]
- 3 **Miskovic D**, Foster J, Agha A, Delaney CP, Francis N, Hasegawa H, Karachun A, Kim SH, Law WL, Marks J, Morino M, Panis Y, Uriburu JC, Wexner SD, Parvaiz A. Standardization of laparoscopic total mesorectal excision for rectal cancer: a structured international expert consensus. *Ann Surg* 2015; **261**: 716-722 [PMID: 25072446 DOI: 10.1097/SLA.0000000000000823]
- 4 **Law WL**, Chu KW. Anterior resection for rectal cancer with mesorectal excision: a prospective evaluation of 622 patients. *Ann Surg* 2004; **240**: 260-268 [PMID: 15273550 DOI: 10.1097/01.sla.0000133185.23514.32]
- 5 **Artus A**, Tabchouri N, Iskander O, Michot N, Muller O, Giger-Pabst U, Bourlier P, Bourbao-Tournois C, Kraemer-Bucur A, Lecomte T, Salamé E, Ouaiissi M. Long term outcome of anastomotic leakage in patients undergoing low anterior resection for rectal cancer. *BMC Cancer* 2020; **20**: 780 [PMID: 32819329 DOI: 10.1186/s12885-020-07109-4]
- 6 **Koedam TWA**, Bootsma BT, Deijen CL, van de Brug T, Kazemier G, Cuesta MA, Fürst A, Lacy AM, Haglind E, Tuynman JB, Daams F, Bonjer HJ; COLOR COLOR II study group. Oncological Outcomes After Anastomotic Leakage After Surgery for Colon or Rectal Cancer: Increased Risk of Local Recurrence. *Ann Surg* 2022; **275**: e420-e427 [PMID: 32224742 DOI: 10.1097/SLA.0000000000003889]
- 7 **Denost Q**, Rouanet P, Faucheron JL, Panis Y, Meunier B, Cotte E, Meurette G, Portier G, Sabbagh C, Loriau J, Benoist S, Piessen G, Sielezneff I, Lelong B, Mauvais F, Romain B, Barussaud ML, Capdepon M, Laurent C, Rullier E. Impact of early biochemical diagnosis of anastomotic leakage after rectal cancer surgery: long-term results from GRECCAR 5 trial. *Br J Surg* 2021; **108**: 605-608 [PMID: 33793764 DOI: 10.1093/bjs/znab003]
- 8 **Kawada K**, Sakai Y. Preoperative, intraoperative and postoperative risk factors for anastomotic leakage after laparoscopic low anterior resection with double stapling technique anastomosis. *World J Gastroenterol* 2016; **22**: 5718-5727 [PMID: 27433085 DOI: 10.3748/wjg.v22.i25.5718]
- 9 **Verduin WM**, Warps AK, van den Helder R, Doodeman HJ, Houdijk APJ; INfluences of Fat And Muscle in colorectal Surgery Collaborative. Visceral Fat and Anastomotic Leakage After Colon Cancer Resection. *Dis Colon Rectum* 2021; **64**: 163-170 [PMID: 33394767 DOI: 10.1097/DCR.0000000000001779]
- 10 **Chen Z**, Yang J, Liu Z, Zhang Y, Sun J, Wang P. Which obesity-associated parameters can better reflect the risk of the occurrence of the anastomotic leakage? *Scand J Gastroenterol* 2020; **55**: 466-471 [PMID: 32285713 DOI: 10.1080/00365521.2020.1748223]
- 11 **Daley BJ**, Cecil W, Clarke PC, Cofer JB, Guillaumondegui OD. How slow is too slow? *J Am Coll Surg* 2015; **220**: 550-558 [PMID: 25728140 DOI: 10.1016/j.jamcollsurg.2014.12.040]
- 12 **Thapa B**, Sutanto E, Bhandari R. Thickness of subcutaneous fat is a risk factor for incisional surgical site infection in acute appendicitis surgery: a prospective study. *BMC Surg* 2021; **21**: 6 [PMID: 33397364 DOI: 10.1186/s12893-020-01029-7]
- 13 **Kim JH**, Kim J, Lee WJ, Seong H, Choi H, Ahn JY, Jeong SJ, Ku NS, Son T, Kim HI, Han SH, Choi JY, Yeom JS, Hyung WJ, Song YG, Noh SH. A High Visceral-To-Subcutaneous Fat Ratio is an Independent Predictor of Surgical Site Infection after Gastrectomy. *J Clin Med* 2019; **8** [PMID: 30979055 DOI: 10.3390/jcm8040494]
- 14 **Zager Y**, Khalilieh S, Ganaiem O, Gorgov E, Horesh N, Anteby R, Kopylov U, Jacoby H, Dreznik Y, Dori A, Gutman M, Nevler A. Low psoas muscle area is associated with postoperative complications in Crohn's disease. *Int J Colorectal Dis* 2021; **36**: 543-550 [PMID: 33236229 DOI: 10.1007/s00384-020-03799-1]
- 15 **Fujihata S**, Ogawa R, Nakaya S, Hayakawa S, Okubo T, Sagawa H, Tanaka T, Takahashi H, Matsuo Y, Takiguchi S. The impact of skeletal muscle wasting during neoadjuvant chemotherapy on postoperative anastomotic leakage in patients with esophageal cancer. *Esophagus* 2021; **18**: 258-266 [PMID: 32889673 DOI: 10.1007/s10388-020-00774-1]
- 16 **Baracos VE**, Arribas L. Sarcopenic obesity: hidden muscle wasting and its impact for survival and complications of cancer therapy. *Ann Oncol* 2018; **29**: ii1-ii9 [PMID: 29506228 DOI: 10.1093/annonc/mdx810]
- 17 **Rahbari NN**, Weitz J, Hohenberger W, Heald RJ, Moran B, Ulrich A, Holm T, Wong WD, Tiret E, Moriya Y, Laurberg S, den Dulk M, van de Velde C, Büchler MW. Definition and grading of anastomotic leakage following anterior resection of the rectum: a proposal by the International Study Group of Rectal Cancer. *Surgery* 2010; **147**: 339-351 [PMID: 20004450 DOI: 10.1016/j.surg.2009.10.012]
- 18 **Fujiwara N**, Nakagawa H, Kudo Y, Tateishi R, Taguri M, Watadani T, Nakagomi R, Kondo M, Nakatsuka T, Minami T, Sato M, Uchino K, Enooku K, Kondo Y, Asaoka Y, Tanaka Y, Ohtomo K, Shiina S, Koike K. Sarcopenia, intramuscular fat deposition, and visceral adiposity independently predict the outcomes of hepatocellular carcinoma. *J Hepatol* 2015; **63**: 131-140 [PMID: 25724366 DOI: 10.1016/j.jhep.2015.02.031]
- 19 **Nachit M**, Kwanten WJ, Thissen JP, Op De Beeck B, Van Gaal L, Vonghia L, Verrijken A, Driessen A, Horsmans Y, Francque S, Leclercq IA. Muscle fat content is strongly associated with NASH: A longitudinal study in patients with morbid obesity. *J Hepatol* 2021; **75**: 292-301 [PMID: 33865909 DOI: 10.1016/j.jhep.2021.02.037]
- 20 **McGovern J**, Dolan RD, Horgan PG, Laird BJ, McMillan DC. Computed tomography-defined low skeletal muscle index and density in cancer patients: observations from a systematic review. *J Cachexia Sarcopenia Muscle* 2021; **12**: 1408-1417

- [PMID: 34664431 DOI: 10.1002/jcsm.12831]
- 21 **Jang JH**, Kim HC, Huh JW, Park YA, Cho YB, Yun SH, Lee WY, Yu JI, Park HC, Park YS, Park JO. Anastomotic Leak Does Not Impact Oncologic Outcomes After Preoperative Chemoradiotherapy and Resection for Rectal Cancer. *Ann Surg* 2019; **269**: 678-685 [PMID: 29112004 DOI: 10.1097/SLA.0000000000002582]
  - 22 **Smith JD**, Paty PB, Guillem JG, Temple LK, Weiser MR, Nash GM. Anastomotic leak is not associated with oncologic outcome in patients undergoing low anterior resection for rectal cancer. *Ann Surg* 2012; **256**: 1034-1038 [PMID: 22584695 DOI: 10.1097/SLA.0b013e318257d2c1]
  - 23 **Scarborough JE**, Mantyh CR, Sun Z, Migaly J. Combined Mechanical and Oral Antibiotic Bowel Preparation Reduces Incisional Surgical Site Infection and Anastomotic Leak Rates After Elective Colorectal Resection: An Analysis of Colectomy-Targeted ACS NSQIP. *Ann Surg* 2015; **262**: 331-337 [PMID: 26083870 DOI: 10.1097/SLA.0000000000001041]
  - 24 **Park JS**, Choi GS, Kim SH, Kim HR, Kim NK, Lee KY, Kang SB, Kim JY, Kim BC, Bae BN, Son GM, Lee SI, Kang H. Multicenter analysis of risk factors for anastomotic leakage after laparoscopic rectal cancer excision: the Korean laparoscopic colorectal surgery study group. *Ann Surg* 2013; **257**: 665-671 [PMID: 23333881 DOI: 10.1097/SLA.0b013e31827b8ed9]
  - 25 **Ghuman A**, Ganga R, Parisi Severino N, Krizzuk D, Li QZ, Wexner SD, Da Silva G. Clinical Factors Contributing to Anastomotic Leak After Mid-to-High Colorectal Anastomosis. *Am Surg* 2021; 31348211041555 [PMID: 34519249 DOI: 10.1177/00031348211041555]
  - 26 **Nattenmüller J**, Böhm J, Bagdassarjan A, Kulu Y, Gigic B, Schneider M, Kauczor HU, Ulrich CM, Ulrich A. CT-Quantified Adipose Tissue Distribution: Risk or Protective Factor for Complications after Rectal Cancer Surgery? *Obes Facts* 2019; **12**: 259-271 [PMID: 31055588 DOI: 10.1159/000499320]
  - 27 **Tsuruta A**, Tashiro J, Ishii T, Oka Y, Suzuki A, Kondo H, Yamaguchi S. Prediction of Anastomotic Leakage After Laparoscopic Low Anterior Resection in Male Rectal Cancer by Pelvic Measurement in Magnetic Resonance Imaging. *Surg Laparosc Endosc Percutan Tech* 2017; **27**: 54-59 [PMID: 28092330 DOI: 10.1097/SLE.0000000000000366]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA  
**Telephone:** +1-925-3991568  
**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
**Help Desk:** <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>

