World Journal of *Gastroenterology*

World J Gastroenterol 2023 June 21; 29(23): 3574-3747



Baishidena



WJG

World Journal of VVoria jos. Gastroenterology

Contents

Weekly Volume 29 Number 23 June 21, 2023

EDITORIAL

3574 Recent advances in treatment of nodal and gastrointestinal follicular lymphoma

Watanabe T

MINIREVIEWS

Prognostic role of intestinal ultrasound in Crohn's disease 3595

Manzotti C, Colombo F, Zurleni T, Danelli P, Maconi G

ORIGINAL ARTICLE

Basic Study

3606 BMI-1 activates hepatic stellate cells to promote the epithelial-mesenchymal transition of colorectal cancer cells

Jiang ZY, Ma XM, Luan XH, Liuyang ZY, Hong YY, Dai Y, Dong QH, Wang GY

3622 18β-glycyrrhetinic acid inhibits proliferation of gastric cancer cells through regulating the miR-345-5p/TGM2 signaling pathway

Li X, Ma XL, Nan Y, Du YH, Yang Y, Lu DD, Zhang JF, Chen Y, Zhang L, Niu Y, Yuan L

Retrospective Study

3645 High expression of the circadian clock gene NPAS2 is associated with progression and poor prognosis of gastric cancer: A single-center study

Cao XM, Kang WD, Xia TH, Yuan SB, Guo CA, Wang WJ, Liu HB

3658 SGK3 overexpression correlates with a poor prognosis in endoscopically resected superficial esophageal squamous cell neoplasia: A long-term study

Xu N, Li LS, Li H, Zhang LH, Zhang N, Wang PJ, Cheng YX, Xiang JY, Linghu EQ, Chai NL

3668 Hot snare polypectomy vs endoscopic mucosal resection using bipolar snare for intermediate size colorectal lesions: Propensity score matching

Minakata N, Murano T, Wakabayashi M, Sasabe M, Watanabe T, Mitsui T, Yamashita H, Inaba A, Sunakawa H, Nakajo K, Kadota T, Shinmura K, Ikematsu H, Yano T

3678 Lymphocyte-to-white blood cell ratio is associated with outcome in patients with hepatitis B virus-related acute-on-chronic liver failure

Zhang Y, Chen P, Zhu X

Observational Study

3688 Spatial cluster mapping and environmental modeling in pediatric inflammatory bowel disease Michaux M, Chan JM, Bergmann L, Chaves LF, Klinkenberg B, Jacobson K



Contents

Weekly Volume 29 Number 23 June 21, 2023

Prospective Study

3703 Novel multi-parametric diagnosis of non-alcoholic fatty liver disease using ultrasonography, body mass index, and Fib-4 index

Funada K, Kusano Y, Gyotoku Y, Shirahashi R, Suda T, Tamano M

3715 Robotic-assisted proctosigmoidectomy for Hirschsprung's disease: A multicenter prospective study Zhang MX, Zhang X, Chang XP, Zeng JX, Bian HQ, Cao GQ, Li S, Chi SQ, Zhou Y, Rong LY, Wan L, Tang ST

SYSTEMATIC REVIEWS

Paradoxical association between dyspepsia and autoimmune chronic atrophic gastritis: Insights into 3733 mechanisms, pathophysiology, and treatment options

Rossi RE, Elvevi A, Sciola V, Mandarino FV, Danese S, Invernizzi P, Massironi S



Contents

Weekly Volume 29 Number 23 June 21, 2023

ABOUT COVER

Editorial Board Member of World Journal of Gastroenterology, Kamran Rostami, FRACP, MD, PhD, Department of Gastroenterology, MidCentral District Health Board Palmerston Hospital, Palmerston North 4472, New Zealand. kamran.rostami@midcentraldhb.govt.nz

AIMS AND SCOPE

The primary aim of World Journal of Gastroenterology (WJG, World J Gastroenterol) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. WJG mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The WJG is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports, Index Medicus, MEDLINE, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJG as 4.3; IF without journal self cites: 4.1; 5-year IF: 5.3; Journal Citation Indicator: 0.82; Ranking: 32 among 93 journals in gastroenterology and hepatology; and Quartile category: Q2. The WJG's CiteScore for 2021 is 8.3 and Scopus CiteScore rank 2022: Gastroenterology is 22/149.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yi-Xuan Cai; Production Department Director: Xiang Li; Editorial Office Director: Jia-Ru Fan.

NAME OF JOURNAL	INSTRUCTIONS TO AUTHORS
World Journal of Gastroenterology	https://www.wjgnet.com/bpg/gerinfo/204
ISSN	GUIDELINES FOR ETHICS DOCUMENTS
ISSN 1007-9327 (print) ISSN 2219-2840 (online)	https://www.wjgnet.com/bpg/GerInfo/287
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH
October 1, 1995	https://www.wignet.com/bpg/gerinfo/240
FREQUENCY	PUBLICATION ETHICS
Weekly	https://www.wjgnet.com/bpg/GerInfo/288
EDITORS-IN-CHIEF	PUBLICATION MISCONDUCT
Andrzej S Tarnawski	https://www.wjgnet.com/bpg/gerinfo/208
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE
http://www.wjgnet.com/1007-9327/editorialboard.htm	https://www.wjgnet.com/bpg/gerinfo/242
PUBLICATION DATE June 21, 2023	STEPS FOR SUBMITTING MANUSCRIPTS https://www.wjgnet.com/bpg/GerInfo/239
COPYRIGHT	ONLINE SUBMISSION
© 2023 Baishideng Publishing Group Inc	https://www.f6publishing.com

© 2023 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: bpgoffice@wjgnet.com https://www.wjgnet.com



WJG

World Journal of Gastroenterology

Submit a Manuscript: https://www.f6publishing.com

World J Gastroenterol 2023 June 21; 29(23): 3703-3714

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

DOI: 10.3748/wjg.v29.i23.3703

ORIGINAL ARTICLE

Prospective Study Novel multi-parametric diagnosis of non-alcoholic fatty liver disease using ultrasonography, body mass index, and Fib-4 index

Kei Funada, Yumi Kusano, Yoshinori Gyotoku, Ryosaku Shirahashi, Toshikuni Suda, Masaya Tamano

Specialty type: Gastroenterology and hepatology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0 Grade B (Very good): B Grade C (Good): C Grade D (Fair): 0 Grade E (Poor): 0

P-Reviewer: Pécsi D, Hungary; Sempokuya T, United States

Received: February 8, 2023 Peer-review started: February 8, 2023 First decision: March 7, 2023 Revised: March 14, 2023 Accepted: May 22, 2023 Article in press: May 22, 2023 Published online: June 21, 2023



Kei Funada, Yumi Kusano, Yoshinori Gyotoku, Ryosaku Shirahashi, Toshikuni Suda, Masaya Tamano, Department of Gastroenterology, Dokkyo Medical University Saitama Medical Center, Koshigaya-shi 343-8555, Saitama, Japan

Corresponding author: Masaya Tamano, PhD, Professor, Department of Gastroenterology, Dokkyo Medical University Saitama Medical Center, 2-1-50 Minami-Koshigaya, Koshigayashi 343-8555, Saitama, Japan. mstamano@dokkyomed.ac.jp

Abstract

BACKGROUND

Shear wave speed (SWS), shear wave dispersion (SWD), and attenuation imaging (ATI) are new diagnostic parameters for non-alcoholic fatty liver disease. To differentiate between non-alcoholic steatohepatitis (NASH) and non-alcoholic fatty liver (NAFL), we developed a clinical index we refer to as the "NASH pentagon" consisting of the 3 abovementioned parameters, body mass index (BMI), and Fib-4 index.

AIM

To investigate whether the area of the NASH pentagon we propose is useful in discriminating between NASH and NAFL.

METHODS

This non-invasive, prospective, observational study included patients diagnosed with fatty liver by abdominal ultrasound between September 2021 and August 2022 in whom shear wave elastography, SWD, and ATI were measured. Histological diagnosis based on liver biopsy was performed in 31 patients. The large pentagon group (LP group) and the small pentagon group (SP group), using an area of 100 as the cutoff, were compared; the NASH diagnosis rate was also investigated. In patients with a histologically confirmed diagnosis, receiveroperating characteristic (ROC) curve analyses were performed.

RESULTS

One hundred-seven patients (61 men, 46 women; mean age 55.1 years; mean BMI 26.8 kg/m^2) were assessed. The LP group was significantly older (mean age: $60.8 \pm$ 15.2 years vs 46.4 ± 13.2 years; P < 0.0001). Twenty-five patients who underwent liver biopsies were diagnosed with NASH, and 6 were diagnosed with NAFL. On ROC curve analyses, the areas under the ROC curves for SWS, dispersion slope, ATI value, BMI, Fib-4 index, and the area of the NASH pentagon were 0.88000,



0.82000, 0.58730, 0.63000, 0.59333, and 0.93651, respectively; the largest was that for the area of the NASH pentagon.

CONCLUSION

The NASH pentagon area appears useful for discriminating between patients with NASH and those with NAFL.

Key Words: Non-alcoholic fatty liver disease; Non-alcoholic steatohepatitis; Attenuation imaging; Shear wave elastography; Shear wave dispersion; Diagnosis

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The non-alcoholic steatohepatitis (NASH) pentagon is a novel clinical index consisting of the five parameters of shear wave speed, dispersion slope, attenuation imaging value, Fib-4 index, and body mass index. It is simple and the calculation of its area is easy. The area of the NASH pentagon is useful for discriminating between patients with NASH and those with non-alcoholic fatty liver.

Citation: Funada K, Kusano Y, Gyotoku Y, Shirahashi R, Suda T, Tamano M. Novel multi-parametric diagnosis of non-alcoholic fatty liver disease using ultrasonography, body mass index, and Fib-4 index. *World J Gastroenterol* 2023; 29(23): 3703-3714

URL: https://www.wjgnet.com/1007-9327/full/v29/i23/3703.htm DOI: https://dx.doi.org/10.3748/wjg.v29.i23.3703

INTRODUCTION

Non-alcoholic fatty liver disease (NAFLD) is a global public health problem[1,2]. It is a grouping of diseases consisting of either non-alcoholic fatty liver (NAFL) or non-alcoholic steatohepatitis (NASH) [3]. If NASH progresses, it can lead to liver cirrhosis and hepatocellular carcinoma; thus, a clinical index needs to be developed that can efficiently discriminate NASH patients from NAFL patients[4,5]. Assessments based on multiple parameters using ultrasound, including shear wave elastography (SWE), are reported to be useful in stratifying the risk of advanced NASH[6].

SWE measures the speed of shear waves (shear wave speed; SWS) generated by push pulses. SWS is slow in soft objects and fast in hard objects. Measurements of SWS by SWE are considered useful in assessing liver fibrosis in viral hepatitis and NAFLD[7-11].

The shear wave dispersion (SWD) developed in recent years obtains the "frequency dependence" (dispersion slope; DS), which is the extent of the changes in the speed of shear waves, from changes in their frequency. Some preliminary evidence has been reported that DS measurements with SWD are an index of necroinflammation of the liver[12,13].

Attenuation imaging (ATI) was developed as a new testing method for evaluation of hepatic steatosis with ultrasound technology. ATI is a technique that uses the principle of attenuation due to phenomena such as absorption and diffusion of ultrasonic pulses emitted within the body when they pass through body tissue, and it produces a value referred to as the attenuation coefficient. Magnetic resonance imaging (MRI)-determined proton density fat fraction (PDFF) values have been reportedly used for assessing histological hepatic steatosis grades[14], and Tada *et al*[15] reported a good correlation between ATI values and MRI-determined PDFF values.

On the other hand, it is important to have parameters that do not require special equipment to measure. The prevalence of NAFLD in Japanese non-obese patients is 15%[16]. In other words, 85% of NAFLD patients are obese. Therefore, the body mass index (BMI), which can be easily used to determine obesity, is one such parameter of importance.

Similarly, the scoring system for liver fibrosis by the biochemical examination of blood is also relevant. The Fib-4 index, calculated by aspartate aminotransferase (AST), alanine aminotransferase (ALT), platelets (Plts), and age, is simple and easy to use[17].

In this study, a pentagon consisting of five parameters was prepared with the addition of the BMI and the Fib-4 index to the three parameters of SWS, SWD, and ATI. Whether the area of this pentagon is useful in discriminating NASH from NAFL was then investigated.

Raishideng® WJG | https://www.wjgnet.com

MATERIALS AND METHODS

Patients

This was a non-invasive, prospective, observational study that conformed with the ethical guidelines of the Declaration of Helsinki in its 2008 revision and was approved by the ethics review board of the hospital with which the authors are affiliated (No. 21057). The study was conducted after complete information disclosure on the website of the authors' hospital.

Of the patients diagnosed with fatty liver by abdominal ultrasound between September 2021 and August 2022, the subjects were patients in whom SWE, SWD, and ATI were measured. Patients with a history of alcohol intake of ethanol ≥ 20 g/d, who had hepatitis B, hepatitis C, or autoimmune liver disease, and patients with concurrent drug-induced liver injury or cholangitis were excluded.

Diagnosis of fatty liver with ultrasonography

The ultrasound scanner used was an Aplio i800 from Canon Medical Systems Corp. (Otawara, Tochigi, Japan). Fatty liver was diagnosed if the following were seen on B-mode ultrasound examination: (1) Bright liver[18]; (2) Positive hepatorenal echo contrast[19]; (3) Deep ultrasound attenuation in the liver [20]; and (4) Vascular blurring in the liver[21]. Specialists with ten or more years of ultrasonographic experience (Suda T and Tamano M) performed the B-mode ultrasound examination and measured SWE, SWD, and ATI.

Measurements of shear wave elastography and SWD

All SWE measurements were done following B-mode scans, using the same diagnostic equipment and transducers. The measurement results are shown as SWS (m/s). A 1-cm-diameter, circular region of interest was placed on the sample box.

SWD was measured at the same time as SWE. Simultaneous measurements could be taken by switching to quad view mode including a SWS map and a SWD slope map. This test was also performed using five or more measurement values, and the measurement results are shown as DS[(m/s)/KHz].

Measurements of attenuation imaging

After SWE and SWD were measured, the same physician performed the ATI test. The measurement result is shown as the ATI value [(dB/cm)/MHz]. A sample box with the default settings was used to acquire the data, set at least 1.5 cm below the liver capsule to avoid reverberation artifacts. The reliability of the results is displayed with R^2 values, categorized as poor ($R^2 < 0.80$), good ($R^2 = 0.81-0.89$), or excellent ($R^2 > 0.90$). According to information provided by the manufacturer, measurements are considered to be valid when $R^2 \ge 0.80$. Attenuation imaging examinations were performed until five valid measurements were obtained, and the mean value of the five measurements was used in the analysis.

Structure of the NASH pentagon

AST, ALT, and Plts were measured on the same day as the abdominal ultrasound. The Fib-4 index was calculated from these values and age. The BMI was calculated from height and weight on the day on which abdominal ultrasound was performed.

A pentagon consisting of these five parameters (SWS, DS, ATI value, BMI, and Fib-4 index) was created. This was defined as the NASH pentagon. The area of the NASH pentagon was calculated automatically by inputting a calculation formula into an Aplio i800 work station.

The standard value for each item was set according to previous studies as follows. The standard value of SWS was set as 1.33 m/s (cutoff value of fibrosis stage 1), that of DS was set to 8.5 (m/s)/KHz (cutoff value of lobular inflammation grade 1)[22]. Similarly, the standard value of ATI was set to 0.66 (dB/cm)/MHz, defining steatosis grade 1[16,22]. The standard value for BMI was set at 25.0 kg/m² based on the index of the Japanese Society for the Study of Obesity[1]. The standard value for the Fib-4 index was set to 1.30, which is the low cutoff value for advanced fibrosis (stage 3-4), based on a report by Shah et al[24] The investigation was done taking the area of this pentagon as 100 (Figure 1).

Diagnosis of NASH

Histological diagnosis based on liver biopsy was performed within three months from SWE measurement in those patients who provided consent. NASH diagnosis was made by an experienced pathologist certified by the Japanese Society of Pathology, based on comprehensive findings including macrovesicular steatosis, ballooning degeneration of hepatocytes, scattered inflammation and apoptotic bodies in the lobules, Mallory-Denk bodies, and fibrosis[25-27]. In this study, all patients without liver biopsy were judged to have NAFL, even if NASH was suspected clinically.

Comparison of two groups classified by the area of the NASH pentagon

Patient characteristics and individual parameters were compared between a large pentagon group (LP group), in which the area of the NASH pentagon was 100 or greater, and a small pentagon group (SP





Figure 1 Non-alcoholic steatohepatitis pentagon. The non-alcoholic steatohepatitis pentagon consists of the five parameters of shear wave speed (SWS), dispersion slope (DS), attenuation imaging value (ATI value), Fib-4 index, and the body mass index (BMI). Based on past reports, the reference values were set as SWS 1.33 m/s, DS 8.5 (m/s)/KHz, ATI value 0.66 (dB/cm)/MHz, BMI 25.0 kg/m², and Fib-4 index 1.3. SWS: Shear wave speed; DS: Dispersion slope; BMI: Body mass index; ATI: Attenuation imaging.

group), in which the area was less than 100. In addition, the NASH diagnosis rate was investigated when the NASH pentagon area of 100 was taken as the reference.

Diagnostic accuracy for NASH using the area of the NASH pentagon

In patients for whom liver biopsies were performed during the study period and a histologically confirm-ed diagnosis was made, a receiver-operating characteristic (ROC) curve was drawn for SWS, DS, ATI value, BMI, Fib-4 index, and area of the NASH pentagon.

Statistical analysis

Continuous data for SWS, DS, ATI value, and other clinical parameters are expressed as mean \pm SD. A Spearman rank-order correlation coefficient test was used to test the independency of these clinical parameters. A non-paired Wilcoxon test was used in comparisons of each parameter between the two groups divided by NASH pentagon area, with *P* < 0.05 taken to indicate a significant difference.

The NASH diagnostic performance of SWS, DS, ATI value, BMI, Fib-4 index, and the area of the pentagon was investigated using the ROC curve. The area under the ROC curve (AUROC) was used to evaluate NASH diagnostic performance.

RESULTS

Patients' characteristics

The number of patients who were diagnosed with fatty liver based on abdominal ultrasound and in whom SWE, SWD, and ATI were measured was 126. After excluding 7 patients with an alcohol intake history of \geq 20 g ethanol/d, 10 patients with concurrent hepatitis B, hepatitis C, or autoimmune liver injury, and 2 patients for other reasons, the investigation was conducted with the remaining 107 patients (Figure 2).

The characteristics for these 107 patients are shown in Table 1. There were 61 men and 46 women, with a mean age of 55.1 years. The mean BMI was 26.8 kg/m². Of the 107 patients, diabetes mellitus was seen in 25, dyslipidemia in 35, and hypertension in 37. Mean AST, ALT, and γ -glutamyltransferase concentrations were all mildly elevated at 44.9 U/L, 70.6 U/L, and 99.6 U/L, respectively. The mean Fib-4 index was 1.53. Twenty-five of the 107 patients were diagnosed with NASH, and the remaining 82 were diagnosed with NAFL.

The Spearman rank correlation coefficient between SWS and the Fib-4 index was 0.3486; no strong correlation was found. Similarly, the rank correlation coefficient between ATI and BMI was 0.1955, showing no correlation.

Table 1 Patients' characteristics	
Patients' characteristics	<i>n</i> = 107
M/F	61/46
Age (yr)	55.1 ± 15.9
BMI (kg/m ²)	26.8 ± 4.1
Metabolic diseases	
Diabetes mellitus (yes/no)	25/82
Dyslipidemia (yes/no)	35/72
Hypertension (yes/no)	37/70
Concomitant drugs	
SGLT2 inhibitor	12
DPP-4 inhibitor	13
Thiazolidinedione	2
GLP-1 agonist	2
Statin	19
Bezafibrate	0
Pemafibrate	8
EPA and DHA preparation	1
AST (U/L)	44.9 ± 38.0
ALT (U/L)	70.6 ± 57.7
GGT (U/L)	99.6 ± 139.7
T-B (mg/dL)	0.9 ± 0.5
Alb (mg/dL)	4.5 ± 0.4
eGFR (mL/min)	73.0 ± 15.0
HbA1c (%)	6.5 ± 1.2
T-chol (mg/dL)	202.1 ± 52.6
TG (mg/dL)	180.0 ± 119.0
WBC (10 ³ /µL)	6731.8 ± 1882.5
Hb (g/dL)	14.7 ± 1.3
Plts $(10^4/\mu L)$	23.9 ± 7.6
Fib-4 index	1.53 ± 1.90

M: Male; F: Female; AST: Aspartate aminotransferase; ALT: Alanine aminotransferase; GGT: γ-glutamyltransferase; T-Bil: Total bilirubin; Alb: Albumin; eGFR: Estimated glomerular filtration rate; HbA1c: Glycated hemoglobin A1c; T-chol: Total cholesterol; TG: Triglycerides; WBC: White blood cells; Hb: Hemoglobin; Plts: Platelets.

Comparison of two groups classified by the area of the NASH pentagon

The LP group with a NASH pentagon area \geq 100 had 64 patients, and the SP group with an area < 100 had 43 patients. A comparison of the two groups is shown in Table 2.

There was no difference in sex between the two groups, but mean age was 60.8 ± 15.2 years in the LP group and 46.4 ± 13.2 years in the SP group, with the LP group significantly older (P < 0.0001).

Concurrent diabetes mellitus was seen in 18/64 patients (28.1%) in the LP group and 6/43 patients (13.9%) in the SP group. While this represents a higher tendency in the LP group, no significant difference was seen (P = 0.0849). Concurrent dyslipidemia was seen in 27/64 patients (42.2%) in the LP group and 8/43 patients (18.6%) in the SP group, and concurrent hypertension was seen in 29/64 patients (45.3%) and 7/43 patients (16.3%), respectively; both were significantly higher in the LP group (P = 0.0108, P = 0.0183).

Table 2 Comparison of the two groups classified by area of non-alcoholic steatohepatitis pentagon				
	Large pentagon group (<i>n</i> = 64)	Small pentagon group (<i>n</i> = 43)	<i>P</i> value	
M/F	33/31	28/15	0.165	
Age (yr)	60.8 ± 15.2	46.4 ± 13.2	< 0.0001	
Metabolic diseases				
Diabetes mellitus (yes/no)	18/46	6/37	0.0849	
Dyslipidemia (yes/no)	27/37	8/35	0.0108	
Hypertension (yes/no)	29/35	7/36	0.0183	
AST (U/L)	52.6 ± 44.2	40.3 ± 24.4	0.0521	
ALT (U/L)	61.1 ± 45.2	83.0 ± 71.6	0.1699	
GGT (U/L)	97.3 ± 123.2	102.9 ± 162.0	0.7238	
T-B (mg/dL)	1.01 ± 0.6	0.77 ± 0.3	0.0044	
Alb (mg/dL)	4.5 ± 0.4	4.6 ± 0.4	0.2843	
eGFR (mL/min)	71.8 ± 16.4	74.7 ± 12.1	0.1639	
HbA1c (%)	6.5 ± 1.1	6.3 ± 1.2	0.2847	
T-chol (mg/dL)	188.5 ± 59.4	221.3 ± 33.8	0.008	
TG (mg/dL)	172.9 ± 126.5	189.9 ± 108.8	0.1278	
WBC (10 ³ /µL)	6225.0 ± 1586.9	7463.9 ± 2050.9	0.008	
Hb (g/dL)	14.7 ± 1.3	14.7 ± 1.5	0.6836	
Plts (10 ⁴ /µL)	21.0 ± 6.5	28.8 ± 6.6	< 0.0001	
SWS (m/s)	1.78 ± 0.36	1.39 ± 0.14	< 0.0001	
DS[(m/s)/kHz]	13.3 ± 3.7	10.8 ± 2.0	< 0.0001	
ATI value [(dB/cm)/MHz]	0.78 ± 0.13	0.74 ± 0.13	0.1047	
Fib-4 index	2.2 ± 2.3	0.7 ± 0.4	< 0.0001	
BMI (kg/m ²)	27.8 ± 4.4	25.6 ± 3.2	0.0249	
NASH/NAFL	22/42	3/40	0.0042	

M: Male; F: Female; AST: Aspartate aminotransferase; ALT: Alanine aminotransferase; GGT: γ-glutamyltransferase; T-chol: Total cholesterol; TG: Triglycerides; WBC: White blood cells; Hb: Hemoglobin; Plts: Platelets; SWS: Shear wave speed; DS: Dispersion slope; ATI value: Attenuation imaging value; BMI: Body mass index; Alb: Albumin; eGFR: Estimated glomerular filtration rate; NASH: Non-alcoholic steatohepatitis; NAFL: Non-alcoholic fatty liver.

On blood biochemistry tests, total bilirubin was significantly higher in the LP group (P = 0.0044), whereas total cholesterol, white blood cells, and Plts were significantly lower in the LP group (P = 0.0080, P = 0.0080, P < 0.0001). There were no significant differences between the two groups in any of the other test results.

Comparisons of the five parameters that make up the NASH pentagon showed no significant difference between the groups in the ATI value (P = 0.1407), and significantly higher values in the LP group for SWS, DS, Fib-4 index, and BMI (P < 0.0001, P = 0.0004, P < 0.0001, P = 0.0249).

The number of patients diagnosed with NASH was 22 (34.3%) of 64 in the LP group and 3 (7.0%) of 43 in the SP group. The LP group had a significantly higher percentage of NASH patients (P = 0.0042).

Diagnostic accuracy of histological NAFLD using the area of the NASH pentagon

Liver biopsy was performed during the period of this study in 31 patients. Twenty-five of these 31 patients were diagnosed with NASH, and six were diagnosed with NAFL. The ROC curves indicating the NASH diagnostic performance in these 31 patients are shown in Figure 3. The AUROCs for SWS, DS, ATI value, BMI, Fib-4 index, and the area of the NASH pentagon were 0.88000, 0.82000, 0.58730, 0.63000, 0.59333, and 0.93651, respectively. Of these six items, the largest AUROC was that for the area of the NASH pentagon.

Boishidena® WJG https://www.wjgnet.com



DOI: 10.3748/wjg.v29.i23.3703 Copyright ©The Author(s) 2023.

Figure 2 Flowchart of the study. A total of 126 patients who were diagnosed with fatty liver based on abdominal ultrasound and in whom shear wave elastography, shear wave dispersion, and attenuation imaging were measured during the period of this study were entered in the study. Of them, 7 patients with an alcohol intake history of 20 g ethanol/d or more, 3 with hepatitis B, 3 with hepatitis C, and 4 with autoimmune hepatitis or primary biliary cholangitis were excluded. In addition, 1 patient with drug-induced liver injury and 1 patient with cholangitis were excluded, bringing the final number of subjects in this investigation to 107. HBV: Hepatitis B virus; HCV: Hepatitis C virus; AIH: Autoimmune hepatitis; PBC: Primary biliary cholangitis; NAFLD: Non-alcoholic fatty liver disease.



Figure 3 Receiver-operating characteristic curve for each parameter. This shows the receiver-operating characteristic (ROC) curve of the 17 patients who underwent liver biopsy during the period of this study. The areas under the ROC curves (AUROCs) of shear wave speed, dispersion slope, attenuation imaging value, body mass index, Fib-4 index, and the area of the non-alcoholic steatohepatitis (NASH) pentagon are 0.88000, 0.82000, 0.58730, 0.63000, 0.59333, and 0.93651, respectively. Of these six parameters, the largest AUROC is that for the area of the NASH pentagon. SWE: Shear wave elastography; SWD: Shear wave dispersion; ATI: Attenuation imaging; BMI: Body mass index; NASH: Non-alcoholic steatohepatitis.

DISCUSSION

Ultrasound elastography has been used to evaluate fibrosis noninvasively. SWE, one type of ultrasound



elastography, includes transient elastography, point SWE, and other techniques[28-30]. The Aplio i800 used in the present study can measure SWE, SWD, and ATI with the same transducer. This machine is also equipped with a function that automatically creates radar charts from these measured values. The radar charts are created using five parameters. The three parameters of SWS, DS, and ATI value are set in the initial settings, and the remaining two can be freely set by the user. In the present study, these two were set as BMI and Fib-4 index, and it was thought that the area of this pentagon (NASH pentagon) may be useful for the diagnosis of NASH.

There are various reports about the cutoff values of SWE and DS[31]. In the present study, the cutoff values of SWE and DS were set from the report of Sugimoto *et al*[22] based on 120 liver biopsies. The cutoff values for steatosis grade 1 of ATI are reported to be 0.62-0.79[32-34]. In the present study, it was set to 0.66 from the report of Tada *et al*[35] based on 148 liver biopsies.

Obesity is the most important factor in the development of NAFLD. In NASH patients, there is a positive correlation between visceral fat and the amount of fat in hepatocytes[36]. Thus, obesity can lead to a diagnosis of NASH. In the present study, the reference value for BMI was set at 25.0 kg/m² based on the index of the Japanese Society for the Study of Obesity and a WHO recommendation[23,37].

The Fib-4 index calculated from age, AST, ALT, and platelet count is regularly used as a noninvasive parameter of hepatic fibrosis[38]. In this study, a value of 1.30, which is the low cutoff value for advanced fibrosis (stage 3-4), based on a report by Shah *et al*[24], was set as the reference value.

SWE and the Fib-4 index are indices of liver fibrosis. However, SWE makes use of ultrasound technology and predicts fibrosis. The Fib-4 index is calculated from biochemical evaluation and age. The correlation of the factors was examined, and no strong correlation was found. ATI is an index of hepatic steatosis assessed using ultrasound technology. BMI is an index of obesity of patients calculated from height and weight. In the present statistical analysis, there was no correlation between these two factors. Due to the above-mentioned reasons, the Fib-4 index and BMI were adopted as constituent factors of the NASH pentagon.

In a comparison of the LP group with a NASH pentagon area of 100 or more and the SP group with an area of less than 100, the LP group was older and had a lower platelet count. This is thought to be a result of the strong involvement of the Fib-4 index among the constituent elements of the pentagon. In fact, a very large difference was seen in the Fib-4 index between the groups, which was 2.2 in the LP group and 0.7 in the SP group (P < 0.0001).

The mean BMI, on the other hand, was 27.8 kg/m² in the LP group and 25.6 kg/m² in the SP group, tending to be higher in the SP group. Obesity is an important element of NAFLD, but as NAFL progresses to NASH, there is a tendency for weight loss. Moreover, even in the SP group, the mean BMI of 25.6 kg/m² was only slightly above the reference of 25.0 kg/m². This result is also consistent with reports that nonobese NAFLD is more common in Asia than in western countries[39,40].

The onset and progression of NAFLD/NASH are related to metabolic syndrome, and they have an especially strong relationship with diabetes mellitus[41]. In the present study, no difference was seen in concurrent diabetes mellitus by differences in the NASH pentagon area, but concurrent dyslipidemia and hypertension were significantly higher in the LP group.

The NASH pentagon consists of five parameters: Fibrosis based on SWS, inflammation as assessed by DS, steatosis as assessed by ATI value, obesity assessed by BMI, and the level of serological and clinical progression based on the Fib-4 index. Once this pentagon is programmed in the ultrasound machine, it can be automatically created and its area subsequently calculated, making it very simple to use. We thus consider it to be a useful method for identifying NASH among NAFLD patients.

In fact, when ROC curves were made for the six parameters of SWS, DS, ATI value, BMI, Fib-4 index, and the area of the NASH pentagon in 31 patients who underwent liver biopsy during the study period, the largest AUROC was the one for the area of the NASH pentagon.

In the early stage of NASH, liver fibrosis is mild, and in these patients, SWS is expected to be either normal or only mildly elevated. In young patients, the Fib-4 index also tends to be low. Even in these patients, however, DS, ATI value, and BMI are expected to be high, so it may be possible to identify them using the NASH pentagon.

Conversely, BMI tends to decrease when NASH progresses to cirrhosis, and even histologically, a state is present in which reduced fat deposition is seen (so-called "burned-out NASH")[42,43]. Such patients are expected to have low BMI and ATI values in conjunction with high SWS and Fib-4 index values. DS may have different values depending on the individual case, but the NASH pentagon is thought to be useful even in these patients.

Limitations

This was a prospective study, but only 107 patients were included, and only 31 underwent liver biopsy during the study period. These numbers are too small to obtain sufficient results. Additional studies with a larger number of patients will be needed to confirm our results. It may also be necessary to conduct a more detailed pathological investigation and a review of the reference values for the five parameters.

Zaishideng® WJG | https://www.wjgnet.com

CONCLUSION

The preparation of a NASH pentagon consisting of five parameters, SWS, DS, ATI value, Fib-4 index, and BMI, is simple and the calculation of its area is easy. The authors believe that this multi-parametric index will be useful for identifying NASH patients among NAFLD patients.

ARTICLE HIGHLIGHTS

Research background

Non-alcoholic fatty liver disease (NAFLD) is a major problem throughout the world. If non-alcoholic steatohepatitis (NASH) progresses, it can lead to hepatocellular carcinoma. Novel clinical index needs to be developed that can efficiently discriminate NASH patients from non-alcoholic fatty liver (NAFL) patients.

Research motivation

Shear wave speed (SWS), shear wave dispersion (SWD), and attenuation imaging (ATI) are new ultrasound diagnostic parameters for NAFLD. We developed a novel clinical index as the "NASH pentagon" consisting of the 3 abovementioned parameters, body mass index (BMI), and Fib-4 index.

Research objectives

Objective of this study is to prove the utility of NASH pentagon area in the differentiation of NASH and NAFL.

Research methods

Patients diagnosed with fatty liver by abdominal ultrasound between September 2021 and August 2022 were enrolled. Histological diagnosis based on liver biopsy was performed in 31 patients. The large pentagon group (LP group) and the small pentagon group (SP group), using an area of 100 as the cutoff, were compared; the NASH diagnosis rate was also investigated. In patients with a histologically confirmed diagnosis, receiver-operating characteristic curve analyses were performed.

Research results

The preparation of a NASH pentagon consisting of five parameters, SWS, DS, ATI value, Fib-4 index, and BMI, is simple and the calculation of its area is easy. The LP group with a NASH pentagon area \geq 100 had 64 patients, and the SP group with an area < 100 had 43 patients. The number of patients diagnosed with NASH was 22 (34.3%) of 64 in the LP group and 3 (7.0%) of 43 in the SP group. The LP group had a significantly higher percentage of NASH patients (P = 0.0042).

Research conclusions

The NASH pentagon is a novel multi-parametric index, and will be useful for identifying NASH patients among NAFLD patients.

Research perspectives

NASH pentagon should be tried in more patients. Besides, the area of NASH pentagon and the details of the histologic diagnosis should be considered.

ACKNOWLEDGEMENTS

The authors would like to thank the institutions that participated in the working group and everyone who helped collect clinical data.

FOOTNOTES

Author contributions: Funada K, Kusano Y, Gyotoku Y and Shirahashi R conceptualized and designed the study, collected data, carried out the initial analysis, and drafted the initial manuscript; Suda T and Tamano M coordinated and supervised data collection and critically reviewed the manuscript for important intellectual content; and all authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

Institutional review board statement: The study was reviewed and approved by the Dokkyo Medical University Saitama Medical Center Institutional Review Board (approval No. 21057).



Clinical trial registration statement: This study is registered at clinical research support office, Dokkyo Medical University Saitama Medical Center. The registration identification number is 21057.

Informed consent statement: Patients were not required to give informed consent for participation in the present study because the analysis used anonymous clinical data that were obtained after each patient gave informed consent to treatment. For full disclosure, the details of this prospective, observational study were published on the home page of the medical center.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Data sharing statement: No additional data are available.

CONSORT 2010 statement: The authors have read the CONSORT 2010 statement, and the manuscript was prepared and revised according to the CONSORT 2010 statement.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is noncommercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country/Territory of origin: Japan

ORCID number: Kei Funada 0000-0003-1190-5588; Yumi Kusano 0000-0001-8000-0884; Yoshinori Gyotoku 0000-0003-2395-3421; Ryosaku Shirahashi 0000-0001-7843-3138; Toshikuni Suda 0000-0003-1017-3725; Masaya Tamano 0000-0001-5595-2330.

S-Editor: Gao CC L-Editor: A P-Editor: Wang JJ

REFERENCES

- 1 Sayiner M, Koenig A, Henry L, Younossi ZM. Epidemiology of Nonalcoholic Fatty Liver Disease and Nonalcoholic Steatohepatitis in the United States and the Rest of the World. Clin Liver Dis 2016; 20: 205-214 [PMID: 27063264 DOI: 10.1016/j.cld.2015.10.001]
- Asrani SK, Devarbhavi H, Eaton J, Kamath PS. Burden of liver diseases in the world. J Hepatol 2019; 70: 151-171 2 [PMID: 30266282 DOI: 10.1016/j.jhep.2018.09.014]
- 3 Yeh MM, Brunt EM. Pathological features of fatty liver disease. Gastroenterology 2014; 147: 754-764 [PMID: 25109884 DOI: 10.1053/j.gastro.2014.07.056]
- Tsochatzis EA, Newsome PN. Non-alcoholic fatty liver disease and the interface between primary and secondary care. 4 Lancet Gastroenterol Hepatol 2018; 3: 509-517 [PMID: 29893235 DOI: 10.1016/S2468-1253(18)30077-3]
- Younossi ZM. Non-alcoholic fatty liver disease A global public health perspective. J Hepatol 2019; 70: 531-544 [PMID: 5 30414863 DOI: 10.1016/j.jhep.2018.10.033]
- Kuroda H, Fujiwara Y, Abe T, Nagasawa T, Oguri T, Noguchi S, Kamiyama N, Takikawa Y. Two-dimensional shear 6 wave elastography and ultrasound-guided attenuation parameter for progressive non-alcoholic steatohepatitis. PLoS One 2021; 16: e0249493 [PMID: 33826669 DOI: 10.1371/journal.pone.0249493]
- Suda T, Okawa O, Masaoka R, Gyotoku Y, Tokutomi N, Katayama Y, Tamano M. Shear wave elastography in hepatitis C patients before and after antiviral therapy. World J Hepatol 2017; 9: 64-68 [PMID: 28105260 DOI: 10.4254/wjh.v9.i1.64]
- Tada T, Kumada T, Toyoda H, Ito T, Sone Y, Okuda S, Tsuji N, Imayoshi Y, Yasuda E. Utility of real-time shear wave 8 elastography for assessing liver fibrosis in patients with chronic hepatitis C infection without cirrhosis: Comparison of liver fibrosis indices. Hepatol Res 2015; 45: E122-E129 [PMID: 25580959 DOI: 10.1111/hepr.12476]
- Meyer G, Dauth N, Grimm M, Herrmann E, Bojunga J, Friedrich-Rust M. Shear Wave Elastography Reveals a High Prevalence of NAFLD-related Fibrosis even in Type 1 Diabetes. Exp Clin Endocrinol Diabetes 2022; 130: 532-538 [PMID: 34784620 DOI: 10.1055/a-1666-0431]
- Bâldea V, Bende F, Popescu A, Şirli R, Sporea I. Comparative study between two 2D-Shear Waves Elastography 10 techniques for the non-invasive assessment of liver fibrosis in patients with chronic hepatitis C virus (HCV) infection. Med Ultrason 2021; 23: 257-264 [PMID: 33657193 DOI: 10.11152/mu-2863]
- Podrug K, Sporea I, Lupusoru R, Pastrovic F, Mustapic S, Bâldea V, Bozin T, Bokun T, Salkic N, Şirli R, Popescu A, 11 Puljiz Z, Grgurevic I. Diagnostic Performance of 2-D Shear-Wave Elastography with Propagation Maps and Attenuation Imaging in Patients with Non-Alcoholic Fatty Liver Disease. Ultrasound Med Biol 2021; 47: 2128-2137 [PMID: 33985827 DOI: 10.1016/j.ultrasmedbio.2021.03.025]
- Sugimoto K, Moriyasu F, Oshiro H, Takeuchi H, Yoshimasu Y, Kasai Y, Furuichi Y, Itoi T. Viscoelasticity Measurement 12 in Rat Livers Using Shear-Wave US Elastography. Ultrasound Med Biol 2018; 44: 2018-2024 [PMID: 29936025 DOI: 10.1016/j.ultrasmedbio.2018.05.008]



- Lee DH, Lee JY, Bae JS, Yi NJ, Lee KW, Suh KS, Kim H, Lee KB, Han JK. Shear-Wave Dispersion Slope from US 13 Shear-Wave Elastography: Detection of Allograft Damage after Liver Transplantation. Radiology 2019; 293: 327-333 [PMID: 31502939 DOI: 10.1148/radiol.2019190064]
- 14 Jeon SK, Lee JM, Joo I, Yoon JH, Lee DH, Lee JY, Han JK. Prospective Evaluation of Hepatic Steatosis Using Ultrasound Attenuation Imaging in Patients with Chronic Liver Disease with Magnetic Resonance Imaging Proton Density Fat Fraction as the Reference Standard. Ultrasound Med Biol 2019; 45: 1407-1416 [PMID: 30975533 DOI: 10.1016/j.ultrasmedbio.2019.02.008]
- Tada T, Kumada T, Toyoda H, Nakamura S, Shibata Y, Yasuda S, Watanuki Y, Tsujii K, Fukuda N, Fujioka M, 15 Takeshima K, Niwa F, Ogawa S, Hashinokuchi S, Kataoka S, Ichikawa H, Iijima H. Attenuation imaging based on ultrasound technology for assessment of hepatic steatosis: A comparison with magnetic resonance imaging-determined proton density fat fraction. Hepatol Res 2020; 50: 1319-1327 [PMID: 32876367 DOI: 10.1111/hepr.13563]
- Nishioji K, Sumida Y, Kamaguchi M, Mochizuki N, Kobayashi M, Nishimura T, Yamaguchi K, Itoh Y. Prevalence of and 16 risk factors for non-alcoholic fatty liver disease in a non-obese Japanese population, 2011-2012. J Gastroenterol 2015; 50: 95-108 [PMID: 24619537 DOI: 10.1007/s00535-014-0948-9]
- Vilar-Gomez E, Chalasani N. Non-invasive assessment of non-alcoholic fatty liver disease: Clinical prediction rules and 17 blood-based biomarkers. J Hepatol 2018; 68: 305-315 [PMID: 29154965 DOI: 10.1016/j.jhep.2017.11.013]
- Joseph AE, Dewbury KC, McGuire PG. Ultrasound in the detection of chronic liver disease (the "bright liver"). Br J 18 Radiol 1979; 52: 184-188 [PMID: 435696 DOI: 10.1259/0007-1285-52-615-184]
- Yajima Y, Ohta K, Narui T, Abe R, Suzuki H, Ohtsuki M. Ultrasonographical diagnosis of fatty liver: significance of the 19 liver-kidney contrast. Tohoku J Exp Med 1983; 139: 43-50 [PMID: 6220488 DOI: 10.1620/tjem.139.43]
- 20 Hamaguchi M, Kojima T, Itoh Y, Harano Y, Fujii K, Nakajima T, Kato T, Takeda N, Okuda J, Ida K, Kawahito Y, Yoshikawa T, Okanoue T. The severity of ultrasonographic findings in nonalcoholic fatty liver disease reflects the metabolic syndrome and visceral fat accumulation. Am J Gastroenterol 2007; 102: 2708-2715 [PMID: 17894848 DOI: 10.1111/j.1572-0241.2007.01526.x
- 21 Saadeh S, Younossi ZM, Remer EM, Gramlich T, Ong JP, Hurley M, Mullen KD, Cooper JN, Sheridan MJ. The utility of radiological imaging in nonalcoholic fatty liver disease. Gastroenterology 2002; 123: 745-750 [PMID: 12198701 DOI: 10.1053/gast.2002.35354]
- Sugimoto K, Moriyasu F, Oshiro H, Takeuchi H, Abe M, Yoshimasu Y, Kasai Y, Sakamaki K, Hara T, Itoi T. The Role 22 of Multiparametric US of the Liver for the Evaluation of Nonalcoholic Steatohepatitis. Radiology 2020; 296: 532-540 [PMID: 32573385 DOI: 10.1148/radiol.2020192665]
- 23 Examination Committee of Criteria for 'Obesity Disease' in Japan; Japan Society for the Study of Obesity. New criteria for 'obesity disease' in Japan. Circ J 2002; 66: 987-992 [PMID: 12419927 DOI: 10.1253/circj.66.987]
- 24 Shah AG, Lydecker A, Murray K, Tetri BN, Contos MJ, Sanyal AJ; Nash Clinical Research Network. Comparison of noninvasive markers of fibrosis in patients with nonalcoholic fatty liver disease. Clin Gastroenterol Hepatol 2009; 7: 1104-1112 [PMID: 19523535 DOI: 10.1016/j.cgh.2009.05.033]
- Angulo P. Nonalcoholic fatty liver disease. N Engl J Med 2002; 346: 1221-1231 [PMID: 11961152 DOI: 25 10.1056/NEJMra011775]
- 26 Matteoni CA, Younossi ZM, Gramlich T, Boparai N, Liu YC, McCullough AJ. Nonalcoholic fatty liver disease: a spectrum of clinical and pathological severity. Gastroenterology 1999; 116: 1413-1419 [PMID: 10348825 DOI: 10.1016/s0016-5085(99)70506-8]
- Kleiner DE, Brunt EM, Van Natta M, Behling C, Contos MJ, Cummings OW, Ferrell LD, Liu YC, Torbenson MS, 27 Unalp-Arida A, Yeh M, McCullough AJ, Sanyal AJ; Nonalcoholic Steatohepatitis Clinical Research Network. Design and validation of a histological scoring system for nonalcoholic fatty liver disease. Hepatology 2005; 41: 1313-1321 [PMID: 15915461 DOI: 10.1002/hep.20701]
- 28 Tamano M, Kojima K, Akima T, Murohisa T, Hashimoto T, Uetake C, Sugaya T, Nakano M, Hiraishi H, Yoneda M. The usefulness of measuring liver stiffness by transient elastography for assessing hepatic fibrosis in patients with various chronic liver diseases. Hepatogastroenterology 2012; 59: 826-830 [PMID: 22469726 DOI: 10.5754/hge11255]
- Yoneda M, Yoneda M, Fujita K, Inamori M, Tamano M, Hiriishi H, Nakajima A. Transient elastography in patients with 29 non-alcoholic fatty liver disease (NAFLD). Gut 2007; 56: 1330-1331 [PMID: 17470477 DOI: 10.1136/gut.2007.126417]
- Fang C, Konstantatou E, Romanos O, Yusuf GT, Quinlan DJ, Sidhu PS. Reproducibility of 2-Dimensional Shear Wave 30 Elastography Assessment of the Liver: A Direct Comparison With Point Shear Wave Elastography in Healthy Volunteers. J Ultrasound Med 2017; 36: 1563-1569 [PMID: 28370146 DOI: 10.7863/ultra.16.07018]
- Suda T, Kanefuji T, Abe A, Nagayama I, Hoshi T, Morita S, Yagi K, Hatakeyama S, Hayatsu M, Hasegawa N, Terai S. A 31 cut-off value of shear wave speed to distinguish nonalcoholic steatohepatitis candidates. Medicine (Baltimore) 2019; 98: e13958 [PMID: 30633176 DOI: 10.1097/MD.00000000013958]
- Sporea I, Bâldea V, Lupușoru R, Bende F, Mare R, Lazăr A, Popescu A, Șirli R. Quantification of Steatosis and Fibrosis 32 using a new system implemented in an ultrasound machine. Med Ultrason 2020; 22: 265-271 [PMID: 32399537 DOI: 10.11152/mu-2495
- 33 Hsu PK, Wu LS, Yen HH, Huang HP, Chen YY, Su PY, Su WW. Attenuation Imaging with Ultrasound as a Novel Evaluation Method for Liver Steatosis. J Clin Med 2021; 10 [PMID: 33801163 DOI: 10.3390/jcm10050965]
- 34 Jang JK, Lee ES, Seo JW, Kim YR, Kim SY, Cho YY, Lee DH. Two-dimensional Shear-Wave Elastography and US Attenuation Imaging for Nonalcoholic Steatohepatitis Diagnosis: A Cross-sectional, Multicenter Study. Radiology 2022; 305: 118-126 [PMID: 35727151 DOI: 10.1148/radiol.220220]
- Tada T, Iijima H, Kobayashi N, Yoshida M, Nishimura T, Kumada T, Kondo R, Yano H, Kage M, Nakano C, Aoki T, 35 Aizawa N, Ikeda N, Takashima T, Yuri Y, Ishii N, Hasegawa K, Takata R, Yoh K, Sakai Y, Nishikawa H, Iwata Y, Enomoto H, Hirota S, Fujimoto J, Nishiguchi S. Usefulness of Attenuation Imaging with an Ultrasound Scanner for the Evaluation of Hepatic Steatosis. Ultrasound Med Biol 2019; 45: 2679-2687 [PMID: 31277922 DOI: 10.1016/j.ultrasmedbio.2019.05.033]
- 36 Koda M, Kawakami M, Murawaki Y, Senda M. The impact of visceral fat in nonalcoholic fatty liver disease: cross-



sectional and longitudinal studies. J Gastroenterol 2007; 42: 897-903 [PMID: 18008034 DOI: 10.1007/s00535-007-2107-z]

- WHO Expert Consultation. Appropriate body-mass index for Asian populations and its implications for policy and 37 intervention strategies. Lancet 2004; 363: 157-163 [PMID: 14726171 DOI: 10.1016/S0140-6736(03)15268-3]
- Sumida Y, Yoneda M, Hyogo H, Itoh Y, Ono M, Fujii H, Eguchi Y, Suzuki Y, Aoki N, Kanemasa K, Fujita K, Chayama 38 K, Saibara T, Kawada N, Fujimoto K, Kohgo Y, Yoshikawa T, Okanoue T; Japan Study Group of Nonalcoholic Fatty Liver Disease (JSG-NAFLD). Validation of the FIB4 index in a Japanese nonalcoholic fatty liver disease population. BMC Gastroenterol 2012; 12: 2 [PMID: 22221544 DOI: 10.1186/1471-230X-12-2]
- 39 Fan JG, Kim SU, Wong VW. New trends on obesity and NAFLD in Asia. J Hepatol 2017; 67: 862-873 [PMID: 28642059 DOI: 10.1016/j.jhep.2017.06.003]
- 40 Wei JL, Leung JC, Loong TC, Wong GL, Yeung DK, Chan RS, Chan HL, Chim AM, Woo J, Chu WC, Wong VW. Prevalence and Severity of Nonalcoholic Fatty Liver Disease in Non-Obese Patients: A Population Study Using Proton-Magnetic Resonance Spectroscopy. Am J Gastroenterol 2015; 110: 1306-14; quiz 1315 [PMID: 26215532 DOI: 10.1038/ajg.2015.235]
- El-Serag HB, Tran T, Everhart JE. Diabetes increases the risk of chronic liver disease and hepatocellular carcinoma. 41 Gastroenterology 2004; 126: 460-468 [PMID: 14762783 DOI: 10.1053/j.gastro.2003.10.065]
- Tokushige K, Ikejima K, Ono M, Eguchi Y, Kamada Y, Itoh Y, Akuta N, Yoneda M, Iwasa M, Otsuka M, Tamaki N, 42 Kogiso T, Miwa H, Chayama K, Enomoto N, Shimosegawa T, Takehara T, Koike K. Evidence-based clinical practice guidelines for nonalcoholic fatty liver disease/nonalcoholic steatohepatitis 2020. J Gastroenterol 2021; 56: 951-963 [PMID: 34533632 DOI: 10.1007/s00535-021-01796-x]
- Tokushige K, Ikejima K, Ono M, Eguchi Y, Kamada Y, Itoh Y, Akuta N, Yoneda M, Iwasa M, Otsuka M, Tamaki N, 43 Kogiso T, Miwa H, Chayama K, Enomoto N, Shimosegawa T, Takehara T, Koike K. Evidence-based clinical practice guidelines for nonalcoholic fatty liver disease/nonalcoholic steatohepatitis 2020. Hepatol Res 2021; 51: 1013-1025 [PMID: 34533266 DOI: 10.1111/hepr.13688]





Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: bpgoffice@wjgnet.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

