World Journal of *Gastrointestinal Surgery*

World J Gastrointest Surg 2024 February 27; 16(2): 260-634





Published by Baishideng Publishing Group Inc

WJGS

World Journal of Gastrointestinal Surgery

Contents

Monthly Volume 16 Number 2 February 27, 2024

EDITORIAL

- 260 Actuality and underlying mechanisms of systemic immune-inflammation index and geriatric nutritional risk index prognostic value in hepatocellular carcinoma Tchilikidi KY
- 266 Prognostic impact of preoperative nutritional and immune inflammatory parameters on liver cancer Bae SU
- 270 Don't forget emergency surgery! Lessons to learn from elective indocyanine green-guided gastrointestinal interventions

Perini D, Martellucci J

276 Mutational landscape of TP53 and CDH1 in gastric cancer Cai HQ, Zhang LY, Fu LM, Xu B, Jiao Y

284 Overview of ectopic pancreas Li CF, Li QR, Bai M, Lv YS, Jiao Y

ORIGINAL ARTICLE

Clinical and Translational Research

289 Phospholipase A2 enzymes PLA2G2A and PLA2G12B as potential diagnostic and prognostic biomarkers in cholangiocarcinoma

Qiu C, Xiang YK, Da XB, Zhang HL, Kong XY, Hou NZ, Zhang C, Tian FZ, Yang YL

Case Control Study

307 Classification of anatomical morphology of cystic duct and its association with gallstone Zhu JH, Zhao SL, Kang Q, Zhu Y, Liu LX, Zou H

Retrospective Cohort Study

- 318 Will partial splenic embolization followed by splenectomy increase intraoperative bleeding? Huang L, Li QL, Yu QS, Peng H, Zhen Z, Shen Y, Zhang Q
- 331 Influence of donor age on liver transplantation outcomes: A multivariate analysis and comparative study Bezjak M, Stresec I, Kocman B, Jadrijević S, Filipec Kanizaj T, Antonijević M, Dalbelo Bašić B, Mikulić D
- 345 Machine learning-based radiomics score improves prognostic prediction accuracy of stage II/III gastric cancer: A multi-cohort study

Xiang YH, Mou H, Qu B, Sun HR



World Journal of Gastrointestinal Surgery				
Conte	nts Monthly Volume 16 Number 2 February 27, 2024			
357	Risk stratification in gastric cancer lung metastasis: Utilizing an overall survival nomogram and comparing it with previous staging			
	Chen ZR, Yang MF, Xie ZY, Wang PA, Zhang L, Huang ZH, Luo Y			
382	Systemic inflammatory response index is a predictor of prognosis in gastric cancer patients: Retrospective cohort and meta-analysis			
	Ren JY, Xu M, Niu XD, Ma SX, Jiao YJ, Wang D, Yu M, Cai H			
	Retrospective Study			
396	Development of a clinical nomogram for prediction of response to neoadjuvant chemotherapy in patients with advanced gastric cancer			
	Liu B, Xu YJ, Chu FR, Sun G, Zhao GD, Wang SZ			
409	Laparoscopic left hemihepatectomy guided by indocyanine green fluorescence: A cranial-dorsal approach			
	Wang XR, Li XJ, Wan DD, Zhang Q, Liu TX, Shen ZW, Tong HX, Li Y, Li JW			
419	Hemoglobin loss method calculates blood loss during pancreaticoduodenectomy and predicts bleeding- related risk factors			
	Yu C, Lin YM, Xian GZ			
429	Short- and long-term outcomes of surgical treatment in patients with intestinal Behcet's disease			
	Park MY, Yoon YS, Park JH, Lee JL, Yu CS			
438	Preoperative neutrophil-to-lymphocyte ratio predicts symptomatic anastomotic leakage in elderly colon cancer patients: Multicenter propensity score-matched analysis			
	Wang CY, Li XL, Ma XL, Yang XF, Liu YY, Yu YJ			
451	Preoperative blood markers and intra-abdominal infection after colorectal cancer resection			
	Liu CQ, Yu ZB, Gan JX, Mei TM			
463	Immune function status of postoperative patients with colon cancer for predicting liver metastasis			
	Xiong L, Liu FC			
471	Efficacy of transjugular intrahepatic portosystemic shunts in treating cirrhotic esophageal-gastric variceal bleeding			
	Hu XG, Dai JJ, Lu J, Li G, Wang JM, Deng Y, Feng R, Lu KP			
481	Correlation between serum markers and transjugular intrahepatic portosystemic shunt prognosis in patients with cirrhotic ascites			
	Hu XG, Yang XX, Lu J, Li G, Dai JJ, Wang JM, Deng Y, Feng R			
491	Development of a new Cox model for predicting long-term survival in hepatitis cirrhosis patients underwent transjugular intrahepatic portosystemic shunts			
	Lv YF, Zhu B, Meng MM, Wu YF, Dong CB, Zhang Y, Liu BW, You SL, Lv S, Yang YP, Liu FQ			
503	"Five steps four quadrants" modularized <i>en bloc</i> dissection technique for accessing hepatic hilum lymph nodes in laparoscopic pancreaticoduodenectomy			
	Hu XS, Wang Y, Pan HT, Zhu C, Chen SL, Liu HC, Pang Q, Jin H			



	World Journal of Gastrointestinal Surgery
Conte	nts
	Monthly Volume 16 Number 2 February 27, 2024
511	Efficacy and safety of endoscopic submucosal dissection for early gastric cancer and precancerous lesions in elderly patients
	Xu WS, Zhang HY, Jin S, Zhang Q, Liu HD, Wang MT, Zhang B
518	Nomogram model including <i>LATS2</i> expression was constructed to predict the prognosis of advanced gastric cancer after surgery
	Sun N, Tan BB, Li Y
	Observational Study
529	To explore the pathogenesis of anterior resection syndrome by magnetic resonance imaging rectal defeco- graphy
	Meng LH, Mo XW, Yang BY, Qin HQ, Song QZ, He XX, Li Q, Wang Z, Mo CL, Yang GH
539	Biopsy forceps are useful for measuring esophageal varices in vitro
	Duan ZH, Zhou SY
	SYSTEMATIC REVIEWS
546	First experience in laparoscopic surgery in low and middle income countries: A systematic review
	Troller R, Bawa J, Baker O, Ashcroft J
554	Comparative effectiveness of several adjuvant therapies after hepatectomy for hepatocellular carcinoma patients with microvascular invasion
	Pei YX, Su CG, Liao Z, Li WW, Wang ZX, Liu JL
	META-ANALYSIS
571	Is tumor necrosis factor-α monoclonal therapy with proactive therapeutic drug monitoring optimized for
571	inflammatory bowel disease? Network meta-analysis
	Zheng FY, Yang KS, Min WC, Li XZ, Xing Y, Wang S, Zhang YS, Zhao QC
585	Poor oral health was associated with higher risk of gastric cancer: Evidence from 1431677 participants
	Liu F, Tang SJ, Li ZW, Liu XR, Lv Q, Zhang W, Peng D
	CASE REPORT
596	Treatment of hemolymphangioma by robotic surgery: A case report
	Li TN, Liu YH, Zhao J, Mu H, Cao L
601	Postoperative encapsulated hemoperitoneum in a patient with gastric stromal tumor treated by exposed endoscopic full-thickness resection: A case report
	Lu HF, Li JJ, Zhu DB, Mao LQ, Xu LF, Yu J, Yao LH
609	Early endoscopic management of an infected acute necrotic collection misdiagnosed as a pancreatic pseudocyst: A case report
	Zhang HY, He CC



Conto	World Journal of Gastrointestinal Surgery
Conte	Monthly Volume 16 Number 2 February 27, 2024
616	Percutaneous ultrasound-guided coaxial core needle biopsy for the diagnosis of multiple splenic lesions: A case report
	Pu SH, Bao WYG, Jiang ZP, Yang R, Lu Q
622	Spilled gallstone mimicking intra-abdominal seeding of gallbladder adenocarcinoma: A case report
	Huang CK, Lu RH, Chen CC, Chen PC, Hsu WC, Tsai MJ, Ting CT
628	Ileal collision tumor associated with gastrointestinal bleeding: A case report and review of literature
	Wu YQ, Wang HY, Shao MM, Xu L, Jiang XY, Guo SJ



Contents

Monthly Volume 16 Number 2 February 27, 2024

ABOUT COVER

Editorial Board Member of World Journal of Gastrointestinal Surgery, Nikolaos Chatzizacharias, FACS, FRCS, MD, PhD, Consultant Surgeon, Department of HPB and liver transplantation, Queen Elizabeth Hospital, University Hospitals Birmingham, Birmingham B15 2TH, United Kingdom. nikolaos.chatzizacharias@uhb.nhs.uk

AIMS AND SCOPE

The primary aim of World Journal of Gastrointestinal Surgery (WJGS, World J Gastrointest Surg) is to provide scholars and readers from various fields of gastrointestinal surgery with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGS mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal surgery and covering a wide range of topics including biliary tract surgical procedures, biliopancreatic diversion, colectomy, esophagectomy, esophagostomy, pancreas transplantation, and pancreatectomy, etc.

INDEXING/ABSTRACTING

The WJGS is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports[®] cites the 2022 impact factor (IF) for WJGS as 2.0; IF without journal self cites: 1.9; 5-year IF: 2.2; Journal Citation Indicator: 0.52; Ranking: 113 among 212 journals in surgery; Quartile category: Q3; Ranking: 81 among 93 journals in gastroenterology and hepatology; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Zi-Hang Xu; Production Department Director: Xiang Li; Editorial Office Director: Jia-Ru Fan.

NAME OF JOURNAL	INSTRUCTIONS TO AUTHORS
World Journal of Gastrointestinal Surgery	https://www.wjgnet.com/bpg/gerinfo/204
ISSN	GUIDELINES FOR ETHICS DOCUMENTS
ISSN 1948-9366 (online)	https://www.wjgnet.com/bpg/GerInfo/287
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH
November 30, 2009	https://www.wjgnet.com/bpg/gerinfo/240
FREQUENCY	PUBLICATION ETHICS
Monthly	https://www.wjgnet.com/bpg/GerInfo/288
EDITORS-IN-CHIEF	PUBLICATION MISCONDUCT
Peter Schemmer	https://www.wjgnet.com/bpg/gerinfo/208
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE
https://www.wjgnet.com/1948-9366/editorialboard.htm	https://www.wignet.com/bpg/gerinfo/242
PUBLICATION DATE	STEPS FOR SUBMITTING MANUSCRIPTS
February 27, 2024	https://www.wjgnet.com/bpg/GerInfo/239
COPYRIGHT	ONLINE SUBMISSION
© 2024 Baishideng Publishing Group Inc	https://www.f6publishing.com

© 2024 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: office@baishideng.com https://www.wjgnet.com



Х

S WU



Submit a Manuscript: https://www.f6publishing.com

World J Gastrointest Surg 2024 February 27; 16(2): 546-553

DOI: 10.4240/wjgs.v16.i2.546

ISSN 1948-9366 (online)

SYSTEMATIC REVIEWS

First experience in laparoscopic surgery in low and middle income countries: A systematic review

Rebekka Troller, Jasmine Bawa, Olivia Baker, James Ashcroft

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0 Grade B (Very good): 0 Grade C (Good): C Grade D (Fair): 0 Grade E (Poor): 0

P-Reviewer: Navarrete Arellano M, Mexico

Received: November 6, 2023 Peer-review started: November 6, 2023 First decision: December 6, 2023 Revised: January 2, 2024 Accepted: January 30, 2024 Article in press: January 30, 2024 Published online: February 27, 2024



Rebekka Troller, Department of Surgery, Medway Maritime Hospital, Gillingham ME7 5NY, Kent, United Kingdom

Jasmine Bawa, Olivia Baker, James Ashcroft, Department of Surgery, University Hospital Cambridge, Cambridge CB2 0QQ, United Kingdom

Corresponding author: Rebekka Troller, MD, Doctor, Surgeon, Department of Surgery, Medway Maritime Hospital, Windmill Rd, Gillingham ME7 5NY, Kent, United Kingdom. rebekka.troller@gmail.com

Abstract

BACKGROUND

Laparoscopic surgery has reduced morbidity and mortality rates, shorter postoperative recovery periods and lower complication rates than open surgery. It is routine practice in high-income countries and is becoming increasingly common in countries with limited resources. However, introducing laparoscopic surgery in low-and-middle-income countries (LMIC) can be expensive and requires resources, equipment, and trainers.

AIM

To report the challenges and benefits of introducing laparoscopic surgery in LMIC as well as to identify solutions to these challenges for countries with limited finances and resources.

METHODS

MEDLINE, EMBASE and Cochrane databases were searched for studies reporting first experience in laparoscopic surgery in LMIC. Included studies were published between 1996 and 2022 with full text available in English. Exclusion criteria were studies considering only open surgery, ear, nose, and throat, endoscopy, arthroscopy, hysteroscopy, cystoscopy, transplant, or bariatric surgery.

RESULTS

Ten studies out of 3409 screened papers, from eight LMIC were eligible for inclusion in the final analysis, totaling 2497 patients. Most reported challenges were related to costs of equipment and training programmes, equipment problems such as faulty equipment, and access to surgical kits. Training-related challenges were reliance on foreign trainers and lack of locally trained surgeons and theatre staff. The benefits of introducing laparoscopic surgery were economic and clinical, including a reduction in hospital stay, complications, and morbidi-



WJGS https://www.wjgnet.com

ty/mortality. The introduction of laparoscopic surgery also provided training opportunities for junior doctors.

CONCLUSION

Despite financial and technical challenges, many studies emphasise the overall benefit of introducing laparoscopic surgery in LMICs such as reduced hospital stay and the related lower cost for patients. While many of the clinical centres in LMICs have proposed practical solutions to the challenges reported, more support is critically required, in particular regarding training.

Key Words: Laparoscopic surgery; Low and middle income country; First experience; Training in laparoscopic surgery

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: While laparoscopic surgery could have major positive effects on patients in low and middle income countries, this type of operation is not widely accessible in many parts of the world. The aim of this study was a systematic review of the literature on laparoscopic surgery in low in middle income countries in order to get a better understanding of first experiences made with minimal invasive surgical techniques and equipment. Out of over 3400 screened papers, 10 studies were selected and analised. We report on common challenges as well as on possible ways to overcome them.

Citation: Troller R, Bawa J, Baker O, Ashcroft J. First experience in laparoscopic surgery in low and middle income countries: A systematic review. World J Gastrointest Surg 2024; 16(2): 546-553 URL: https://www.wjgnet.com/1948-9366/full/v16/i2/546.htm DOI: https://dx.doi.org/10.4240/wjgs.v16.i2.546

INTRODUCTION

Laparoscopic surgery has become routine practice in many places in the world and has numerous benefits when compared to open surgery, including: Reduced morbidity and mortality, length of hospital stay, blood loss, infection rate and postoperative pain[1-3]. However, laparoscopic surgery is still not routine in many low- and middle income countries (LMICs).

The term LMIC includes low, lower-middle, and upper-middle income countries, defined by their gross national income (GNI). The GNI fluctuates every year, and as of 2022 a LMIC is defined by a GNI per capita of less than \$12695[4] (Unless otherwise stated, all monetary values in this paper are in American Dollars: \$). This economic deficit can limit access to high-cost technology and its support and represents a great challenge to introduce laparoscopic surgery.

LMICs perform a greater number of procedures with an open approach, rather than a laparoscopic approach. Access to laparoscopic surgery in LMICs is difficult and thus much of the developing world cannot benefit from minimal invasive surgery. Introducing laparoscopic surgical services requires a range of specialist input necessitating that surgical teams are constituted from a broad multidisciplinary background.

This systematic review aims to assess the challenges in introducing laparoscopic surgery service in LMICs and to produce a consensus regarding commencing a high quality, replicable, and collaborative laparoscopic service.

MATERIALS AND METHODS

Search strategy

This Systematic Review utilised the preferred reporting items for systematic reviews and meta-analyses statement (PRISMA Statement) to guide all methodology and write up. Before undertaking this review, inclusion criteria were predetermined. Reports met inclusion criteria if they described: (1) A report of first experience in laparoscopic surgery; (2) performed in a low-and-middle income country; (3) were reported in the English language or had an English language translation; and (4) were published after 1996.

Exclusion criteria defined were as follows: (1) Studies focused on open surgery, endoscopy, arthroscopy, cystoscopy, hysteroscopy, hand-assisted surgery, ear, nose, and throat surgery, bariatric surgery, breast surgery, thoracic surgery, robotic surgery, transplant surgery, obstetric; (2) reviews, case reports, letters to the editor or conference abstracts; (3) studies where the patient cohort was equal or smaller than ten; and (4) studies which reported advanced laparoscopic techniques. Studies for which the online full-text was not available were also excluded.

A systematic search of MEDLINE, EMBASE and Cochrane was undertaken until November 2022. An initial search for the identification of laparoscopic surgical reports in LMIC utilised the search terms: "laparoscopic", "laparoscopy", and "hand-assisted laparoscopy" in addition to the terms "LMIC" or "developed/developing country". This broad search strategy generated 4026 articles.

WJGS | https://www.wjgnet.com

Study selection, data extraction, and synthesis of results

Duplicates were removed (n = 617) and 4 reviewers independently screened abstracts and/or full texts of 3409 publications in reference to study design, study population, and study location as outlined in the inclusion and exclusion criteria. Full screening resulted in the capture of 10 articles which met the predetermined criteria for inclusion into the systematic review. This process is displayed in full in Figure 1. Prior to data extraction it was decided that a meta-analysis would not be undertaken due to the heterogeneity of both laparoscopic implementations and outcome measurements.

Three reviewers independently extracted relevant information from each laparoscopic report using a standardised data extraction method. One author with surgical training expertise reviewed all extracted data. This review captured: Cost, equipment, training, clinical-, economic- and systemic details in addition to quantitative and qualitative outcome measures.

RESULTS

The process of identification, detailing the search and screening strategy is shown in the PRISMA flow diagram (Figure 1).

Characteristics of included studies

Ten studies were identified as meeting the inclusion criteria. The majority of these studies were conducted in Africa (n = 9) with four studies being conducted in Nigeria (Table 1)[5-14]. The studies were conducted between 1999 and 2018 and had a study duration of 2.8 years on average. Half were retrospective studies (n = 5), the remainder were prospective (n = 4) and one case series (n = 1). A total of 2497 patients were eligible for inclusion in the final analysis. No multinational studies were identified for inclusion during the screening process.

Challenges

There were three common themes for the challenges faced when introducing laparoscopic surgery; these related to training, costs and equipment (Table 2)[5-14].

Training

Minimal access surgery requires additional training and skills and this can be challenging in many LMICs[11,12]. As laparoscopic surgery is less often practised, the result is that junior surgeons have poor knowledge of laparoscopy and lack the confidence to use laparoscopy outside of normal working hours, even for emergency cases, such as for appendicitis[7]. Once more surgeons develop an interest and improve skills in laparoscopic surgery, the operating time will reduce[6].

In a study defining the learning curve for laparoscopic colorectal resections in the United States, it was felt that a minimum of 30 procedures was required for laparoscopic colorectal resection in an expert unit, with early cases being associated with higher conversion to open, operative time, and post-operative stay[15].

Apart from the additional training for surgeons, many centres found it challenging to introduce laparoscopic surgery because they did not have adequately trained staff such as theatre nurses, technicians and scrub nurses[6,7,9,11]. As laparoscopic surgery is not widely practised in LMICs, it may be that these skills are not taught to trainees[16] and in turn there is no motivation for training support staff in these skills.

Some studies reported on the training of local staff either abroad or in their hospital by surgeons from abroad. Training was provided not only for the surgeons, but also the theatre staff, nurses and paramedics[9]. A gynaecology team from Cameroon were trained over seven years by a surgeon from France[8]. The anaesthetists and theatre staff of the same team also benefited from several months of training[8]. In some cases, it falls on the operating surgeon to train the entire team (endoscopy nurses and technicians) as there is inadequate training of support staff[11].

Some centres are struggling with a complete absence of basic training in laparoscopic surgery resulting in underperformance and a very slow learning curve[12]. Once more surgeons develop an interest and improve their skills in laparoscopic surgery, a reduction in operating time can be achieved[6]. This is reflected in the units that were only able to perform laparoscopic procedures intermittently, yet still reported reduced operation times and improvement of surgical skills[9,12].

Training is not only an issue for clinical teams, but also for the engineering teams that maintain the laparoscopic equipment. Local biometric engineers are often not knowledgeable about the endoscopic instruments or trained to fix them[12,14]. Maintenance of laparoscopic equipment by untrained staff can lead to malfunction and contribute to equipment problems encountered during surgery[17]. To reduce cost, reusable instruments are used and therefore careful sterilisation is necessary which requires advanced know-how[14].

Costs

While the long term costs of laparoscopic surgery are comparable or even cheaper than the ones of open surgery, the initial investment for equipment, training and infrastructure can pose a problem. In one of the first studies introducing laparoscopic appendicectomy in the United States, it was found that laparoscopic surgery required a significant increase in intraoperative equipment charge ($$125.32 \pm 3.99 open *vs* $$1078.70 \pm 24.06 Lap), operative time charge ($$3022.16 \pm 57.51 *vs* $$4065.24 \pm 122.64), and total hospital charge ($$12310 \pm 772 *vs* $$16773 \pm 1319)[18].

Zaishideng® WJGS | https://www.wjgnet.com

Table 1 Characteristics of included studies						
Ref.	Country	Year published	Study duration	Specialty	Study design	No. cases
Bendinelli et al[5]	Senegal	2002	5 years	Various	Retrospective	826
Parkar et al[6]	Kenya	2003	2 years	Gynae and general	Retrospective	408
Patel <i>et al</i> [7]	Kenya	2003	6 years	General	Case series	106
Raiga et al[<mark>8</mark>]	Cameroon	1999	4 years	Gynae	Retrospective	735
Arung et al[9]	Democratic Republic of Congo	2015	10 months	General	Prospective	75
Adisa et al[10]	Nigeria	2013	2 years	General	Retrospective	175
Ray-Offor <i>et al</i> [11]	Nigeria	2014	17 months	General and paediatric	Prospective	15
Onoh et al[12]	Nigeria	2018	2 years	Gynae	Prospective	36
Ishaq et al[<mark>13</mark>]	Pakistan	2016	31 months	Paediatric	Retrospective	100
Ismaila et al[14]	Nigeria	2013	12 months	General	Prospective	21

Table 2 Commonly reported challenges for the introduction of laparoscopic surgery

Ref.	Training	Costs	Equipment	Others
Bendinelli et al[5]		Y	Y	
Parkar <i>et al</i> [6]	Y		Υ	
Patel et al[7]	Υ			Υ
Raiga et al[8]	Υ	Υ		
Arung et al[9]	Y	Y	Υ	
Adisa <i>et al</i> [10]		Υ	Y	
Ray-Offor <i>et al</i> [11]	Y	Y	Υ	Υ
Onoh <i>et al</i> [12]	Y	Y	Υ	Υ
Ishaq et al[13]	Y			
Ismaila et al[14]	Υ		Y	Y

Y: Yes. Challenges were reported in the literature.

Six studies report the cost to acquire laparoscopic equipment and consumables in LMICs as an issue to establish a laparoscopic surgery service in their respective hospitals. The main challenge in LMICs is the high initial costs to acquire laparoscopic equipment and consumables^[9]. This is in part due to the absence of local production companies resulting in a lack of supply of instruments on demand and in part due to fewer alternative options being available to purchase affordable equipment[12].

Regarding the operational costs, Ray-Offor et al[11] reported that after the initial setup, which includes training of personnel, acquisition of equipment and building the relevant infrastructure, the operational costs for laparoscopic surgery is not huge (though without exact specification or comparison to open surgery)[11]. Similarly, Bendinelli et al[5] reported that the hospital could recover more than 87% of its investment in 6 years and the higher surgical charges were compensated by reduced hospitalisation costs postoperatively[5].

Adisa et al[10] and other centres described how they were able to reduce costs with local adaptations like investing in reusable instruments, which helped to reduce the cost per patient[10,11]. They also made camera covers using drapes that can be washed and sterilised and limited the use of retrieval bags for purulent appendix specimens. No advanced energy devices such as harmonic or Ligasure were used[10].

Depending on the pricing model of the hospital laparoscopic surgery can even be more financially viable than open surgery. For example, Bendinelli et al[5] report that in hospitals with fixed charges per surgery and per day, the costs for patients in the laparoscopic surgery group were lower when compared to the open surgery group, due to shorter length of hospital stay and fewer complications[5].

Another barrier to introducing laparoscopic surgery in LMICs is the lack of a well established healthcare system with insurance where patients must pay for the surgery and hospital stay themselves [10,12]. In these settings, hospitals often operate with extremely low margins and generate very little profit. As a result, for a long time, laparoscopic surgery in LMICs was only offered in private hospitals who historically have been more likely to be able to afford the higher upfront investment cost required for modern equipment and training[9,10].

WJGS | https://www.wjgnet.com

Troller R et al. Systematic review: Laparoscopic surgery in LMICs




Figure 1 Study identification process.

One of the overall benefits of laparoscopic surgery compared to open surgery is the reduced hospital stay which means lower costs for patients in LMICs[5,7,10,12]. Diagnostic laparoscopy is one of the most convenient procedures; it is simple, cost-effective and avoids an unnecessary laparotomy which leads to a shorter postoperative stay and lower costs[5,9,12].

Equipment

Equipment challenges evolve mostly around the acquisition of equipment and the reliability of equipment and infrastructure [5,6]. Most of the hospitals received donations of endoscopic instruments from corporations, companies or visiting surgeons [5,8,12,14]. Relying on such donations and investing in reusable equipment were two methods to achieve cost effectiveness.

Some studies noted that the costs for laparoscopic surgery are not much higher than conventional surgery after the initial set-up, training of staff and acquisition of relevant infrastructure[11,19]. For the initial set-up, cooperation with developed countries and public institutions can be helpful to obtain laparoscopic supplies [5]. However, the lack of local companies producing laparoscopic equipment can lead to maintenance problems. Maintenance of laparoscopic equipment by untrained staff contributes to some problems encountered during surgery as well as challenges with effective sterilisation and care of the instruments^[14].

Another commonly reported challenge is unreliable infrastructure which can lead to power outages or unreliable power supplies[11,12,14], which creates a huge problem as laparoscopic surgery requires a reliable source of electricity. Introduction of solar power or electricity by a generator or an alternate power supply during surgeries could provide a solution to the power outages[11].

Adisa et al[10] introduced laparoscopic surgery with a basic setup using regular TV manual carbon dioxide insufflations. They reported four conversions due to equipment failure and technical difficulties[10]. In two cases loss of view due to a faulty cable that could not be immediately resolved and two conversions happened due to excessive intraoperative bleeding. This study highlighted how the costs and difficulties of replacing damaged laparoscopic equipment can be a limiting factor to continuing minimally invasive surgery[12].

Specific issues with equipment such as diathermy malfunctioning, faulty suction, gas leak from a malfunctioning port, and a non-functioning clip applicator were reported by Ismaila *et al*[14].

Other challenges and outcomes

One study reported lack of acceptance of laparoscopic surgery being a challenge; surgeons had a preference for nonendoscopic techniques and were not keen on trying a new technique[12]. In addition, the initial set-up for laparoscopic surgery can be time-consuming, which can prolong operating time and cause problems in view of competing theatre usage[11].

Bureaucratic issues regarding procurement and approving endoscopic equipment can be another challenge in introducing minimally invasive surgery [12,14].

Clinical outcomes

Most authors looked at complication and conversion rates. Complication rate is known to be related to the level of training and experience in laparoscopic surgery^[20]. As Parkar et al^[6] pointed out, laparoscopic training and collaboration with local trainers or those from overseas is crucial for the development of minimal invasive surgery [5,12].

An increased conversion rate related to inadequate equipment, malfunction and poor patient selection was reported by Onoh et al[12]. One hospital reported a conversion rate of 38% (16 conversions of 42 Laparoscopic appendicectomy), most of them due to intraoperative difficulties such as retrocaecal appendix, necrosed fragmented appendix or adhesions[7].



That being said, most authors reported a conversion rate comparable to the number in developed countries^[21] between 1 and 10%^[5,6].

One study reported six port site complications (1.47%) of which two needed subsequent laparotomy only seven days post laparoscopic surgery[6]. This is comparable to the other studies reporting port site complications between 3% and 9% within three months after surgery[22,23].

A number of studies highlighted the reduction in operating time with experience[6,9,10] as well as shorter postoperative stay and less pain after laparoscopic surgery[5,9,10].

Diagnostic laparoscopy is one of the most convenient procedures and can avoid an unnecessary laparotomy which leads to reduced wound infection rates [7,9,10], minimises scar formation [7], allows earlier return to normal activities [7, 12], less pain [9] and reduces postoperative adhesions [7,8]. The routine use of diagnostic laparoscopy can also reduce morbidity and mortality for patients with suspected intra-abdominal malignancies compared to diagnostic laparotomies for these patients [10].

DISCUSSION

Laparoscopic surgery presents significant opportunities for enhancing healthcare delivery in low and middle-income countries (LMICs). Its advantages, including shorter hospital stays, improved patient outcomes due to lower infection rates and fewer laparotomies which enable a quicker return to work, highlight the potentially transformative impact on healthcare systems in resource-limited settings.

While these opportunities are promising, numerous challenges must be addressed to fully realise the benefits of laparoscopic surgery in LMICs.

Costs for equipment, training and infrastructure are a major barrier due to low income and the associated low economic development in LMICs. The economic difficulties that face many LMICs may have been a result of conflicts, inequality, lack of education and poor basic infrastructure. Therefore, support of LMICs in introducing laparoscopic surgery and training needs to be built and adapted to local practice to be sustainable.

One of the primary hurdles is the substantial initial investment required for acquiring and maintaining laparoscopic equipment. The higher acquisition and maintenance costs, and limited financial resources in LMICs, pose a significant barrier to widespread adoption. Moreover, the absence of skilled local labour for equipment maintenance aggravate these challenges, leading to potential equipment downtime and reduced accessibility to laparoscopic equipment.

The absence of a local support industry further complicates the sustainability of laparoscopic surgery in LMICs. Inadequate infrastructure, including unstable power and water supply, presents operational difficulties and jeopardises the continuity of surgical procedures. The need for larger sterilisation equipment adds to the infrastructure demands, requiring substantial modifications that may not align with existing healthcare facilities in these regions.

A critical challenge remains the scarcity of trained surgeons in laparoscopic surgery in LMICs associated with a lack of local surgical training opportunities and educational resources. Efforts to establish comprehensive training programs tailored to the specific needs of these regions are imperative to overcome this limitation. Collaborations between local and international medical institutions could facilitate knowledge transfer and skill development, ultimately empowering local healthcare professionals. Previous reports have documented successful introduction of a laparoscopic surgical service, such as the Ministry of Health in Botswana where 12 surgeons were trained through a combination of hands-on teaching, tele-simulation, workshops and mentorship to begin a laparoscopic cholecystectomy program[24].

Strategies to address these challenges should focus not only on infrastructure shortcomings but also establishing sustainable training programs and support systems for equipment maintenance. Innovative financing models and technology-sharing initiatives could alleviate the financial burden associated with the adoption and maintenance of laparoscopic surgery in LMICs.

CONCLUSION

On the whole, laparoscopic surgery could have a huge potential in LMICs. Collaborative efforts between international surgeons, companies, and local units emerge as a cornerstone for success in navigating the challenges.

Identification of essential equipment needs and skill gaps, coupled with comprehensive training programs for both clinical staff and equipment engineers, are needed for a sustainable implementation. Reusable equipment not only addresses financial constraints but also aligns with the principles of resource efficiency which is crucial in resource-limited settings. Given the lack of local resources and skills, the use of remote technology, such as video-based training materials or remote video-assistance to train and mentor local teams could also be considered.

The introduction of solar electricity or alternative power supplies could enable better coping with disruptions posed by unreliable power supply, ensuring uninterrupted surgical procedures.

Further studies around the cost-effectiveness of laparoscopic surgery in LMICs are required, to evaluate approaches that might enable LMICs to benefit from minimal invasive surgery. Lastly, a future survey which also includes studies from less economically prosperous areas within otherwise more developed countries, might help yield a larger number of studies and hence increase the validity of the review.

Zaishidena® WJGS | https://www.wjgnet.com

ARTICLE HIGHLIGHTS

Research background

Laparoscopic surgery has become routine practice in many places in the world and has numerous benefits when compared to open surgery, including: Reduced morbidity and mortality, length of hospital stay, blood loss, infection rate and postoperative pain. However, laparoscopic surgery is still not routine in many low- and middle income countries (LMIC) as introducing minimal invasive surgery can be expensive and requires resources. The economic deficit in lowand middle income countries can limit access to high-cost technology and its support and represents a great challenge to introduce laparoscopic surgery. In low-and middle income countries most surgeries are performed with an open approach, rather than a laparoscopically. Access to laparoscopic surgery in LMICs is difficult and thus much of the developing world cannot benefit from minimal invasive surgery. Introducing laparoscopic surgical services requires a range of specialist input necessitating that surgical teams are constituted from a broad multidisciplinary background.

Research motivation

This systematic review aims to assess the challenges in introducing laparoscopic surgery service in LMICs and to produce a consensus regarding commencing a high quality, replicable, and collaborative laparoscopic service.

Research objectives

The main objective was to assess the challenges in surgical services in Low- and middle-income countries. We could identify the main gaps and problems and analysed potential solutions. But also identified the need for future studies addressing certain questions and design.

Research methods

Systematic review of MEDLINE, EMBASE and Cochrane databases.

Research results

Ten studies have been included in this systematic review. They main challenges found were related to costs, training and equipment. Lack of training opportunities, faulty equipment or access to equipment and the associated costs. Some studies highlighted the benefits of introducing laparoscopic surgery, particularly the reduction in hospital stay, lower complications and lower morbidity and mortality.

Research conclusions

Identification of essential equipment needs and skill gaps, coupled with comprehensive training programs for both clinical staff and equipment engineers, are needed for a sustainable implementation. Reusable equipment, the use of remote technology, such as video-based training materials or remote video-assistance to train and mentor local teams, introduction of solar electricity or alternative power supplies should be considered. Further studies to assess cost-effectiveness of laparoscopic surgery and studies which also includes papers from less economically prosperous areas within developed countries might help to increase the validity of the review.

Research perspectives

Further studies to assess cost-effectiveness of laparoscopic surgery and studies which also includes papers from less economically prosperous areas within developed countries might help to increase the validity of the review.

FOOTNOTES

Author contributions: Troller R contributed to identification and screening of papers, analysis, supervision of writing the abstract, completion of manuscript and revision; Bawa J contributed to study screening, data analysis, writing of abstract and part of manuscript; Baker O contributed to study screening, data analysis; Ashcroft J contributed to study screening, data analysis, writing part of manuscript.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

PRISMA 2009 Checklist statement: The authors have read the PRISMA 2009 Checklist, and the manuscript was prepared and revised according to the PRISMA 2009 Checklist.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country/Territory of origin: United Kingdom

ORCID number: Rebekka Troller 0000-0002-6167-3386; Jasmine Bawa 0000-0002-9332-8851; Olivia Baker 0000-0002-0038-2657; James Ashcroft



0000-0003-2964-5032.

S-Editor: Li L L-Editor: A P-Editor: Yuan YY

REFERENCES

- 1 Buia A, Stockhausen F, Hanisch E. Laparoscopic surgery: A qualified systematic review. World J Methodol 2015; 5: 238-254 [PMID: 26713285 DOI: 10.5662/wjm.v5.i4.238]
- 2 Braga M, Vignali A, Zuliani W, Frasson M, Di Serio C, Di Carlo V. Laparoscopic vs open colorectal surgery: cost-benefit analysis in a singlecenter randomized trial. Ann Surg 2005; 242: 890-895, discussion 895 [PMID: 16327499 DOI: 10.1097/01.sla.0000189573.23744.59]
- Jaschinski T, Mosch CG, Eikermann M, Neugebauer EA, Sauerland S. Laparoscopic vs open surgery for suspected appendicitis. Cochrane 3 Database Syst Rev 2018; 11: CD001546 [PMID: 30484855 DOI: 10.1002/14651858.CD001546.pub4]
- Worldbank, World Bank GNI per capita in middle income countries, 2022. Aug 29, 2022. [cited 13 January 2024]. Available from: https:// 4 www.worldbank.org/en/country/mic/overview#1
- Bendinelli C, Leal T, Moncade F, Dieng M, Toure CT, Miccoli P. Endoscopic surgery in Senegal. Benefits, costs and limits. Surg Endosc 5 2002; 16: 1488-1492 [PMID: 11988789 DOI: 10.1007/s00464-001-9188-1]
- Parkar RB, Thagana NG, Baraza R, Otieno D. Experience with laparoscopic surgery at the Aga Khan Hospital, Nairobi. East Afr Med J 2003; 6 **80**: 44-50 [PMID: 12755241 DOI: 10.4314/eamj.v80i1.8665]
- Patel SC, Jumba GF, Akmal S. Laparoscopic appendicectomy at the Aga Khan Hospital, Nairobi. East Afr Med J 2003; 80: 447-451 [PMID: 7 14640164 DOI: 10.4314/eamj.v80i9.8740]
- Raiga J, Kasia JM, Bruhat MA. Laparoscopic surgery in the Cameroon. Int J Gynaecol Obstet 1999; 65: 65-66 [PMID: 10390103 DOI: 8 10.1016/s0020-7292(98)00206-9]
- Arung W, Dinganga N, Ngoie E, Odimba E, Detry O. First steps of laparoscopic surgery in Lubumbashi: problems encountered and 9 preliminary results. Pan Afr Med J 2015; 21: 210 [PMID: 26448805 DOI: 10.11604/pamj.2015.21.210.6689]
- Adisa AO, Lawal OO, Arowolo OA, Alatise OI. Local adaptations aid establishment of laparoscopic surgery in a semiurban Nigerian hospital. 10 Surg Endosc 2013; 27: 390-393 [PMID: 22806524 DOI: 10.1007/s00464-012-2463-5]
- Ray-Offor E, Okoro P, Gbobo I, Allison A. Pilot study on laparoscopic surgery in port-harcourt, Nigeria. Niger J Surg 2014; 20: 23-25 11 [PMID: 24665198 DOI: 10.4103/1117-6806.127104]
- 12 Onoh RC, Ezeonu PO, Lawani LO, Ajah LO, Ezegwui HU, Ejikeme BN. Experiences and challenges of gynecological endoscopy in a lowresource setting, Southeast Nigeria. Trop J Obstet Gynaecol 2018; 35: 30-37 [DOI: 10.4103/TJOG.TJOG_34_17]
- Ishaq H, Qazi SH, Dogar S, Khan Durrani MY, Faruque AV. Pediatric laparoscopic surgery; initial experience from Pakistan; first 100 cases 13 in single center. J Pak Med Assoc 2016; 66: S116-S118 [PMID: 27895374]
- Ismaila BO, Shuaibu SI, Ale AA. Laparoscopic surgery in a Nigerian teaching hospital for 1 year: challenges and effect on outcomes. Niger J 14 *Med* 2013; **22**: 134-137 [PMID: 23829125]
- Schlachta CM, Mamazza J, Seshadri PA, Cadeddu M, Gregoire R, Poulin EC. Defining a learning curve for laparoscopic colorectal resections. 15 Dis Colon Rectum 2001; 44: 217-222 [PMID: 11227938 DOI: 10.1007/BF02234296]
- Wilkinson E, Aruparayil N, Gnanaraj J, Brown J, Jayne D. Barriers to training in laparoscopic surgery in low- and middle-income countries: A 16 systematic review. Trop Doct 2021; **51**: 408-414 [PMID: 33847545 DOI: 10.1177/0049475521998186]
- 17 Asbun HJ, Berguer R, Altamirano R, Castellanos H. Successfully establishing laparoscopic surgery programs in developing countries. Clinical results and lessons learned. Surg Endosc 1996; 10: 1000-1003 [PMID: 8864094 DOI: 10.1007/s004649900223]
- Cothren CC, Moore EE, Johnson JL, Moore JB, Ciesla DJ, Burch JM. Can we afford to do laparoscopic appendectomy in an academic 18 hospital? Am J Surg 2005; 190: 950-954 [PMID: 16307952 DOI: 10.1016/j.amjsurg.2005.08.026]
- Gehrman J, Angenete E, Björholt I, Lesén E, Haglind E. Cost-effectiveness analysis of laparoscopic and open surgery in routine Swedish care 19 for colorectal cancer. Surg Endosc 2020; 34: 4403-4412 [PMID: 31630289 DOI: 10.1007/s00464-019-07214-x]
- Brosens I, Gordon A, Campo R, Gordts S. Bowel injury in gynecologic laparoscopy. J Am Assoc Gynecol Laparosc 2003; 10: 9-13 [PMID: 20 12554987 DOI: 10.1016/s1074-3804(05)60227-7]
- Moghadamyeghaneh Z, Masoomi H, Mills SD, Carmichael JC, Pigazzi A, Nguyen NT, Stamos MJ. Outcomes of conversion of laparoscopic 21 colorectal surgery to open surgery. JSLS 2014; 18 [PMID: 25587213 DOI: 10.4293/JSLS.2014.00230]
- Madan Mohan Mudgal, Pradeep Kumar Kothiya, Naveen Kushwah, Raghvendra Singh. Port site complications following laparoscopic 22 surgeries: a prospective study. Int Surg J 2018; 5: 598-601 [DOI: 10.18203/2349-2902.isj20180360]
- Karthik S, Augustine AJ, Shibumon MM, Pai MV. Analysis of laparoscopic port site complications: A descriptive study. J Minim Access Surg 23 2013; 9: 59-64 [PMID: 23741110 DOI: 10.4103/0972-9941.110964]
- Bedada AG, Hsiao M, Bakanisi B, Motsumi M, Azzie G. Establishing a contextually appropriate laparoscopic program in resource-restricted 24 environments: experience in Botswana. Ann Surg 2015; 261: 807-811 [PMID: 24915782 DOI: 10.1097/SLA.00000000000691]



WJGS | https://www.wjgnet.com



Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: office@baishideng.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

