

Supplementary Table 1: Elixhauser comorbidities for present hospitalizations.

Factor	Alcoholic cirrhosis	No alcoholic cirrhosis (before matching)	P value	No alcoholic cirrhosis (after matching)	P value
N	1325	736685		1135	
Congestive heart failure	295 (22.3%)	108970 (14.8%)	<0.001	255 (22.5%)	0.90
Cardiac arrhythmias	340 (25.7%)	164510 (22.3%)	0.004	245 (21.6%)	0.018
Valvular heart disease	75 (5.7%)	27195 (3.7%)	<0.001	85 (7.5%)	0.067
Pulmonary circulation disorder	95 (7.2%)	36500 (5.0%)	<0.001	120 (10.6%)	0.003
Peripheral vascular disease	90 (6.8%)	30935 (4.2%)	<0.001	60 (5.3%)	0.12
Uncomplicated hypertension	435 (32.8%)	325495 (44.2%)	<0.001	345 (30.4%)	0.20
Chronic pulmonary diseases	320 (24.2%)	170925 (23.2%)	0.41	235 (20.7%)	0.042
Uncomplicated diabetes	160 (12.1%)	123285 (16.7%)	<0.001	170 (15.0%)	0.035
Complicated diabetes	375 (28.3%)	175575 (23.8%)	<0.001	305 (26.9%)	0.43
Hypothyroidism	155 (11.7%)	100015 (13.6%)	0.046	135 (11.9%)	0.88
Chronic kidney disease	280 (21.1%)	116295 (15.8%)	<0.001	260 (22.9%)	0.29
AIDS/HIV	15 (1.1%)	1155 (0.2%)	<0.001	15 (1.3%)	0.67
Rheumatoid arthritis/collagen vascular disorder	20 (1.5%)	20955 (2.8%)	0.003	25 (2.2%)	0.20
Obesity	270 (20.4%)	216220 (29.4%)	<0.001	230 (20.3%)	0.94
Weight loss	130 (9.8%)	36015 (4.9%)	<0.001	110 (9.7%)	0.92
Fluid and electrolyte disorder	765 (57.7%)	332875 (45.2%)	<0.001	675 (59.5%)	0.38
Blood loss anemia	25 (1.9%)	2175 (0.3%)	<0.001	10 (0.9%)	0.036
Deficiency anemia	160 (12.1%)	22970 (3.1%)	<0.001	115 (10.1%)	0.13
Drug abuse	110 (8.3%)	12175 (1.7%)	<0.001	115 (10.1%)	0.12
Psychoses	35 (2.6%)	12160 (1.7%)	0.005	45 (4.0%)	0.065
Depression	190 (14.3%)	81690 (11.1%)	<0.001	210 (18.5%)	0.005
Complicated hypertension	380 (28.7%)	164430 (22.3%)	<0.001	350 (30.8%)	0.24

AIDS, acquired immunodeficiency syndrome; HIV, human immunodeficiency virus