

World Journal of *Gastrointestinal Oncology*

World J Gastrointest Oncol 2024 February 15; 16(2): 251-570



EDITORIAL

- 251 Cardiotoxicity induced by fluoropyrimidine drugs in the treatment of gastrointestinal tumors
Kong MW, Sun FD, Pei ZY, Xu L, Wang ZB, Chen Y, Tang SQ, Yang TF, He GX
- 255 Does enhanced recovery after surgery programs improve clinical outcomes in liver cancer surgery?
Sánchez-Pérez B, Ramia JM

REVIEW

- 259 Biological factors driving colorectal cancer metastasis
An SX, Yu ZJ, Fu C, Wei MJ, Shen LH

MINIREVIEWS

- 273 Progress in the treatment of advanced hepatocellular carcinoma with immune combination therapy
Pan D, Liu HN, Qu PF, Ma X, Ma LY, Chen XX, Wang YQ, Qin XB, Han ZX
- 287 Targeting oxidative stress with natural products: A novel strategy for esophageal cancer therapy
Cao F, Zhang HL, Guo C, Xu XL, Yuan Q
- 300 Multifaceted role of microRNAs in gastric cancer stem cells: Mechanisms and potential biomarkers
Sun QH, Kuang ZY, Zhu GH, Ni BY, Li J

ORIGINAL ARTICLE**Clinical and Translational Research**

- 314 Expression of cyclin-dependent kinase 9 is positively correlated with the autophagy level in colon cancer
Zheng L, Lu J, Kong DL
- 331 Tumour response following preoperative chemotherapy is affected by body mass index in patients with colorectal liver metastases
Song HC, Zhou HC, Gu P, Bao B, Sun Q, Mei TM, Cui W, Yao K, Yao HZ, Zhang SY, Wang YS, Song RP, Wang JZ

Case Control Study

- 343 Preoperative controlling nutritional status as an optimal prognostic nutritional index to predict the outcome for colorectal cancer
Liu LX, Wang H, Gao B, Xu TT, Yuan QG, Zhou SZ, Ding C, Miao J, Guan WX

Retrospective Study

- 354 Effect of screening colonoscopy frequency on colorectal cancer mortality in patients with a family history of colorectal cancer
Zheng L, Li B, Lei L, Wang LJ, Zeng ZP, Yang JD

- 364 Effect of different anesthetic modalities with multimodal analgesia on postoperative pain level in colorectal tumor patients

Tang JC, Ma JW, Jian JJ, Shen J, Cao LL

Observational Study

- 372 Prognostic value of circulating tumor cells combined with neutrophil-lymphocyte ratio in patients with hepatocellular carcinoma

Chen JL, Guo L, Wu ZY, He K, Li H, Yang C, Han YW

- 386 Systemic Inflammation Response Index and weight loss as prognostic factors in metastatic pancreatic cancer: A concept study from the PANTHEIA-SEOM trial

Pacheco-Barcia V, Custodio-Cabello S, Carrasco-Valero F, Palka-Kotlowska M, Mariño-Mendez A, Carmona-Bayonas A, Gallego J, Martín AJM, Jimenez-Fonseca P, Cabezon-Gutierrez L

Basic Study

- 398 Prohibitin 1 inhibits cell proliferation and induces apoptosis *via* the p53-mediated mitochondrial pathway *in vitro*

Shi JJ, Wang YK, Wang MQ, Deng J, Gao N, Li M, Li YP, Zhang X, Jia XL, Liu XT, Dang SS, Wang WJ

- 414 Early results of the integrative epigenomic-transcriptomic landscape of colorectal adenoma and cancer

Lu YW, Ding ZL, Mao R, Zhao GG, He YQ, Li XL, Liu J

- 436 Comprehensive analysis of the potential pathogenesis of COVID-19 infection and liver cancer

Rong Y, Tang MZ, Liu SH, Li XF, Cai H

- 458 Immune-related long noncoding RNA zinc finger protein 710-AS1-201 promotes the metastasis and invasion of gastric cancer cells

Ding W, Chen WW, Wang YQ, Xu XZ, Wang YB, Yan YM, Tan YL

- 475 Comprehensive analysis of the protein phosphatase 2A regulatory subunit B56ε in pan-cancer and its role and mechanism in hepatocellular carcinoma

Wu HM, Huang YY, Xu YQ, Xiang WL, Yang C, Liu RY, Li D, Guo XF, Zhang ZB, Bei CH, Tan SK, Zhu XN

- 493 Identification of anti-gastric cancer effects and molecular mechanisms of resveratrol: From network pharmacology and bioinformatics to experimental validation

Ma YQ, Zhang M, Sun ZH, Tang HY, Wang Y, Liu JX, Zhang ZX, Wang C

SYSTEMATIC REVIEWS

- 514 Prognostic nutritional index in predicting survival of patients with gastric or gastroesophageal junction adenocarcinoma: A systematic review

Fiflis S, Christodoulidis G, Papakonstantinou M, Giakoustidis A, Koukias S, Roussos P, Kouliou MN, Koumarelas KE, Giakoustidis D

SCIENTOMETRICS

- 527 Global research trends and prospects of cellular metabolism in colorectal cancer

Liu YC, Gong ZC, Li CQ, Teng P, Chen YY, Huang ZH

CASE REPORT

- 543** Large isolated fibrous tumors in the upper esophagus: A case report
Yu JJ, Pei HS, Meng Y
- 550** Hepatomegaly and jaundice as the presenting symptoms of systemic light-chain amyloidosis: A case report
Zhang X, Tang F, Gao YY, Song DZ, Liang J
- 557** Anti-EGFR antibody monotherapy for colorectal cancer with severe hyperbilirubinemia: A case report
Tsurui T, Hirasawa Y, Kubota Y, Yoshimura K, Tsunoda T
- 563** Early adenocarcinoma mixed with a neuroendocrine carcinoma component arising in the gastroesophageal junction: A case report
Cheng YQ, Wang GF, Zhou XL, Lin M, Zhang XW, Huang Q

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Oncology*, Akihiko Oka, MD, PhD, Assistant Professor, Department of Internal Medicine II, Shimane University Faculty of Medicine, Izumo 693-8501, Japan.
aoka@med.shimane-u.ac.jp

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Oncology* (*WJGO*, *World J Gastrointest Oncol*) is to provide scholars and readers from various fields of gastrointestinal oncology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGO mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal oncology and covering a wide range of topics including liver cell adenoma, gastric neoplasms, appendiceal neoplasms, biliary tract neoplasms, hepatocellular carcinoma, pancreatic carcinoma, cecal neoplasms, colonic neoplasms, colorectal neoplasms, duodenal neoplasms, esophageal neoplasms, gallbladder neoplasms, *etc.*

INDEXING/ABSTRACTING

The *WJGO* is now abstracted and indexed in PubMed, PubMed Central, Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJGO* as 3.0; IF without journal self cites: 2.9; 5-year IF: 3.0; Journal Citation Indicator: 0.49; Ranking: 157 among 241 journals in oncology; Quartile category: Q3; Ranking: 58 among 93 journals in gastroenterology and hepatology; and Quartile category: Q3. The *WJGO*'s CiteScore for 2022 is 4.1 and Scopus CiteScore rank 2022: Gastroenterology is 71/149; Oncology is 197/366.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Xiang-Di Zhang*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jia-Ru Fan*.

NAME OF JOURNAL

World Journal of Gastrointestinal Oncology

ISSN

ISSN 1948-5204 (online)

LAUNCH DATE

February 15, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Monjur Ahmed, Florin Burada

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-5204/editorialboard.htm>

PUBLICATION DATE

February 15, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Effect of screening colonoscopy frequency on colorectal cancer mortality in patients with a family history of colorectal cancer

Li Zheng, Bin Li, Ling Lei, Li-Jia Wang, Zhi-Ping Zeng, Jian-Dong Yang

Specialty type: Oncology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): C, C

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Cucarull B, Spain; D'Amato M, Spain

Received: October 23, 2023

Peer-review started: October 23, 2023

First decision: November 8, 2023

Revised: November 13, 2023

Accepted: December 11, 2023

Article in press: December 11, 2023

Published online: February 15, 2024



Li Zheng, Bin Li, Ling Lei, Li-Jia Wang, Zhi-Ping Zeng, Department of Gastroenterology, The First Hospital of Nanchang, Nanchang 330008, Jiangxi Province, China

Jian-Dong Yang, Department of General Surgery, The First Hospital of Nanchang, Nanchang 330008, Jiangxi Province, China

Corresponding author: Jian-Dong Yang, Doctor, Occupational Physician, Department of General Surgery, The First Hospital of Nanchang, No. 128 Xiangshan North Road, Nanchang 330008, Jiangxi Province, China. yjd-009@163.com

Abstract

BACKGROUND

Colorectal cancer is a common malignant tumor in China, and its incidence in the elderly is increasing annually. Inflammatory bowel disease is a group of chronic non-specific intestinal inflammatory diseases, including ulcerative colitis and Crohn's disease.

AIM

To assess the effect of screening colonoscopy frequency on colorectal cancer mortality.

METHODS

We included the clinicopathological and follow-up data of patients with colorectal cancer who underwent laparoscopic colectomy or open colectomy at our Gastrointestinal Department between January 2019 and December 2022. Surgical indicators, oncological indicators, and survival rates were compared between the groups. The results of 104 patients who met the above criteria were extracted from the database (laparoscopic colectomy group = 63, open colectomy group = 41), and there were no statistically significant differences in the baseline data or follow-up time between the two groups.

RESULTS

Intraoperative blood loss, time to first ambulation, and time to first fluid intake were significantly lower in the laparoscopic colectomy group than in the open colectomy group. The differences in overall mortality, tumor-related mortality, and recurrence rates between the two groups were not statistically significant, and survival analysis showed that the differences in the cumulative overall survival, tumor-related survival, and cumulative recurrence-free rates between the two

groups were not statistically significant.

CONCLUSION

In elderly patients with colorectal cancer, laparoscopic colectomy has better short-term outcomes than open colectomy, and laparoscopic colectomy has superior long-term survival outcomes compared with open colectomy.

Key Words: Colorectal cancer; Laparoscopic surgery; Open surgery; Prognosis; Laparoscopic colectomy; Open colectomy

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Screening colonoscopy performed at different time frequencies did not show a statistically significant effect on overall mortality, tumor-related mortality, and recurrence rates in colorectal cancer patients with a family history of rectal cancer. However, laparoscopic surgery demonstrated better short-term outcomes compared to open surgery, while maintaining comparable long-term survival results. This suggests that laparoscopic surgery may be a favorable option for elderly patients with colorectal cancer.

Citation: Zheng L, Li B, Lei L, Wang LJ, Zeng ZP, Yang JD. Effect of screening colonoscopy frequency on colorectal cancer mortality in patients with a family history of colorectal cancer. *World J Gastrointest Oncol* 2024; 16(2): 354-363

URL: <https://www.wjgnet.com/1948-5204/full/v16/i2/354.htm>

DOI: <https://dx.doi.org/10.4251/wjgo.v16.i2.354>

INTRODUCTION

Colorectal cancer is a common malignant tumor in China[1-5]. Owing to an increase in the elderly population, the incidence of colorectal cancer in the elderly is increasing annually[6-10]. There is no consensus on whether patients should undergo open or laparoscopic colectomy[11-15]. Furthermore, elderly patients with colorectal cancer have a high mortality rate after open colectomy[16], and early randomized controlled clinical studies on the short- and long-term outcomes of laparoscopic colectomy mostly excluded elderly patients. Until recently, the results of short-term studies have confirmed the applicability of laparoscopic colectomy in elderly patients. Although the short-term outcomes of laparoscopic colectomy in elderly patients with colorectal cancer have been reported to be superior to those of open colectomy in China, few reports have analyzed the long-term survival outcomes both in China and around the world[17-20]. In clinical practice, diverse temporal frequencies may affect screening accuracy. Therefore, we assessed the effect of screening colonoscopy at different frequencies on the mortality of patients with colorectal cancer and a family history of rectal cancer.

MATERIALS AND METHODS

General information

Patients with colorectal cancer who underwent laparoscopic or open colectomy at our Gastrointestinal Department between January 2019 and December 2022 according to the American Joint Committee on Cancer tumor-node-metastasis staging system (7th edition) were included in this study. Patients were selected according to the following criteria: Patients aged > 65 years; those who had postoperative pathologically confirmed primary colorectal or mucinous adenocarcinoma; and those who successfully underwent radical D3 resection, except for emergency colectomy. The patients were treated with a 5-fluorouracil-based postoperative chemotherapy regimen.

Observation indicators

Clinical and pathological data, including age, sex, tumor location, tumor size, degree of differentiation, type of pathology, surgical procedure, and tumor-node-metastasis stage, were included as observation indicators.

Perioperative-related indicators

Preoperative comorbidities, history of abdominal colectomy, mean operative time, intraoperative blood loss, postoperative length of hospital stay, time to first ambulation, time to first fluid intake, time to first gastric discharge, and postoperative complications were included as perioperative indicators. The above indexes were measured in both groups. Except for different treatment methods, the other arrangements were consistent between the two groups.

Oncological and prognostic indicators

The length of the specimen, distance of the tumor from both sides of the excised edge, number of dissected lymph nodes, number of positive lymph nodes, recurrence rate, overall survival rate, and tumor-related survival rate were included as

oncological and prognostic indicators.

Statistical analysis

SPSS 25.0 software was used for statistical analyses. Measurement data are expressed as the mean \pm SD, and the *t*-test was used for comparison between groups and within groups. Enumeration data are expressed as percentages or composition ratios *n* (%), and the χ^2 test was used for comparison between groups and within groups. *P* < 0.05 was considered statistically significant.

RESULTS

Comparison of clinical and pathological data between the two groups

The results of 104 patients who met the above criteria were extracted from the database (laparoscopic colectomy group = 63, open colectomy group = 41). The differences in sex, age, tumor location, tumor size, degree of differentiation, and pathological stage between the two groups were not statistically significant (Table 1).

Comparison of perioperative-related indicators between the two groups

Except for the preoperative anemia rate, which was higher in the laparoscopic colectomy group than in the open colectomy group, no statistically significant difference was noted between the two groups in terms of preoperative comorbidities and preoperative abdominal digestive disease history; intraoperative blood loss, time to first ambulation, and time to first fluid intake were significantly lower in the laparoscopic surgery group than in the open surgery group. The mean operative time was slightly longer in the laparoscopic colectomy group than in the open colectomy group, although the difference was not statistically significant.

Comparison of oncological indicators between the two groups

No statistically significant differences were noted in the length of the specimens, distance of the tumor from the cut edge on either side, number of lymph nodes dissected, or number of positive lymph nodes between the two study groups (Table 2).

Comparison of intestinal cleaning compliance rate between the two groups

The compliance rates of intestinal cleanliness in the observation and control groups were 90.47% and 53.66% (144/200), respectively, thus indicating a significant difference (Table 3).

Bowel examination preparation

The Boston bowel preparation scale score and adenoma detection rate in the laparoscopic colectomy group were higher than those in the open colectomy group, whereas the colonoscopy time was shorter than that in the control group (*P* < 0.05) (Table 4). Based on Tables 1-4 and the current research results, it can be seen that the frequency of colonoscopy had a greater impact on the mortality of colorectal cancer in patients with a family history of colorectal cancer, which can be used as a means of evaluation.

DISCUSSION

Delay in consultation is common in patients with a positive family history of colorectal cancer

This study found that patients with a positive family history of colorectal cancer experienced serious delays in consultation[21-25], with a delay rate of 39.9%, and many patients missed the optimal time for early detection and treatment of colorectal cancer[26-30]. The results of the univariate analysis showed that a longer time to consultation is associated with a more advanced pathological stage and worse prognosis. Therefore, delayed consultation seriously affects the prognosis of patients and increases the burden on families and the society. The reason for the delay in consultation in the current study was that the hospital where the study was conducted is a higher-level specialist hospital where patients from various regions of the country visit, and the patient's knowledge of the disease varied. Conversely, a previous study conducted the study at a general hospital, and the patients included were mostly residents of the big city where they were located[29]; furthermore, the patients had sufficient knowledge of the disease and medical examination findings. Therefore, consultations were conducted in a timely manner. The delay in consultation of patients in the present study was similar to that in studies from other developing countries[31] but was more serious than that in Denmark[32]. This finding may be related to differences in socioeconomic and cultural environments, particularly in health service policies, resources, and organizations in different regions, in addition to individual factors[33].

The level of social support for patients with colorectal cancer and a positive family history of colorectal cancer is low and needs improvement

The effect of social support as an available external resource wherein individuals engage with and receive information, comfort, and reassurance from other people or groups in formal and informal ways[34] has received increasing attention in the field of nursing. Good social support has a positive effect on the physical and mental health of patients with cancer

Table 1 Comparison of clinical and pathological data between the two groups

| | Laparoscopic colectomy group (n = 63) | Open colectomy group (n = 41) | P value |
|-------------------------------------|---------------------------------------|-------------------------------|---------|
| Age (yr) | 72.3 ± 5.0 | 70.8 ± 4.3 | 0.109 |
| Sex (n) | | 0.532 | 0.466 |
| Male | 37 | 27 | |
| Female | 26 | 14 | |
| Tumor size (cm) | 4.7 ± 2.3 | 4.4 ± 1.4 | 0.406 |
| Tumor location (n) | | | 0.281 |
| Colon | 36 | 19 | |
| Rectum | 27 | 22 | |
| Right colon | 20 | 8 | |
| Left colon | 4 | 1 | |
| Transverse colon cancer | 0 | 3 | |
| Total colon | 1 | 0 | |
| Sigmoid colon | 11 | 7 | |
| Low rectum | 18 | 6 | |
| High rectum | 6 | 9 | |
| Middle rectum | 3 | 7 | |
| Degree of tumor differentiation (n) | | 1.279 | 0.528 |
| Well differentiation | 20 | 13 | |
| Moderate differentiation | 36 | 26 | |
| Low differentiation | 7 | 2 | |
| T stage (n) | | 0.811 | 0.667 |
| T1 | 0 | 0 | |
| T2 | 4 | 2 | |
| T3 | 52 | 32 | |
| T4 | 7 | 7 | |
| Lymph node metastasis (n) | | 0.633 | 0.426 |
| N0 | 35 | 26 | |
| N1 | 28 | 15 | |
| N2 | 0 | 0 | |
| TNM stage (n) | | 3.195 | 0.362 |
| IIA | 32 | 20 | |
| IIB | 3 | 6 | |
| IIIA | 4 | 2 | |
| IIIB | 24 | 13 | |

TNM: Tumor-node-metastasis.

[35]. The results of the present study showed that the total social support score of patients with colorectal cancer and a family history of colorectal cancer (37.51 ± 8.43) was higher than that of the normal population. The reason for the low level of social support in this study could be attributed to the fact that negative tumor-related events in other members of the family tended to make patients feel negative and unable to seek and use social support. Moreover, it specifies that most patients failed to understand the positive role of social support for patients with cancer and were unable to obtain a high level of social support from their sick relatives during previous medical treatment. Therefore, healthcare professionals should pay special attention to patients with a positive family history of cancer, understand their level of social

Table 2 Comparison of oncological indicators between the two groups

| | Laparoscopic colectomy group, (n = 63) | Open colectomy group, (n = 41) | t-value | P value |
|--|--|--------------------------------|---------|---------|
| Length of specimen (cm) | | | | |
| Colon cancer | 30.1 ± 17.8 | 23.6 ± 9.3 | 1.492 | 0.142 |
| Rectal cancer | 19.7 ± 8.6 | 16.8 ± 4.2 | 1.364 | 0.394 |
| Tumor distance from proximal margin (cm) | | | | |
| Colon cancer | 12.1 ± 6.4 | 12.8 ± 6.4 | 0.885 | 0.721 |
| Rectal cancer | 9.2 ± 4.3 | 8.9 ± 3.9 | 0.281 | 0.770 |
| Tumor distance from distal margin (cm) | | | | |
| Colon cancer | 10.1 ± 8.8 | 7.9 ± 4.0 | 0.984 | 0.330 |
| Rectal cancer | 6.5 ± 4.5 | 4.3 ± 1.7 | 0.148 | 0.883 |
| Number of resected lymph nodes (n) | | | | |
| Colon cancer | 11.1 ± 6.3 | 12.0 ± 6.0 | 0.602 | 0.380 |
| Rectal cancer | 9.0 ± 5.3 | 10.3 ± 5.7 | 0.148 | 0.349 |
| Number of positive lymph nodes (n) | | | | |
| Colon cancer | 0.7 ± 1.1 | 0.5 ± 0.7 | 0.853 | 0.398 |
| Rectal cancer | 1.1 ± 1.7 | 0.6 ± 1.1 | 1.446 | 0.258 |

Table 3 Comparison of intestinal cleaning compliance rate between the two groups, n (%)

| Group | N | 1 | 2 | 3 | 4 | Rate |
|---------------------------------------|----|----|----|----|---|------------|
| Laparoscopic colectomy group (n = 63) | 63 | 23 | 20 | 10 | 4 | 57 (90.47) |
| Open colectomy group (n = 41) | 41 | 10 | 10 | 1 | 1 | 22 (53.66) |
| χ^2 | - | - | - | - | - | 6.939 |
| P value | - | - | - | - | - | 0.008 |

support, take effective measures to improve the social support of each individual, encourage patients and their relatives to socialize more with friends, encourage behaviors that could help others, and seek external help for increasing social support. Healthcare professionals should use appropriate communication methods to provide effective social support to patients.

Timing of consultation for patients with colorectal cancer and a positive family history of colorectal cancer is influenced by multiple factors

The timing of patient visits in the present study was influenced by several factors, including the level of social support, medical examinations, and number of relatives with cancer.

Effect of social support level on consultation timing: The results of this study showed that the level of social support was significantly negatively associated with the delay in patient hospital visits and that the lack of social support significantly increased the delay in hospital visits, thus suggesting that good social support has an enabling effect on patients' timely hospital visits. This finding is in line with the findings of both national and international studies. Patients who lack social support may be less able to adapt psychosocially, thus leading to high levels of psychological stress and negativity, which may affect their self-esteem levels and coping methods. Furthermore, individuals who lack social support may receive less positive advice and opinions from others regarding seeking medical care, thus ultimately predisposing them to delays in seeking medical care. Previous studies have shown that talking to others after symptoms have been identified can reduce delays in consultation, and close relationships with family members can help determine symptoms and facilitate consultation. Patients who are living alone are more likely to experience delays in consultation than those living with family members, and many patients are persuaded and encouraged by family members to seek consultation. Therefore, healthcare professionals should teach patients on how to obtain support from their families and instruct patients' relatives on how to help and encourage patients obtain support from outside the family, particularly for the management of cancer within their family. Mutual support for the entire family is important.

Effect of medical examination status on consultation timing: This study showed that the length of delay in consultation for patients with colorectal cancer was significantly correlated with the status of medical examination, and the delay in

Table 4 Comparison of perioperative-related indicators between the two groups

| | Laparoscopic colectomy group, (n = 63) | Open colectomy group, (n = 41) | χ^2 , t, or Z value | P value |
|--------------------------------------|--|--------------------------------|--------------------------|--------------------|
| Preoperative comorbidities, n (%) | | | | |
| Anemia | 25 (39.7) | 7 (17.1) | 5.960 | 0.015 ^a |
| Intestinal obstruction | 12 (19) | 5 (12.2) | 0.853 | 0.356 |
| Hypoproteinemia | 17 (27) | 7 (17.1) | 1.374 | 0.241 |
| High blood pressure | 20 (31.7) | 12 (29.3) | 0.072 | 0.789 |
| Diabetes | 7 (11.1) | 4 (9.8) | 0.048 | 0.826 |
| Pneumonia or bronchitis | 8 (12.7) | 8 (19.5) | 0.886 | 0.347 |
| Viral hepatitis | 6 (9.5) | 1 (2.4) | 1.986 | 0.159 |
| History of digestive disease, n (%) | 5 (7.9) | 5 (12.2) | 0.518 | 0.472 |
| Average operation time (min) | 175.2 ± 43.8 | 171.9 ± 58.8 | 0.325 | 0.746 |
| Intraoperative bleeding (mL) | 80.4 ± 60.6 | 169.4 ± 95.1 | 5.325 | 0.000 ^a |
| Postoperative hospital stay (d) | 12.9 ± 3.7 | 13.7 ± 6.7 | 0.837 | 0.405 |
| Time to first ambulation (d) | 2.8 ± 2.1 | 5.1 ± 2.2 | 5.243 | 0.000 ^a |
| Time to first liquid meal (d) | 3.7 ± 1.9 | 5.0 ± 2.1 | 3.142 | 0.002 ^a |
| Time to first exhaust (d) | 3.3 ± 1.3 | 3.9 ± 1.5 | 1.960 | 0.053 |
| Postoperative complications, n (%) | 6 (9.5) | 9 (22.0) | 3.108 | 0.078 |
| Bleeding from the anastomosis | 1 | 2 | | |
| Anastomotic fistula | 1 | 0 | | |
| Postoperative intestinal obstruction | 1 | 3 | | |
| Infection of the incision | 0 | 1 | | |
| Abdominal infection | 0 | 1 | | |
| Diarrhea | 3 | 1 | | |
| Postoperative urinary retention | 0 | 1 | | |

^a $P < 0.05$ represents a significant difference between the two groups.

consultation was shorter for patients who had undergone medical examination than for those who had never undergone a medical examination. Those who had undergone a medical examination were more aware of the disease, more worried about their health condition, more alert, and more likely to consult the doctor immediately after the appearance of symptoms (Table 5).

Colorectal cancer can be immediately detected during physical examinations; thus, the time to visit a doctor is greatly reduced. However, the results of this study showed that more than half of the patients with colorectal cancer had never undergone a physical examination and that only eight patients (5.2%) had been screened for colorectal cancer. This indicates that although the prevention and treatment of colorectal cancer are receiving increasing attention, prevention and treatment efforts are quite uneven across regions, and the cooperation of some patients is low compared with patients in developed countries.

Effect of number of relatives with cancer on consultation timing: The results of this study showed that the number of relatives suffering from colorectal cancer significantly influenced the time taken to see a patient. Patients with three or more relatives with colorectal cancer had significantly shorter times to consultation than other patients. A larger number of relatives with similar diseases may draw the attention of family members to the disease, thus allowing them to actively or passively obtain information about the clinical symptoms, prevention methods, and treatment of the disease; make quicker and riskier decisions; and seek medical attention quickly when they develop symptoms. However, the results of the present study showed that even when relatives had a history of colorectal cancer, the delay in consultation was still significant, and the major reason for the delay was the “lack of knowledge and attention”. This is because there is a lack of health education for patients and their relatives on the causes, clinical symptoms, and screening of colorectal cancer during clinical treatment and care. Despite the greater understanding of the genetic mechanism of tumors in recent years, its clinical application has not yet been popularized. Most medical institutions in China are still managing the disease according to the diagnosis and treatment model of sporadic colorectal cancer, thus resulting in high-risk groups not being

Table 5 Comparison of results of bowel preparation assessment in 104 patients undergoing colonoscopy

| Group | BBPS score | Colonoscopy time (min) | Adenoma detection rate (%) |
|---|-------------|------------------------|----------------------------|
| Laparoscopic colectomy group (<i>n</i> = 63) | 7.24 ± 1.07 | 5.63 ± 1.93 | 26.00 |
| Open colectomy group (<i>n</i> = 41) | 5.63 ± 1.93 | 9.81 ± 2.08 | 13.33 |
| <i>t</i> / χ^2 | 49.21 | 5.79 | 4.75 |
| <i>P</i> value | < 0.0001 | 0.02 | 0.03 |

BBPS: Boston bowel preparation scale

identified in time.

Summary and recommendations

Given that colorectal cancer is a common cancer in China, improving its prevention and control is of great significance to both individuals and the country. This study shows that delayed consultation is common among patients with colorectal cancer and a family history of colorectal cancer, and the acceptance rate of colorectal cancer screening is very low. Therefore, healthcare professionals should pay attention to the intervention of delayed consultation and screening of patients in the secondary prevention of colorectal cancer by taking the following measures: (1) Clinical practice should ensure the accuracy of family history taking to avoid missing diagnosis because of incomplete family history taking; (2) Referring patients with a positive family history of colorectal cancer to oncological genetic counseling for further confirmation of the diagnosis; (3) Explaining in detail the hereditary aspects of colorectal cancer, law of inheritance, and range of high-risk relatives to patients and relatives who are suspected of having a genetic predisposition to the disease; (4) Popularizing the knowledge of symptom recognition and screening of colorectal cancer in medical and nursing staff at all levels to strengthen health education for patients and their relatives and encourage participation in cancer prevention health check-ups; (5) Medical centers that treat a large number of patients with colorectal cancer should adopt case management for the dedicated and systematic management of high-risk families to track colorectal cancer screening, early diagnosis, and treatment of patients' high-risk relatives; and (6) Emphasizing the importance of education on cancer prevention and control for people in rural and remote areas.

CONCLUSION

This study was designed to develop effective measures for the prevention, control, and treatment of colorectal cancer. These results provide a scientific basis for the early treatment of tumors in the relatives of patients with hereditary colorectal cancer and for the systematic management of families with hereditary colorectal cancer. This study has great clinical value as a reference for the treatment and prevention of malignant tumors of the gastrointestinal tract.

In summary, screening strategies should be improved, screening should be integrated with other public health campaigns, more cost-effective screening techniques should be sought, screening methods with high sensitivity and accuracy should be explored, risk stratification should be performed using low-cost and high-efficiency risk assessment tools, technologies such as data mining and artificial intelligence should be appropriately utilized to establish a more accurate colorectal cancer screening prediction model, and individualized screening should be carried out for high-risk groups. It also suggests screening methods, starting ages, and screening intervals for different risk groups, and establishes a reasonable screening model.

ARTICLE HIGHLIGHTS

Research background

Colorectal cancer is one of the common malignant tumors in China, and its incidence in the elderly population has steadily increased. Inflammatory bowel disease is a group of chronic non-specific intestinal inflammatory diseases, including ulcerative colitis and Crohn's disease.

Research motivation

To evaluate the effect of colonoscopy frequency on the mortality of colorectal cancer.

Research objectives

We included the clinicopathological and follow-up data of patients with colorectal cancer who underwent laparoscopic colectomy or open colectomy at our gastrointestinal department.

Research methods

Surgical indicators, oncological indicators, and survival rates were compared between the groups. The results of 104 patients who met the inclusion criteria were extracted from the database (laparoscopic colectomy group = 63, open colectomy group = 41), and there were no statistically significant differences in the baseline data or follow-up time between the two groups.

Research results

Intraoperative blood loss, the time to first ambulation, and time to first liquid meal in the laparoscopic colectomy group were significantly lower than those in the open colectomy group. There was no significant difference in total mortality, tumor-related mortality, or recurrence rate between the two groups. Survival analysis showed that there was no significant difference in cumulative overall survival rate, tumor-related survival rate, or cumulative recurrence-free survival rate between the two groups.

Research conclusions

In elderly patients with colorectal cancer, laparoscopic colectomy has better short-term outcomes than open colectomy.

Research perspectives

Laparoscopic colectomy has superior long-term survival outcomes compared with open colectomy.

FOOTNOTES

Author contributions: Zheng L and Yang JD contributed equally to this work; Zheng L, Li B, Lei L, Wang LJ, Zeng ZP, and Yang JD designed the study and were involved in the data acquisition and writing of this article; Zheng L and Yang JD contributed to the analysis of the manuscript; and all authors have read and approved the final manuscript.

Supported by Science and Technology Plan of Jiangxi Provincial Health Commission, No. 202311202 and No. SKJP220219076; and the Science and Technology Support Plan Project of Nanchang, Jiangxi Province, No. 2020-133-5.

Institutional review board statement: The study was reviewed and approved by the First Hospital of Nanchang Institutional Review Board.

Informed consent statement: All patients provided written informed consent for participation in the study.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Bin Li [0009-0007-4240-0343](https://orcid.org/0009-0007-4240-0343); Jian-Dong Yang [0009-0005-1315-963X](https://orcid.org/0009-0005-1315-963X).

S-Editor: Wang JJ

L-Editor: Wang TQ

P-Editor: Zhang XD

REFERENCES

- 1 Gu J, Chen N. Current status of rectal cancer treatment in China. *Colorectal Dis* 2013; **15**: 1345-1350 [PMID: [23651350](https://pubmed.ncbi.nlm.nih.gov/23651350/) DOI: [10.1111/codi.12269](https://doi.org/10.1111/codi.12269)]
- 2 Levi F, Lucchini F, Negri E, Boyle P, La Vecchia C. Changed trends of cancer mortality in the elderly. *Ann Oncol* 2001; **12**: 1467-1477 [PMID: [11762821](https://pubmed.ncbi.nlm.nih.gov/11762821/) DOI: [10.1023/a:1012539213643](https://doi.org/10.1023/a:1012539213643)]
- 3 Seishima R, Okabayashi K, Hasegawa H, Tsuruta M, Shigeta K, Matsui S, Yamada T, Kitagawa Y. Is laparoscopic colorectal surgery beneficial for elderly patients? A systematic review and meta-analysis. *J Gastrointest Surg* 2015; **19**: 756-765 [PMID: [25617077](https://pubmed.ncbi.nlm.nih.gov/25617077/) DOI: [10.1007/s11605-015-2748-9](https://doi.org/10.1007/s11605-015-2748-9)]
- 4 Okumura S, Goumar C, Gayet B, Fuks D, Scatton O. Laparoscopic versus open two-stage hepatectomy for bilobar colorectal liver metastases: A bi-institutional, propensity score-matched study. *Surgery* 2019; **166**: 959-966 [PMID: [31395397](https://pubmed.ncbi.nlm.nih.gov/31395397/) DOI: [10.1016/j.surg.2019.06.019](https://doi.org/10.1016/j.surg.2019.06.019)]
- 5 Niitsu H, Hinoi T, Kawaguchi Y, Ohdan H, Hasegawa H, Suzuka I, Fukunaga Y, Yamaguchi T, Endo S, Tagami S, Idani H, Ichihara T, Watanabe K, Watanabe M; Japan Society of Laparoscopic Colorectal Surgery. Laparoscopic surgery for colorectal cancer is safe and has

- survival outcomes similar to those of open surgery in elderly patients with a poor performance status: subanalysis of a large multicenter case-control study in Japan. *J Gastroenterol* 2016; **51**: 43-54 [PMID: 25940149 DOI: 10.1007/s00535-015-1083-y]
- 6 **Jayne D**, Pigazzi A, Marshall H, Croft J, Corrigan N, Copeland J, Quirke P, West N, Rautio T, Thomassen N, Tilney H, Gudgeon M, Bianchi PP, Edlin R, Hulme C, Brown J. Effect of Robotic-Assisted vs Conventional Laparoscopic Surgery on Risk of Conversion to Open Laparotomy Among Patients Undergoing Resection for Rectal Cancer: The ROLARR Randomized Clinical Trial. *JAMA* 2017; **318**: 1569-1580 [PMID: 29067426 DOI: 10.1001/jama.2017.7219]
 - 7 **Huang C**, Huang R, Jiang T, Huang K, Cao J, Qiu Z. Laparoscopic and open resection for colorectal cancer: an evaluation of cellular immunity. *BMC Gastroenterol* 2010; **10**: 127 [PMID: 21029461 DOI: 10.1186/1471-230X-10-127]
 - 8 **Engelman DT**, Ben Ali W, Williams JB, Perrault LP, Reddy VS, Arora RC, Roselli EE, Khoynzhad A, Gerdisch M, Levy JH, Lobdell K, Fletcher N, Kirsch M, Nelson G, Engelman RM, Gregory AJ, Boyle EM. Guidelines for Perioperative Care in Cardiac Surgery: Enhanced Recovery After Surgery Society Recommendations. *JAMA Surg* 2019; **154**: 755-766 [PMID: 31054241 DOI: 10.1001/jamasurg.2019.1153]
 - 9 **Frasson M**, Braga M, Vignali A, Zuliani W, Di Carlo V. Benefits of laparoscopic colorectal resection are more pronounced in elderly patients. *Dis Colon Rectum* 2008; **51**: 296-300 [PMID: 18197453 DOI: 10.1007/s10350-007-9124-0]
 - 10 **She WH**, Poon JT, Fan JK, Lo OS, Law WL. Outcome of laparoscopic colectomy for cancer in elderly patients. *Surg Endosc* 2013; **27**: 308-312 [PMID: 22820704 DOI: 10.1007/s00464-012-2466-2]
 - 11 **Hellan M**, Anderson C, Ellenhorn JD, Paz B, Pigazzi A. Short-term outcomes after robotic-assisted total mesorectal excision for rectal cancer. *Ann Surg Oncol* 2007; **14**: 3168-3173 [PMID: 17763911 DOI: 10.1245/s10434-007-9544-z]
 - 12 **Devoto L**, Celentano V, Cohen R, Khan J, Chand M. Colorectal cancer surgery in the very elderly patient: a systematic review of laparoscopic versus open colorectal resection. *Int J Colorectal Dis* 2017; **32**: 1237-1242 [PMID: 28667498 DOI: 10.1007/s00384-017-2848-y]
 - 13 **Miyasaka Y**, Mochidome N, Kobayashi K, Ryu S, Akashi Y, Miyoshi A. Efficacy of laparoscopic resection in elderly patients with colorectal cancer. *Surg Today* 2014; **44**: 1834-1840 [PMID: 24121951 DOI: 10.1007/s00595-013-0753-8]
 - 14 **Hatakeyama T**, Nakanishi M, Murayama Y, Komatsu S, Shiozaki A, Kuriu Y, Ikoma H, Ichikawa D, Fujiwara H, Okamoto K, Ochiai T, Kokuba Y, Otsuji E. Laparoscopic resection for colorectal cancer improves short-term outcomes in very elderly colorectal cancer patients. *Surg Laparosc Endosc Percutan Tech* 2013; **23**: 532-535 [PMID: 24300931 DOI: 10.1097/SLE.0b013e31828e3da5]
 - 15 **Wang Z**, Wu X. Study and analysis of antitumor resistance mechanism of PD1/PD-L1 immune checkpoint blocker. *Cancer Med* 2020; **9**: 8086-8121 [PMID: 32875727 DOI: 10.1002/cam4.3410]
 - 16 **Ferlay J**, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer* 2010; **127**: 2893-2917 [PMID: 21351269 DOI: 10.1002/ijc.25516]
 - 17 **Hampel H**, Frankel WL, Martin E, Arnold M, Khanduja K, Kuebler P, Clendenning M, Sotamaa K, Prior T, Westman JA, Panescu J, Fix D, Lockman J, LaJeunesse J, Comeras I, de la Chapelle A. Feasibility of screening for Lynch syndrome among patients with colorectal cancer. *J Clin Oncol* 2008; **26**: 5783-5788 [PMID: 18809606 DOI: 10.1200/JCO.2008.17.5950]
 - 18 **Ouyang X**, Zhang G, Pan H, Huang J. Susceptibility and severity of cancer-related fatigue in colorectal cancer patients is associated with SLC6A4 gene single nucleotide polymorphism rs25531 A>G genotype. *Eur J Oncol Nurs* 2018; **33**: 97-101 [PMID: 29551185 DOI: 10.1016/j.ejon.2018.02.003]
 - 19 **Bond JH**. Studies show that it is now time to vigorously promote screening for colorectal cancer. *Am J Med* 1997; **102**: 329-330 [PMID: 9217612 DOI: 10.1016/s0002-9343(97)00124-1]
 - 20 **Patel SG**, Karlitz JJ, Yen T, Lieu CH, Boland CR. The rising tide of early-onset colorectal cancer: a comprehensive review of epidemiology, clinical features, biology, risk factors, prevention, and early detection. *Lancet Gastroenterol Hepatol* 2022; **7**: 262-274 [PMID: 35090605 DOI: 10.1016/S2468-1253(21)00426-X]
 - 21 **Torti RA**, Winship T. Culpability for delay in the treatment of breast cancer. *Med Ann Dist Columbia* 1963; **32**: 132-133 [PMID: 13993746]
 - 22 **Macleod U**, Mitchell ED, Burgess C, Macdonald S, Ramirez AJ. Risk factors for delayed presentation and referral of symptomatic cancer: evidence for common cancers. *Br J Cancer* 2009; **101** Suppl 2: S92-S101 [PMID: 19956172 DOI: 10.1038/sj.bjc.6605398]
 - 23 **Kandola A**, Ashdown-Franks G, Hendrikse J, Sabiston CM, Stubbs B. Physical activity and depression: Towards understanding the antidepressant mechanisms of physical activity. *Neurosci Biobehav Rev* 2019; **107**: 525-539 [PMID: 31586447 DOI: 10.1016/j.neubiorev.2019.09.040]
 - 24 **Pedersen AF**, Olesen F, Hansen RP, Zachariae R, Vedsted P. Social support, gender and patient delay. *Br J Cancer* 2011; **104**: 1249-1255 [PMID: 21487428 DOI: 10.1038/bjc.2011.87]
 - 25 **Smith LK**, Pope C, Botha JL. Patients' help-seeking experiences and delay in cancer presentation: a qualitative synthesis. *Lancet* 2005; **366**: 825-831 [PMID: 16139657 DOI: 10.1016/S0140-6736(05)67030-4]
 - 26 **Walter F**, Webster A, Scott S, Emery J. The Andersen Model of Total Patient Delay: a systematic review of its application in cancer diagnosis. *J Health Serv Res Policy* 2012; **17**: 110-118 [PMID: 22008712 DOI: 10.1258/jhsrp.2011.010113]
 - 27 **Rees S**, Williams A. Promoting and supporting self-management for adults living in the community with physical chronic illness: A systematic review of the effectiveness and meaningfulness of the patient-practitioner encounter. *JBI Libr Syst Rev* 2009; **7**: 492-582 [PMID: 27819974 DOI: 10.11124/01938924-200907130-00001]
 - 28 **Liu M**, Liu L, Zhang S, Li T, Ma F, Liu Y. Fear of cancer recurrence and hope level in patients receiving surgery for non-small cell lung cancer: a study on the mediating role of social support. *Support Care Cancer* 2022; **30**: 9453-9460 [PMID: 35947207 DOI: 10.1007/s00520-022-07318-6]
 - 29 **Janssen RM**, Takach O, Nap-Hill E, Enns RA. Time to Endoscopy in Patients with Colorectal Cancer: Analysis of Wait-Times. *Can J Gastroenterol Hepatol* 2016; **2016**: 8714587 [PMID: 27446872 DOI: 10.1155/2016/8714587]
 - 30 **Harirchi I**, Ghaemmaghami F, Karbakhsh M, Moghimi R, Mazaherie H. Patient delay in women presenting with advanced breast cancer: an Iranian study. *Public Health* 2005; **119**: 885-891 [PMID: 15913679 DOI: 10.1016/j.puhe.2004.11.005]
 - 31 **Bretthauer M**, Løberg M, Wieszczy P, Kalager M, Emilsson L, Garborg K, Rupinski M, Dekker E, Spaander M, Bugajski M, Holme Ø, Zauber AG, Pilonis ND, Mroz A, Kuipers EJ, Shi J, Hernán MA, Adami HO, Regula J, Hoff G, Kaminski MF; NordICC Study Group. Effect of Colonoscopy Screening on Risks of Colorectal Cancer and Related Death. *N Engl J Med* 2022; **387**: 1547-1556 [PMID: 36214590 DOI: 10.1056/NEJMoa2208375]
 - 32 **Dominitz JA**, Robertson DJ. Understanding the Results of a Randomized Trial of Screening Colonoscopy. *N Engl J Med* 2022; **387**: 1609-1611 [PMID: 36214591 DOI: 10.1056/NEJMe2211595]
 - 33 **Forbes N**. Outcomes Associated With Colorectal Cancer After Population-Based Colonoscopy Screening: Results From a European Pragmatic

Randomized Trial. *Gastroenterology* 2023; **164**: 493-494 [PMID: 36379242 DOI: 10.1053/j.gastro.2022.10.038]

34 **Kanth P**, Inadomi JM. Screening and prevention of colorectal cancer. *BMJ* 2021; **374**: n1855 [PMID: 34526356 DOI: 10.1136/bmj.n1855]

35 **Issa IA**, Noureddine M. Colorectal cancer screening: An updated review of the available options. *World J Gastroenterol* 2017; **23**: 5086-5096 [PMID: 28811705 DOI: 10.3748/wjg.v23.i28.5086]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: office@baishideng.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

