

World Journal of *Gastroenterology*

World J Gastroenterol 2024 January 21; 30(3): 199-285



Contents

Weekly Volume 30 Number 3 January 21, 2024

EDITORIAL

- 199 Toward less invasive coloproctology: The future is out there
Emile SH, Ragheb J

OPINION REVIEW

- 204 Garg incontinence scores: New scoring system on the horizon to evaluate fecal incontinence. Will it make a difference?
Tsarkov P, Tulina I, Sheikh P, Shlyk DD, Garg P

REVIEW

- 211 Why is early detection of colon cancer still not possible in 2023?
Tonini V, Zanni M
- 225 Relationship among Parkinson's disease, constipation, microbes, and microbiological therapy
Yuan XY, Chen YS, Liu Z

ORIGINAL ARTICLE

Retrospective Cohort Study

- 238 Effectiveness of antibiotic prophylaxis for acute esophageal variceal bleeding in patients with band ligation: A large observational study
Ichita C, Shimizu S, Goto T, Haruki U, Itoh N, Iwagami M, Sasaki A

Basic Study

- 252 SLC6A14 promotes ulcerative colitis progression by facilitating NLRP3 inflammasome-mediated pyroptosis
Gu Q, Xia H, Song YQ, Duan J, Chen Y, Zhang Y, Chen HP, Zhang L
- 268 Calcium/calcimimetic *via* calcium-sensing receptor ameliorates cholera toxin-induced secretory diarrhea in mice
Tang LQ, Fraebel J, Jin S, Winesett SP, Harrell J, Chang WH, Cheng SX

LETTER TO THE EDITOR

- 280 Use of curcumin and its nanopreparations in the treatment of inflammatory bowel disease
Meng ZW, Chang B, Sang LX
- 283 Cholecystokinin and cholecystokinin-A receptor: An attractive treatment strategy for biliary dyskinesia?
Chang J, Liu Y, Jiang TC, Zhao L, Liu JW

ABOUT COVER

Editorial Board Member of *World Journal of Gastroenterology*, Fernando J Corrales, PhD, Professor, Functional Proteomics Laboratory, National Biotechnology Center (CNB-CSIC), Madrid 28049, Spain. fcorrales@cnb.csic.es

AIMS AND SCOPE

The primary aim of *World Journal of Gastroenterology* (WJG, *World J Gastroenterol*) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. WJG mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The WJG is now abstracted and indexed in Science Citation Index Expanded (SCIE), MEDLINE, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJG as 4.3; Quartile category: Q2. The WJG's CiteScore for 2021 is 8.3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yi-Xuan Cai; Production Department Director: Xu Guo; Editorial Office Director: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Andrzej S Tarnawski

EXECUTIVE ASSOCIATE EDITORS-IN-CHIEF

Xian-Jun Yu (Pancreatic Oncology), Jian-Gao Fan (Chronic Liver Disease), Hou-Bao Liu (Biliary Tract Disease)

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

PUBLICATION DATE

January 21, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

PUBLISHING PARTNER

Shanghai Pancreatic Cancer Institute and Pancreatic Cancer Institute, Fudan University
Biliary Tract Disease Institute, Fudan University

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

POLICY OF CO-AUTHORS

<https://www.wjgnet.com/bpg/GerInfo/310>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

PUBLISHING PARTNER's OFFICIAL WEBSITE

<https://www.shca.org.cn>
<https://www.zs-hospital.sh.cn>



Retrospective Cohort Study

Effectiveness of antibiotic prophylaxis for acute esophageal variceal bleeding in patients with band ligation: A large observational study

Chikamasa Ichita, Sayuri Shimizu, Tadahiro Goto, Uojima Haruki, Naoya Itoh, Masao Iwagami, Akiko Sasaki

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): A
Grade B (Very good): 0
Grade C (Good): C, C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Abraham P, India; Haj Ali S, Jordan; Sukocheva OA, Australia

Received: October 4, 2023

Peer-review started: October 4, 2023

First decision: December 6, 2023

Revised: December 12, 2023

Accepted: January 3, 2024

Article in press: January 3, 2024

Published online: January 21, 2024



Chikamasa Ichita, Uojima Haruki, Akiko Sasaki, Gastroenterology Medicine Center, Shonan Kamakura General Hospital, Kamakura 247-8533, Kanagawa, Japan

Chikamasa Ichita, Sayuri Shimizu, Tadahiro Goto, Department of Health Data Science, Yokohama City University, Yokohama 236-0027, Kanagawa, Japan

Tadahiro Goto, TXP Research, TXP Medical Co., Ltd., Chiyoda-ku 101-0042, Tokyo, Japan

Tadahiro Goto, Department of Clinical Epidemiology and Health Economics, School of Public Health, The University of Tokyo, Bunkyo-ku 113-0033, Tokyo, Japan

Uojima Haruki, Department of Genome Medical Sciences Project, Research Institute, National Center for Global Health and Medicine, Ichikawa 272-8516, Chiba, Japan

Naoya Itoh, Division of Infectious Diseases, Aichi Cancer Center, Nagoya 464-8681, Aichi, Japan

Masao Iwagami, Department of Health Services Research, Institute of Medicine, University of Tsukuba, Tsukuba 305-8575, Ibaraki, Japan

Corresponding author: Chikamasa Ichita, MD, Chief, Gastroenterology Medicine Center, Shonan Kamakura General Hospital, No. 1370-1 Okamoto, Kamakura 247-8533, Kanagawa, Japan. ichikamasa@yahoo.co.jp

Abstract

BACKGROUND

Esophageal variceal bleeding is a severe complication associated with liver cirrhosis and typically necessitates endoscopic hemostasis. The current standard treatment is endoscopic variceal ligation (EVL), and Western guidelines recommend antibiotic prophylaxis following hemostasis. However, given the improvements in prognosis for variceal bleeding due to advancements in the management of bleeding and treatments of liver cirrhosis and the global concerns regarding the emergence of multidrug-resistant bacteria, there is a need to reassess the use of routine antibiotic prophylaxis after hemostasis.

AIM

To evaluate the effectiveness of antibiotic prophylaxis in patients treated for EVL.

METHODS

We conducted a 13-year observational study using the Tokushukai medical database across 46 hospitals. Patients were divided into the prophylaxis group (received antibiotics on admission or the next day) and the non-prophylaxis group (did not receive antibiotics within one day of admission). The primary outcome was composed of 6-wk mortality, 4-wk rebleeding, and 4-wk spontaneous bacterial peritonitis (SBP). The secondary outcomes were each individual result and in-hospital mortality. A logistic regression with inverse probability of treatment weighting was used. A subgroup analysis was conducted based on the Child-Pugh classification to determine its influence on the primary outcome measures, while sensitivity analyses for antibiotic type and duration were also performed.

RESULTS

Among 980 patients, 790 were included (prophylaxis: 232, non-prophylaxis: 558). Most patients were males under the age of 65 years with a median Child-Pugh score of 8. The composite primary outcomes occurred in 11.2% of patients in the prophylaxis group and 9.5% in the non-prophylaxis group. No significant differences in outcomes were observed between the groups (adjusted odds ratio, 1.11; 95% confidence interval, 0.61-1.99; $P = 0.74$). Individual outcomes such as 6-wk mortality, 4-wk rebleeding, 4-wk onset of SBP, and in-hospital mortality were not significantly different between the groups. The primary outcome did not differ between the Child-Pugh subgroups. Similar results were observed in the sensitivity analyses.

CONCLUSION

No significant benefit to antibiotic prophylaxis for esophageal variceal bleeding treated with EVL was detected in this study. Global reassessment of routine antibiotic prophylaxis is imperative.

Key Words: Esophageal varices; Endoscopic hemostasis; Antibiotic prophylaxis; Liver cirrhosis; Inverse probability of treatment weighting

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Esophageal variceal bleeding, a serious condition linked to liver cirrhosis, often requires endoscopic treatment. While western guidelines suggest using antibiotics after endoscopic treatment, data from multiple Japanese medical centers indicates that these prophylactic antibiotics are not associated with 6-wk mortality. Based on advances in cirrhosis treatment and the appropriate use of antibiotics, the necessity of routine prophylaxis must be reassessed.

Citation: Ichita C, Shimizu S, Goto T, Haruki U, Itoh N, Iwagami M, Sasaki A. Effectiveness of antibiotic prophylaxis for acute esophageal variceal bleeding in patients with band ligation: A large observational study. *World J Gastroenterol* 2024; 30(3): 238-251

URL: <https://www.wjgnet.com/1007-9327/full/v30/i3/238.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v30.i3.238>

INTRODUCTION

Esophageal variceal bleeding is a life-threatening complication in patients with liver cirrhosis[1], and endoscopic hemostasis is recommended as the first line of treatment[2]. However, even after hemostasis, there is a risk for infection, such as spontaneous bacterial peritonitis (SBP)[3], and rebleeding triggered by these infections[4,5], which are believed to contribute to increased mortality. While current Japanese guidelines do not specifically address antibiotic prophylaxis[6], western guidelines advocate prophylaxis for all patients[7-10].

The rationale for this recommendation lies in several studies conducted prior to the early 2000s that reported high mortality and an infection incidence of approximately 30% after upper gastrointestinal bleeding in patients with cirrhosis [11,12]. However, since the late 2000s, both mortality and infection incidence following upper gastrointestinal bleeding have improved, to less than 10%[13,14]. This marked improvement can be attributed to the shift in the recommended hemostatic method from endoscopic injection sclerotherapy (EIS) to endoscopic variceal ligation (EVL) and advancements in the treatment of liver cirrhosis[15]. These findings have prompted a reconsideration of the current practice of universal antibiotic prophylaxis across all clinical scenarios. Recent reports suggest that such prophylaxis may not always be necessary in modern medical settings[16-19]. Yet, these assertions are primarily from single-center observational studies; no multi-center study has been conducted. Furthermore, the inappropriate use of antibiotics, which has been identified as a cause of the emergence of multidrug-resistant bacteria, is a global issue[20].

Therefore, we aim to reassess the effectiveness of antibiotic prophylaxis in patients with esophageal variceal bleeding treated with EVL using data from several centers in Japan over a 13-year period. It is crucial to conduct research in regions such as Japan, where the guidelines do not recommend antibiotic prophylaxis after hemostasis.

MATERIALS AND METHODS

Data source

We conducted a retrospective cohort study using data from the Tokushukai medical database[21]. The Tokushukai group is a large hospital group in Japan that manages more than 70 hospitals nationwide. 50 hospitals are part of the Diagnosis Procedure Combination (DPC) system. The DPC system is a comprehensive payment system used in Japan that is specifically designed for acute care[22,23]. The Tokushukai Medical Database primarily comprises administrative claims data (specifically, DPC inpatient data) and electronic health records, including inpatient and outpatient blood test results.

The DPC inpatient data includes patient age; sex; admission and discharge dates; discharge status; main diagnosis; comorbidities at admission; post-admission complications recorded by the attending physician using the 2003 version of the International Classification of Diseases, 10th revision (ICD-10) codes; types of surgery (coded with original codes and text data in Japanese); and daily records of drugs and procedures. A distinguishing feature of this database is its capacity to access individual patient medical records. If necessary, additional details can be retrieved directly from these records.

Patient selection

This study included adult patients (aged ≥ 18 years) with esophageal variceal bleeding (ICD-10 code, I850) who underwent emergency EVL on the day of admission between January 2010 and December 2022. We excluded patients with the following criteria: (1) Death occurring on the day of admission or the following day; (2) discharge on the day of admission or the following day; (3) use of a mechanical ventilator on the day of admission or the following day; (4) use of continuous renal replacement therapy (CRRT) on the day of admission or the following day; (5) interventional radiology (IVR) or a surgical procedure on the day of admission or the following day; and (6) the presence of symptoms of infection, defined as having a fever of ≥ 38 °C or obtaining a blood culture, on the day of admission or the following day.

Exposure

The patients were divided into prophylaxis and non-prophylaxis groups. The prophylaxis group included patients who received antibiotics on the day of admission or the following day. The non-prophylaxis group included patients who did not receive antibiotics on the day of admission or the following day. The types of antibiotics considered in this study are detailed in Table 1, and the duration of administration was assumed to be at least one day.

Variables and outcomes

The variables included age, sex, the Barthel Index[24], the Child-Pugh Score and classification[25,26], the Charlson Comorbidity Index[27], maintenance hemodialysis, hepatic cancer, malignancy history, alcohol-related disease, and past varix rupture history. We also collected data regarding the use of antiplatelets, anticoagulants, nonsteroidal anti-inflammatory drugs, corticosteroids, and acid blockers prescribed on the day of admission or the following day or as part of the regular medications of the patient. The antiplatelet drugs used included aspirin, ticlopidine, clopidogrel, prasugrel, and ticagrelor. The anticoagulants prescribed included warfarin, dabigatran, edoxaban, rivaroxaban, apixaban, and heparin. Laboratory data collected on the day of admission included total bilirubin (mg/dL), aspartate aminotransferase (U/L), alanine aminotransferase (U/L), albumin (g/dL), white blood cells (/ μ L), hemoglobin (g/dL), platelets (10^3 / μ L), C-reactive protein (mg/dL), prothrombin time percentage (PT, %), activated partial thromboplastin time (APTT, sec), and estimated glomerular filtration rate (eGFR, mL/min/1.73 m²). Additionally, the shock index[28,29] was evaluated based on the vital signs at the hospital visit. The use of vasopressors and red blood cell transfusion volume on the day of admission were also obtained. Age was classified into four categories: < 65 , 65-4, 75-84, and ≥ 85 years. The Barthel Index was categorized into three groups: 0 (worst disability), 1-99, and 100 (full ability). Albumin and PT% were categorized according to the Child-Pugh score, while APTT was categorized into < 40 , 40-60, and ≥ 60 sec groups. All variables, excluding the Child-Pugh classification, were used as confounders in the analysis.

The primary outcome was a composite of 6-wk mortality, 4-wk rebleeding, and 4-wk onset of SBP. We defined rebleeding as cases when patients underwent endoscopic hemostasis procedures, such as EVL, EIS, or endoscopic clip hemostasis, two or more days after admission. To ensure outcome accuracy, all hemostatic procedures were verified by an endoscopy specialist using electronic medical records. Hemostatic procedures not associated with active bleeding but instead performed for future bleeding prevention, such as EVL or EIS on other varices and argon plasma coagulation, were excluded. SBP was defined as a polymorphonuclear cell count of 250/ μ L or greater[7,30], resulting from an ascites puncture performed during hospitalization. The secondary outcomes were the individual assessments of 6-wk mortality, 4-wk rebleeding, and 4-wk onset of SBP each assessed individually and in-hospital mortality. Also included were the 4-wk onset of clostridium difficile infection (CDI) and the length of hospital stay. CDI was defined as a diagnosis of ICD-10 code A047 on the second day of hospitalization or later and patients who were administered metronidazole or oral vancomycin.

Statistical methods

Continuous variables are reported as median and interquartile range (IQR), and categorical variables are reported as numbers and percentage. We determined the average treatment effect on the treated-based inverse probability of treatment weighting (IPTW) for the prophylaxis and non-prophylaxis groups. This method minimizes the effects of selection bias and imbalances in patient backgrounds between groups[31,32]. We estimated the propensity scores using logistic regression with prophylaxis as the dependent variable and all covariates as independent variables. Balances in baseline variables were also examined using standardized MD (SMD), and absolute values $< 10\%$ were considered balanced[33]. We used logistic regression to evaluate odds ratios (ORs) with 95% confidence intervals (CIs) to assess the

Table 1 List of antibiotics included in the study

ATC code	Type of antibiotics
J01AA	Tetracyclines
J01BA	Amphenicols
J01BB	Macrolides, lincosamides and streptogramins
J01CA	Penicillins with extended spectrum
J01CE	Beta-lactamase sensitive penicillins
J01CF	Beta-lactamase resistant penicillins
J01CR	Combinations of penicillins, including beta-lactamase inhibitors
J01DB	First-generation cephalosporins
J01DC	Second-generation cephalosporins
J01DD	Third-generation cephalosporins
J01DE	Fourth-generation cephalosporins
J01DF	Monobactams
J01DH	Carbapenems
J01EA	Trimethoprim and derivatives
J01EB	Short-acting sulfonamides
J01EC	Intermediate-acting sulfonamides
J01ED	Long-acting sulfonamides
J01EE	Combinations of sulfonamides
J01FA	Macrolides
J01FF	Lincosamides
J01FG	Streptogramins
J01GA	Aminoglycoside antibacterials
J01GB	Other aminoglycosides
J01MA	Fluoroquinolones
J01MB	Other quinolone antibacterials
J01XA	Glycopeptide antibacterials
J01XB	Polymyxins
J01XC	Steroid antibacterials
J01XD	Imidazole derivatives
J01XE	Nitrofurans derivatives
J01XX	Other antibacterials

ATC: Anatomical Therapeutic Chemical.

outcomes for categorical variables. The length of hospital stay was evaluated using negative binomial regression with rate ratios and 95%CI. The two-sided significance level for all tests was set at $P < 0.05$.

For the subgroup analysis, we evaluated the interaction effect between antibiotic prophylaxis and the Child-Pugh classification on the primary composite outcome. We employed logistic regression with IPTW, consistent with our primary analysis approach, using a dataset derived from multiple imputation data.

We conducted several sensitivity analyses to determine the robustness of our inferences. First, we performed both propensity score matching (PSM) to evaluate the robustness of the results. For PSM, we used the same propensity scores estimated for IPTW. A one-to-one PSM was conducted utilizing the nearest neighbor method without replacement. The caliper width was set at 20% of the standard deviation of the propensity scores on the logit scale. Second, considering the absence of a clear consensus on the duration of antibiotic prophylaxis, we narrowed the exposure period to those who received antibiotics for 2 d, 3 d, and 4 or more days. For these analyses, the exposure timing, definition of the control group, and analysis methods were identical to those used in the main analysis. Third, there is no consensus regarding the

appropriate type of antibiotic for prophylaxis. Therefore, to investigate the potential differences in outcomes due to the type of antibiotic used, we conducted a similar analysis with only third-generation cephalosporins that have a relatively large amount of evidence as the exposure[8].

In this study, we handled missing data by making a missing at-random assumption and conducting multiple imputations. These multiple imputations were conducted using chained equations with 100 imputed datasets and 200 iterations (maxit = 200) for each dataset. The imputation models included all the variables of interest and relevant auxiliary variables. Pooled estimates were obtained by combining the results across the imputed datasets, according to Rubin's rules[33,34].

All analyses were performed using R software (version 4.2.3; R Foundation for Statistical Computing, Vienna, Austria).

Sample size calculation

Based on previous reports, we assumed the incidence of the composite outcome to be 15%[11], and the antibiotic administration rate to be > 30%[16], expecting an unexposed to exposure ratio of approximately 2:1. We set a clinically meaningful risk ratio of 0.5 that was clinically meaningful for the composite outcomes associated with antibiotic prophylaxis[11, 35]. With an α error of 0.05 and a power of 80%, using Kelsey's equation, the required sample size was calculated to be 687 cases. Our sample size became larger than the predefined sample size, as described in the Results section.

Ethics

This study was conducted in accordance with the Declaration of Helsinki. It was reviewed and approved by the Institutional Review Board of the Future Medical Research Centre Ethical Committee (Approval Number: No. TGE02100-02). Due to the observational nature of the study, where patient data were accessed from hospital medical records without taking biological samples from patients, informed patient consent was deemed not necessary. Instead, an opt-out method was used and provided on the website of each hospital. This study is based on the Strengthening the Reporting of Observational Studies in Epidemiology guidelines.

RESULTS

A total of 980 patients from 46 hospitals who met the inclusion criteria were considered for inclusion in this study (Table 2). After applying the exclusion criteria, 790 patients were included in the analyses (Figure 1). The patients were divided into the prophylaxis ($n = 232$) and non-prophylaxis ($n = 558$) groups. Antibiotic prophylaxis was administered in 29.4% of patients. Most patients were male, under 65 years of age, and had a moderate level of functional independence (Table 3). The prevalence of alcohol-related diseases, varix rupture history, and β -blocker usage were higher in the prophylaxis group. Other variables, including the Child-Pugh score and the Charlson Comorbidity Index, were similar between the groups. The antibiotics used in the prophylaxis group included four carbapenems, 32 first-generation cephalosporins, 51 s-generation cephalosporins, 106 third-generation cephalosporins, 14 beta-lactamase inhibitor combinations, 22 macrolides, and three lincosamides (Table 4). The mean duration of administration was 4.59 d.

We ensured that the baseline conditions for the analysis were appropriately met. Figure 2 illustrates the overlap of the propensity scores for each group within one of the imputed datasets. The average C-statistic across the imputed datasets was 0.64. A comparison of patient characteristics before and after IPTW, as indicated by SMD, is outlined in Table 3 and Figure 3. Upon the application of IPTW, a balanced equivalence in the baseline characteristics was achieved between the groups.

Table 5 presents the outcomes before and after adjustment using IPTW. Before the application of IPTW, the composite outcome was 11.2% in the prophylaxis group and 9.5% in the non-prophylaxis group; the 6-wk mortality was 6.9% in the prophylaxis group and 6.6% in the non-prophylaxis group; the 4-wk rebleeding was 3.9% in the prophylaxis group and 2.9% in the non-prophylaxis group; the 4-wk onset of SBP was 2.2% in the prophylaxis group and 1.8% in the non-prophylaxis group; and the in-hospital mortality was 6.0% in the prophylaxis group and 6.1% in the non-prophylaxis group. There was one case of CDI in each group (0.4% in the prophylaxis group and 0.2% in the non-prophylaxis group). The median length of hospital stay was 8 d (IQR: 5-15 d) in the prophylaxis group and 9 d (IQR: 6-15 d) in the non-prophylaxis group.

Upon adjustment with IPTW, no significant differences regarding the composite outcome (adjusted OR, 1.11; 95%CI, 0.61-1.99; $P = 0.74$), 6-wk mortality (adjusted OR, 0.97; 95%CI, 0.47-1.98; $P = 0.93$), 4-wk rebleeding (adjusted OR, 1.21; 95%CI, 0.45-3.24; $P = 0.71$), 4-wk onset of SBP (adjusted OR, 1.20; 95%CI, 0.32-4.46; $P = 0.78$), or in-hospital mortality (adjusted OR, 0.89; 95%CI, 0.42-1.87; $P = 0.75$) were observed between the groups. The length of hospital stay did not significantly differ between the groups (adjusted rate ratio, 1.06; 95%CI, 0.94-1.19; $P = 0.34$).



In the subgroup analysis, there was no significant interaction between antibiotic prophylaxis and the Child-Pugh classification in relation to the composite outcome (P for interaction = 0.32) (Table 6). The sensitivity analyses of the PSM results and antibiotic duration were consistent with the main analysis (Table 7, Figures 4 and 5).

DISCUSSION

This long-term observational study involving data from 46 acute care hospitals across Japan explored the effectiveness of antibiotic prophylaxis for esophageal variceal bleeding treated with EVL. No benefits of antibiotic prophylaxis in terms of

Table 2 Distribution of facilities and cases across regions in Japan

Region	Number of facilities	Number of cases
Hokkaido	3	34
Tohoku	2	5
Kanto	14	319
Chubu	5	32
Kansai	11	359
Chugoku/Shikoku	1	2
Kyushu/Okinawa	10	229

**Figure 1** Patient flow.**Figure 2** Overlap of the propensity score of each group.

composite outcomes, individual outcomes, or length of hospital stay were identified. The effectiveness of prophylactic antibiotics in terms of composite outcomes were not significantly affected by the Child-Pugh classification.

Our findings underscore the diminishing role of universal prophylactic antibiotic administration in modern medical settings, aligning more closely with post-2010 results rather than older data. In previous randomized controlled trials (RCTs) regarding variceal bleeding that were conducted until the early 2000s, early mortality ranged from 4.2%-24%, rebleeding from 12.5%-20.8%, and the incidence of infections from 15-27.5%[35-39]. In contrast, only one RCT reported after 2010 reported early mortality and rebleeding rates of 3% and 8.5%, though the infection incidence was not assessed [13]. In this study, the 6-wk mortality was 6.7%, 4-wk rebleeding was 3.2%, and 4-wk onset of SBP was 1.9%, highlighting the improving treatment outcomes. Although a 2022 systematic review advocated for the benefits of antibiotic prophylaxis[15], it included one RCT published after 2010. The majority of studies reported after 2010 are single-center observa-

Table 3 Patient characteristics, missing data, and comparison of standardized mean differences, *n* (%)

	Before Imputation and IPTW			After imputation and IPTW	
	Prophylactic groups	Non-prophylactic groups	Missing (%)	SMD	SMD
Variables	<i>n</i> = 232	<i>n</i> = 558			
Age, yr			0	0.18	0.01
< 65	143 (61.6)	322 (57.7)			
65-74	46 (19.8)	144 (25.8)			
75-84	39 (16.8)	75 (13.4)			
≥ 85	4 (1.7)	17 (3.0)			
Sex, male (%)	181 (78.0)	417 (74.7)	0	0.08	< 0.01
Barthel index (%)			17.3	0.10	< 0.01
100 (full activity)	83 (40.5)	186 (41.5)			
1-99	63 (30.7)	152 (33.9)			
0 (worst disability)	59 (28.8)	110 (24.6)			
Child-Pugh score, median (IQR)	8 (7-10)	8 (7-10)	12.9	0.05	0.03
Child-Pugh classification (%)			10.1	0.06	0.02
A	42 (19.4)	93 (18.8)			
B	110 (50.9)	266 (53.7)			
C	64 (29.6)	136 (27.5)			
Presence of ascites	67 (31.0)	171 (34.5)			
Comorbidities					
Charlson Comorbidity Index, median (IQR)	4 (4-5)	4 (4-5)	0	< 0.01	0.01
Maintenance hemodialysis	3 (1.3)	8 (1.4)	0	0.01	< 0.01
Hepatic cancer	38 (16.4)	112 (20.1)	0	0.10	< 0.01
Malignant tumor history	29 (12.5)	65 (11.6)	0	0.03	< 0.01
Alcohol-related disease	127 (54.7)	246 (44.1)	0	0.21	< 0.01
Past varix rupture history	63 (27.2)	127 (22.8)	0	0.10	< 0.01
Medications					
Antiplatelet use	3 (1.3)	8 (1.4)	0	0.01	< 0.01
Anticoagulant use	5 (2.2)	8 (1.4)	0	0.05	< 0.01
NSAIDs use	5 (2.2)	13 (2.3)	0	0.01	< 0.01
Corticosteroid use	0 (0)	2 (0.4)	0	0.09	< 0.01
Acid blocker use	214 (91.8)	486 (87.1)	0	0.15	< 0.01
β blocker use	26 (11.2)	26 (4.7)	0	0.24	< 0.01
Laboratory data					
Total bilirubin, mg/dL, median (IQR)	1.6 (1-2.9)	1.4 (0.9-2.4)	3.2	0.13	0.02
Aspartate aminotransferase, U/L, median (IQR)	54.5 (32.2-94.8)	47 (31-83)	2.2	0.09	0.01
Alanine aminotransferase, U/L, median (IQR)	30.5 (20-47)	27 (19-42)	2.2	0.08	< 0.01
Albumin			4.6	0.11	0.02
> 3.5 g/dL	37 (16.5)	72 (13.6)			

2.8–3.5 g/dL	97 (43.3)	256 (48.3)			
< 2.8 g/dL	90 (40)	202 (38.1)			
White blood cell, / μ L, median (IQR)	7720 (5900–10700)	7300 (5200–10400)	1.8	0.15	0.01
Hemoglobin, g/dL, median (IQR)	9 (7.3–10.5)	8.5 (6.9–10.2)	1.8	0.12	< 0.01
Platelet, 10^3 / μ L, median (IQR)	99 (72–139)	103 (75–144)	1.8	< 0.01	0.02
C-reactive protein, mg/dL, median (IQR)	0.3 (0.1–0.7)	0.3 (0.1–0.8)	4.7	0.03	0.04
Prothrombin time			5.9	0.02	< 0.01
> 70%	48 (21.5)	113 (21.7)			
40%–70%	138 (61.9)	324 (62.3)			
< 40%	37 (16.6)	83 (16.0)			
Activated partial thromboplastin time			11.4	0.08	0.01
≤ 40 s	185 (87.3)	436 (89.3)			
40–60 s	23 (10.8)	47 (9.6)			
> 60 s	4 (1.9)	5 (1.0)			
eGFR < 30 mL/min/1.73 m ²	19 (8.3)	40 (7.3)	1.9	0.03	< 0.01
Shock index > 1	94 (41.4)	197 (36.5)	3	0.10	< 0.01
Vasopressor use	7 (3.0)	19 (3.4)	0	0.02	< 0.01
RBC transfusion, Unit, median (IQR)	4 (0–4)	2.5 (0–4)	0	0.09	< 0.01

IPTW: Inverse probability of treatment weighting; SMD: Standardized mean difference; IQR: Interquartile range; eGFR: Glomerular filtration rate; RBC: Red blood cell.

Table 4 Antibiotic use in prophylaxis group

Antibiotic class	Number of patients
Carbapenems	4
First-Generation Cephalosporins	32
Second-Generation Cephalosporins	51
Third-Generation Cephalosporins	106
Beta-Lactamase Inhibitor Combinations	14
Macrolides	22
Lincosamides	3

tional studies, indicating a lack of strong evidence supporting the routine use of antibiotics prophylaxis in contemporary settings.

The outcomes of our study can be understood through several underlying factors. The predominant role of EVL in hemostasis may have played a significant role in our findings. EVL results in fewer complications compared to EIS and offers superior control over bleeding[7–10,40]. While several previous RCTs incorporated EIS into their hemostatic protocols[35–39], both the current investigation and the most recent RCT focused exclusively on EVL[13]. This shift in technique may have contributed to a reduced incidence of complications, such as infections, suggesting that the need for antibiotic prophylaxis may be less pronounced when EVL is conducted. Additionally, the exclusion criteria of this study provides context. Severe patients, including those requiring mechanical ventilation, CRRT, IVR, or surgery, may have an inherent increased need for antibiotics. By design, our study did not include these patients. When patients that have effectively undergone hemostasis using EVL are included and critically ill patients are excluded, prophylactic antibiotics may not be as crucial as previously reported.

In recent years, a study evaluating the effectiveness of antibiotic prophylaxis for patients with cirrhosis presenting with upper gastrointestinal bleeding was conducted in Japan using a large-scale database[41]. In this study, the rate of antibiotic prophylaxis was 11.5%. Although the target was upper gastrointestinal bleeding and not esophageal variceal bleeding, it is evident that prophylactic antibiotics are not typically administered to patients with cirrhosis in Japan. Similar to our findings, their study did not demonstrate the utility of prophylactic antibiotic administration[41]. It may be necessary to reconsider the use of antibiotic prophylaxis for patients with cirrhosis in current medical settings.

Table 5 Crude and inverse probability of treatment weighting outcomes, *n* (%)

Outcomes	Before imputation and IPTW		After imputation and IPTW		
	Prophylaxis group (<i>n</i> = 232)	Non-prophylaxis group (<i>n</i> = 558)	Odds ratio (95%CI)	Odds ratio (95%CI)	<i>P</i> value
Composite outcome	26 (11.2)	53 (9.5)	1.20 (0.72-1.96)	1.11 (0.61-1.99)	0.74
6-wk mortality	16 (6.9)	37 (6.6)	1.04 (0.55-1.88)	0.97 (0.47-1.98)	0.93
4-wk rebleeding	9 (3.9)	16 (2.9)	1.37 (0.57-3.08)	1.21 (0.45-3.24)	0.71
4-wk onset of SBP	5 (2.2)	10 (1.8)	1.21 (0.37-3.44)	1.20 (0.32-4.46)	0.78
In-hospital mortality	14 (6.0)	34 (6.1)	0.99 (0.50-1.84)	0.89 (0.42-1.87)	0.75
			Rate ratio (95%CI)	Rate ratio (95%CI)	
Length of hospital, median (IQR)	8 (5-15)	9 (6-15)	1.01 (0.90-1.14)	1.06 (0.94-1.19)	0.34

IPTW: Inverse probability of treatment weighting; CI: Confidence interval; IQR: Interquartile range.

Table 6 Outcomes of subgroup analysis

Child-Pugh classification	Odds ratio (95%CI)	<i>P</i> value	<i>P</i> for interaction
A	0.87 (0.22-3.34)	0.84	0.32
B	0.79 (0.46-1.38)	0.41	
C	1.91 (1.20-3.02)	0.01	

Table 7 Outcomes of sensitivity analysis

Analysis method	Odds ratio (95%CI)	<i>P</i> value
IPTW	1.11 (0.61-1.99)	0.74
Propensity score matching	1.12 (0.62-2.03)	0.71
Duration of antibiotics		
2 d or more	1.14 (0.59-2.20)	0.70
3 d or more	1.05 (0.50-2.21)	0.91
4 d or more	0.96 (0.43-2.18)	0.93
Third-generation cephalosporins only	1.57 (0.64-3.87)	0.33

CI: Confidence interval; IPTW: Inverse probability of treatment weighting.

Our study's findings, revealing no significant benefit of antibiotic prophylaxis in patients undergoing EVL for esophageal variceal bleeding, add to the critical discourse on the necessity of routine prophylactic antibiotics in an era marked by escalating antibiotic resistance. The burgeoning concern for multidrug-resistant organisms (MDROs), highlighted in recent studies, is a pressing global health issue[42-44]. While our study did not directly address the intricate challenge of MDROs' emergence, the results imply that indiscriminate antibiotic use might not offer additional advantages and may, in fact, exacerbate the threat of antibiotic resistance. Consequently, our findings support a prudent reevaluation of antibiotic prophylaxis practices, especially in clinical environments where MDRO prevalence is high, and the risk of fostering resistance is a significant worry.

Patient groups for whom prophylactic antibiotic administration is beneficial must be identified. In our study, we demonstrated only the average effect across the population, showing that antibiotic prophylaxis is not effective. Previous reports have indicated differences in effectiveness based on the severity of the Child-Pugh classification. Although we conducted a subgroup analysis evaluating the interaction effect between antibiotic prophylaxis and the Child-Pugh classification on the primary composite outcome, we did not observe any significant results. Machine learning models are currently being used to identify heterogenous effects of antibiotic prophylaxis[45-47]. Using such methods, patients who would benefit from antibiotic prophylaxis must be identified.




Figure 3 Comparison of standardized mean difference before and after inverse probability of treatment weighting. IPTW: Inverse probability of treatment weighting; RBC: Red blood cell; eGFR: Glomerular filtration rate; NSAIDs: Nonsteroidal anti-inflammatory drugs.

Our study has several strengths. First, focusing on esophageal varix bleeding, this study was conducted on an unprecedented scale and comprised a wide sample of patients from multiple hospitals throughout various regions in Japan, bolstering the generalizability of our results. Second, the Tokushukai medical database offered us unique access to detailed blood test data, vital signs, and the ability to review electronic medical records in-depth. This enabled us to conduct a study with heightened precision.

Limitations

However, this study is not without limitations. First, due to the observational nature of this study, potential unmeasured confounding factors may be present. Second, the study is based on data from Japanese individuals, which limits the ability to generalize these findings to other populations or races. Third, the study encompasses only hemostasis information resulting from EVL. Patients who received treatment solely through EIS, balloon tamponade, or pharmacological interventions, such as somatostatin and vasopressors, were excluded. Fourth, treatment approaches differ notably between Japan and other countries. In Japan, a transjugular intrahepatic portosystemic shunt is not covered by insurance; therefore, no patients in our study received this treatment.

CONCLUSION

Our extensive multicenter observational study did not find a significant benefit to antibiotic prophylaxis for esophageal variceal bleeding treated with EVL. These results suggest that the recommendation for routine prophylactic antibiotic administration may not be universally essential. With growing concerns regarding the misuse of antibiotics and the consequential emergence of multidrug-resistant bacteria combined with advances in the management of esophageal variceal bleeding and liver cirrhosis treatment, there is a compelling need for a global reassessment of the necessity of routine antibiotic prophylaxis.




Figure 4 Comparison of standardized mean difference before and after propensity score matching. RBC: Red blood cell; eGFR: Glomerular filtration rate; NSAIDs: Nonsteroidal anti-inflammatory drugs; PTW: Inverse probability of treatment weighting.




Figure 5 Forest plot of sensitivity analysis. IPTW: Inverse probability of treatment weighting.

ARTICLE HIGHLIGHTS

Research background

Esophageal variceal bleeding is a critical complication of liver cirrhosis, typically managed with endoscopic variceal ligation (EVL). While current Western guidelines advocate antibiotic prophylaxis post-EVL, the evolving landscape of cirrhosis management and the rise of multidrug-resistant bacteria necessitate a reevaluation of this practice.

Research motivation

This study was motivated by the need to reassess the effectiveness of routine antibiotic prophylaxis following EVL in the context of improved cirrhosis treatments and increasing concerns regarding antibiotic resistance. Understanding the real-

world impact of prophylaxis on patient outcomes may result in a more effective and judicious use of antibiotics.

Research objectives

The primary objective was to evaluate the effectiveness of antibiotic prophylaxis in patients undergoing EVL for esophageal variceal bleeding using data from multiple Japanese medical centers. The study aimed to provide evidence that could influence future guideline recommendations and clinical practice.

Research methods

A 13-year observational study was conducted, using the Tokushukai medical database that includes data from 46 hospitals. Patients were categorized into prophylaxis and non-prophylaxis groups, with outcomes measured in terms of mortality, rebleeding, and spontaneous bacterial peritonitis. Logistic regression, inverse probability of treatment weighting, subgroup, and sensitivity analyses were conducted.

Research results

The study included 790 patients, and the primary outcomes were not significantly different between the prophylaxis and non-prophylaxis groups. These findings persisted across various subgroups and sensitivity analyses, suggesting that routine antibiotic prophylaxis post-EVL may not be beneficial.

Research conclusions

These findings challenge the current standard of prescribing antibiotics following EVL for esophageal variceal bleeding. They highlight the need for a global reassessment of this practice, considering the minimal impact on patient outcomes and the broader context of antibiotic resistance.

Research perspectives

Future research should focus on personalized approaches to antibiotic use in cirrhosis-related procedures, considering patient-specific factors and broader public health concerns. Further studies should also explore alternative strategies for managing complications in patients with liver cirrhosis.

ACKNOWLEDGEMENTS

We would like to express our gratitude to the gastroenterologists at the Tokushukai Hospitals throughout Japan who supported the treatment of esophageal variceal bleeding by providing emergency care without exception. We also express our gratitude to the staff at the Tokushukai Information Systems for their cooperation in extracting data and tracing back the electronic medical records.

FOOTNOTES

Author contributions: Ichita C contributed to the planning, data gathering, literature review, writing, and editing of the manuscript; Shimizu S and Goto T are experts in epidemiological statistics and were responsible for the causal inference and analysis methods in this study; Haruki U is an expert in portal hypertension and provided appropriate advice in this field, whereas Itoh N is an expert in infectious diseases and offered appropriate guidance on antibiotics; Iwagami M is a leading expert in the use of the Tokushukai medical database and provided appropriate advice on its utilization; Sasaki A is an expert in endoscopy and provided valuable advice on endoscopic hemostasis; all the authors reviewed the various drafts of the manuscript and have approved the final version of the manuscript.

Institutional review board statement: The study was reviewed and approved by the Institutional Review Board of the Future Medical Research Centre Ethical Committee (Approval No. TGE02100-02).

Informed consent statement: Owing to the observational nature of the study, where patient data were accessed from hospital medical records without taking biological samples from patients, informed patient consent was not deemed to be necessary. Instead, an opt-out method directed at patients was employed on the website of each hospital.

Conflict-of-interest statement: All other authors have nothing to disclose.

Data sharing statement: Due to privacy and ethical considerations, the data supporting the findings of this study is not publicly available. However, the study protocol and analysis code can be made available upon contacting the corresponding author. As for the participant data, ethical approval is required for access. The corresponding author can facilitate this process upon receipt of an appropriate request. After ethical approval, data sharing will be possible. The process of accessing the data will be carried out in accordance with ethical guidelines, ensuring respect for participant privacy and confidentiality.

STROBE statement: The authors have read the STROBE Statement-checklist of items, and the manuscript was prepared and revised according to the STROBE Statement-checklist of items.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Japan

ORCID number: Chikamasa Ichita 0000-0001-9210-7371; Sayuri Shimizu 0000-0003-0661-1171; Tadahiro Goto 0000-0002-5880-2968; Uojima Haruki 0000-0003-1719-1352; Naoya Itoh 0000-0001-6269-7884; Masao Iwagami 0000-0001-7079-0640; Akiko Sasaki 0000-0003-4219-554X.

S-Editor: Qu XL

L-Editor: A

P-Editor: Cai YX

REFERENCES

- Garcia-Tsao G, Bosch J. Management of varices and variceal hemorrhage in cirrhosis. *N Engl J Med* 2010; **362**: 823-832 [PMID: 20200386 DOI: 10.1056/NEJMra0901512]
- de Franchis R. Updating consensus in portal hypertension: report of the Baveno III Consensus Workshop on definitions, methodology and therapeutic strategies in portal hypertension. *J Hepatol* 2000; **33**: 846-852 [PMID: 11097497 DOI: 10.1016/s0168-8278(00)80320-7]
- Lee YY, Tee HP, Mahadeva S. Role of prophylactic antibiotics in cirrhotic patients with variceal bleeding. *World J Gastroenterol* 2014; **20**: 1790-1796 [PMID: 24587656 DOI: 10.3748/wjg.v20.i7.1790]
- Wong F, Bernardi M, Balk R, Christman B, Moreau R, Garcia-Tsao G, Patch D, Soriano G, Hoefs J, Navasa M; International Ascites Club. Sepsis in cirrhosis: report on the 7th meeting of the International Ascites Club. *Gut* 2005; **54**: 718-725 [PMID: 15831923 DOI: 10.1136/gut.2004.038679]
- Goulis J, Patch D, Burroughs AK. Bacterial infection in the pathogenesis of variceal bleeding. *Lancet* 1999; **353**: 139-142 [PMID: 10023916 DOI: 10.1016/S0140-6736(98)06020-6]
- Yoshiji H, Nagoshi S, Akahane T, Asaoka Y, Ueno Y, Ogawa K, Kawaguchi T, Kurosaki M, Sakaida I, Shimizu M, Tanai M, Terai S, Nishikawa H, Hiasa Y, Hidaka H, Miwa H, Chayama K, Enomoto N, Shimosegawa T, Takehara T, Koike K. Evidence-based clinical practice guidelines for Liver Cirrhosis 2020. *J Gastroenterol* 2021; **56**: 593-619 [PMID: 34231046 DOI: 10.1007/s00535-021-01788-x]
- European Association for the Study of the Liver. European Association for the Study of the Liver. EASL Clinical Practice Guidelines for the management of patients with decompensated cirrhosis. *J Hepatol* 2018; **69**: 406-460 [PMID: 29653741 DOI: 10.1016/j.jhep.2018.03.024]
- Tripathi D, Stanley AJ, Hayes PC, Patch D, Millson C, Mehrzad H, Austin A, Ferguson JW, Olliff SP, Hudson M, Christie JM; Clinical Services and Standards Committee of the British Society of Gastroenterology. U.K. guidelines on the management of variceal haemorrhage in cirrhotic patients. *Gut* 2015; **64**: 1680-1704 [PMID: 25887380 DOI: 10.1136/gutjnl-2015-309262]
- Garcia-Tsao G, Abraldes JG, Berzigotti A, Bosch J. Portal hypertensive bleeding in cirrhosis: Risk stratification, diagnosis, and management: 2016 practice guidance by the American Association for the study of liver diseases. *Hepatology* 2017; **65**: 310-335 [PMID: 27786365 DOI: 10.1002/hep.28906]
- de Franchis R, Bosch J, Garcia-Tsao G, Reiberger T, Ripoll C; Baveno VII Faculty. Baveno VII - Renewing consensus in portal hypertension. *J Hepatol* 2022; **76**: 959-974 [PMID: 35120736 DOI: 10.1016/j.jhep.2021.12.022]
- Chavez-Tapia NC, Barrientos-Gutierrez T, Tellez-Avila F, Soares-Weiser K, Mendez-Sanchez N, Gluud C, Uribe M. Meta-analysis: antibiotic prophylaxis for cirrhotic patients with upper gastrointestinal bleeding - an updated Cochrane review. *Aliment Pharmacol Ther* 2011; **34**: 509-518 [PMID: 21707680 DOI: 10.1111/j.1365-2036.2011.04746.x]
- Chavez-Tapia NC, Barrientos-Gutierrez T, Tellez-Avila FI, Soares-Weiser K, Uribe M. Antibiotic prophylaxis for cirrhotic patients with upper gastrointestinal bleeding. *Cochrane Database Syst Rev* 2010; **2010**: CD002907 [PMID: 20824832 DOI: 10.1002/14651858.CD002907.pub2]
- Lee TH, Huang CT, Lin CC, Chung CS, Lin CK, Tsai KC. Similar rebleeding rate in 3-day and 7-day intravenous ceftriaxone prophylaxis for patients with acute variceal bleeding. *J Formos Med Assoc* 2016; **115**: 547-552 [PMID: 26899745 DOI: 10.1016/j.jfma.2016.01.006]
- Zullo A, Soncini M, Bucci C, Marmo R; Gruppo Italiano per lo Studio dell'Emorragia Digestiva (GISED) (Appendix). Clinical outcomes in cirrhotics with variceal or nonvariceal gastrointestinal bleeding: A prospective, multicenter cohort study. *J Gastroenterol Hepatol* 2021; **36**: 3219-3223 [PMID: 34189770 DOI: 10.1111/jgh.15601]
- Wong YJ, Tan CK, Yui YL, Wong Y, Tam YC, Chan E, Abraldes JG. Antibiotic prophylaxis in cirrhosis patients with upper gastrointestinal bleeding: An updated systematic review and meta-analysis. *PH and C* 2022; **1**: 167-177 [DOI: 10.1002/poh2.35]
- Ueno M, Kayahara T, Sunami T, Takayama H, Takabatake H, Morimoto Y, Yamamoto H, Mizuno M. Universal antibiotic prophylaxis may no longer be necessary for patients with acute variceal bleeding: A retrospective observational study. *Medicine (Baltimore)* 2020; **99**: e19981 [PMID: 32443300 DOI: 10.1097/MD.00000000000019981]
- Ueno M, Mano T, Kayahara T, Mizuno M. Antibiotic prophylaxis for cirrhotic patients with upper gastrointestinal bleeding: Is evidence adequate? *J Gastroenterol Hepatol* 2021; **36**: 3249-3250 [PMID: 34368981 DOI: 10.1111/jgh.15652]
- Chang TS, Tsai YH, Lin YH, Chen CH, Lu CK, Huang WS, Yang YH, Chen WM, Hsieh YY, Wu YC, Tung SY, Huang YH. Limited effects of antibiotic prophylaxis in patients with Child-Pugh class A/B cirrhosis and upper gastrointestinal bleeding. *PLoS One* 2020; **15**: e0229101 [PMID: 32084186 DOI: 10.1371/journal.pone.0229101]
- Tandon P, Abraldes JG, Keough A, Bastiampillai R, Jayakumar S, Carbonneau M, Wong E, Kao D, Bain VG, Ma M. Risk of Bacterial Infection in Patients With Cirrhosis and Acute Variceal Hemorrhage, Based on Child-Pugh Class, and Effects of Antibiotics. *Clin Gastroenterol Hepatol* 2015; **13**: 1189-96.e2 [PMID: 25460564 DOI: 10.1016/j.cgh.2014.11.019]
- Fernández J, Piano S, Bartoletti M, Wey EQ. Management of bacterial and fungal infections in cirrhosis: The MDRO challenge. *J Hepatol*

- 2021; **75** Suppl 1: S101-S117 [PMID: [34039482](#) DOI: [10.1016/j.jhep.2020.11.010](#)]
- 21 **Iwagami M**, Moriya H, Doi K, Yasunaga H, Isshiki R, Sato I, Mochida Y, Ishioka K, Ohtake T, Hidaka S, Noiri E, Kobayashi S. Seasonality of acute kidney injury incidence and mortality among hospitalized patients. *Nephrol Dial Transplant* 2018; **33**: 1354-1362 [PMID: [29462342](#) DOI: [10.1093/ndt/gfy011](#)]
 - 22 **Hayashida K**, Murakami G, Matsuda S, Fushimi K. History and Profile of Diagnosis Procedure Combination (DPC): Development of a Real Data Collection System for Acute Inpatient Care in Japan. *J Epidemiol* 2021; **31**: 1-11 [PMID: [33012777](#) DOI: [10.2188/jea.JE20200288](#)]
 - 23 **Wang K**, Li P, Chen L, Kato K, Kobayashi M, Yamauchi K. Impact of the Japanese diagnosis procedure combination-based payment system in Japan. *J Med Syst* 2010; **34**: 95-100 [PMID: [20192060](#) DOI: [10.1007/s10916-008-9220-2](#)]
 - 24 **Shah S**, Vanclay F, Cooper B. Improving the sensitivity of the Barthel Index for stroke rehabilitation. *J Clin Epidemiol* 1989; **42**: 703-709 [PMID: [2760661](#) DOI: [10.1016/0895-4356\(89\)90065-6](#)]
 - 25 **Child CG**, Turcotte JG. Surgery and portal hypertension. *Major Probl Clin Surg* 1964; **1**: 1-85 [PMID: [4950264](#)]
 - 26 **Pugh RN**, Murray-Lyon IM, Dawson JL, Pietroni MC, Williams R. Transection of the oesophagus for bleeding oesophageal varices. *Br J Surg* 1973; **60**: 646-649 [PMID: [4541913](#) DOI: [10.1002/bjs.1800600817](#)]
 - 27 **Quan H**, Sundararajan V, Halfon P, Fong A, Burnand B, Luthi JC, Saunders LD, Beck CA, Feasby TE, Ghali WA. Coding algorithms for defining comorbidities in ICD-9-CM and ICD-10 administrative data. *Med Care* 2005; **43**: 1130-1139 [PMID: [16224307](#) DOI: [10.1097/01.mlr.0000182534.19832.83](#)]
 - 28 **Allgöwer M**, Burri C. ["Shock index"]. *Dtsch Med Wochenschr* 1967; **92**: 1947-1950 [PMID: [5299769](#) DOI: [10.1055/s-0028-1106070](#)]
 - 29 **Koch E**, Lovett S, Nghiem T, Riggs RA, Rech MA. Shock index in the emergency department: utility and limitations. *Open Access Emerg Med* 2019; **11**: 179-199 [PMID: [31616192](#) DOI: [10.2147/OAEM.S178358](#)]
 - 30 **Runyon BA**; AASLD Practice Guidelines Committee. Management of adult patients with ascites due to cirrhosis: an update. *Hepatology* 2009; **49**: 2087-2107 [PMID: [19475696](#) DOI: [10.1002/hep.22853](#)]
 - 31 **Austin PC**. An Introduction to Propensity Score Methods for Reducing the Effects of Confounding in Observational Studies. *Multivariate Behav Res* 2011; **46**: 399-424 [PMID: [21818162](#) DOI: [10.1080/00273171.2011.568786](#)]
 - 32 **Robins JM**, Hernán MA, Brumback B. Marginal structural models and causal inference in epidemiology. *Epidemiology* 2000; **11**: 550-560 [PMID: [10955408](#) DOI: [10.1097/00001648-200009000-00011](#)]
 - 33 **Rubin DB**. Multiple Imputation for Nonresponse in Surveys. Wiley Series in Probability and Statistics. 1987 [DOI: [10.1002/9780470316696](#)]
 - 34 **White IR**, Royston P, Wood AM. Multiple imputation using chained equations: Issues and guidance for practice. *Stat Med* 2011; **30**: 377-399 [PMID: [21225900](#) DOI: [10.1002/sim.4067](#)]
 - 35 **Jun CH**, Park CH, Lee WS, Joo YE, Kim HS, Choi SK, Rew JS, Kim SJ, Kim YD. Antibiotic prophylaxis using third generation cephalosporins can reduce the risk of early rebleeding in the first acute gastroesophageal variceal hemorrhage: a prospective randomized study. *J Korean Med Sci* 2006; **21**: 883-890 [PMID: [17043424](#) DOI: [10.3346/jkms.2006.21.5.883](#)]
 - 36 **Rolando N**, Gimson A, Philpott-Howard J, Sahathevan M, Casewell M, Fagan E, Westaby D, Williams R. Infectious sequelae after endoscopic sclerotherapy of oesophageal varices: role of antibiotic prophylaxis. *J Hepatol* 1993; **18**: 290-294 [PMID: [8228122](#) DOI: [10.1016/s0168-8278\(05\)80272-7](#)]
 - 37 **Selby WS**, Norton ID, Pokorny CS, Benn RA. Bacteremia and bacterascites after endoscopic sclerotherapy for bleeding esophageal varices and prevention by intravenous cefotaxime: a randomized trial. *Gastrointest Endosc* 1994; **40**: 680-684 [PMID: [7859964](#)]
 - 38 **Hong SN**, Kim BJ, Lee SY, Lee CY, Ryu MK, Choi MS, Lee JH, Rhee PL, Koh KC, Kim JJ, Paik SW, Rhee JC, Choi KW. [Prospective randomized trial of intravenous ciprofloxacin for prevention of bacterial infection in cirrhotic patients with esophageal variceal bleeding]. *Taehan Kan Hakhoe Chi* 2002; **8**: 288-296 [PMID: [12499786](#)]
 - 39 **Hou MC**, Lin HC, Liu TT, Kuo BI, Lee FY, Chang FY, Lee SD. Antibiotic prophylaxis after endoscopic therapy prevents rebleeding in acute variceal hemorrhage: a randomized trial. *Hepatology* 2004; **39**: 746-753 [PMID: [14999693](#) DOI: [10.1002/hep.20126](#)]
 - 40 **Lo GH**, Lai KH, Cheng JS, Chen MH, Chiang HT. A prospective, randomized trial of butyl cyanoacrylate injection versus band ligation in the management of bleeding gastric varices. *Hepatology* 2001; **33**: 1060-1064 [PMID: [11343232](#) DOI: [10.1053/jhep.2001.24116](#)]
 - 41 **Ueno M**, Fujiwara T, Tokumasu H, Mano T, Kayahara T, Takabatake H, Morimoto Y, Matsueda K, Fukuoka T, Mizuno M. Real-world efficacy of antibiotic prophylaxis for upper gastrointestinal bleeding in cirrhotic patients in Japan. *J Gastroenterol* 2023; **58**: 766-777 [PMID: [37171554](#) DOI: [10.1007/s00535-023-02000-y](#)]
 - 42 **Caron WP**, Mousa SA. Prevention strategies for antimicrobial resistance: a systematic review of the literature. *Infect Drug Resist* 2010; **3**: 25-33 [PMID: [21694891](#) DOI: [10.2147/idr.s10018](#)]
 - 43 **Morris S**, Cerceio E. Trends, Epidemiology, and Management of Multi-Drug Resistant Gram-Negative Bacterial Infections in the Hospitalized Setting. *Antibiotics (Basel)* 2020; **9** [PMID: [32326058](#) DOI: [10.3390/antibiotics9040196](#)]
 - 44 **Mücke VT**, Peiffer KH, Kessel J, Schwarzkopf KM, Bojunga J, Zeuzem S, Finkelmeier F, Mücke MM. Impact of colonization with multidrug-resistant organisms on antibiotic prophylaxis in patients with cirrhosis and variceal bleeding. *PLoS One* 2022; **17**: e0268638 [PMID: [35609050](#) DOI: [10.1371/journal.pone.0268638](#)]
 - 45 **Inoue K**, Seeman TE, Horwich T, Budoff MJ, Watson KE. Heterogeneity in the Association Between the Presence of Coronary Artery Calcium and Cardiovascular Events: A Machine-Learning Approach in the MESA Study. *Circulation* 2023; **147**: 132-141 [PMID: [36314118](#) DOI: [10.1161/CIRCULATIONAHA.122.062626](#)]
 - 46 **Inoue K**, Athey S, Tsugawa Y. Machine-learning-based high-benefit approach versus conventional high-risk approach in blood pressure management. *Int J Epidemiol* 2023; **52**: 1243-1256 [PMID: [37013846](#) DOI: [10.1093/ije/dyad037](#)]
 - 47 **Shiba K**, Daoud A, Hikichi H, Yazawa A, Aida J, Kondo K, Kawachi I. Uncovering Heterogeneous Associations Between Disaster-Related Trauma and Subsequent Functional Limitations: A Machine-Learning Approach. *Am J Epidemiol* 2023; **192**: 217-229 [PMID: [36255224](#) DOI: [10.1093/aje/kwac187](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

