

# World Journal of *Gastrointestinal Surgery*

*World J Gastrointest Surg* 2024 March 27; 16(3): 635-973



**EDITORIAL**

- 635 *Ex vivo* liver resection and auto-transplantation and special systemic therapy in perihilar cholangiocarcinoma treatment  
*Tchilikidi KY*
- 641 Indocyanine green: The guide to safer and more effective surgery  
*Fransvea P, Chiarello MM, Fico V, Cariati M, Brisinda G*

**MINIREVIEWS**

- 650 Alcohol associated liver disease and bariatric surgery: Current perspectives and future directions  
*Cooper KM, Colletta A, Hebda N, Devuni D*
- 658 Applications of gastric peroral endoscopic myotomy in the treatment of upper gastrointestinal tract disease  
*Chang SY, Jin GH, Sun HB, Yang D, Tang TY*

**ORIGINAL ARTICLE****Retrospective Cohort Study**

- 670 Evaluation of bacterial contamination and medium-term oncological outcomes of intracorporeal anastomosis for colon cancer: A propensity score matching analysis  
*Kayano H, Mamuro N, Kamei Y, Ogimi T, Miyakita H, Nakagohri T, Koyanagi K, Mori M, Yamamoto S*
- 681 Rescue from complications after pancreaticoduodenectomies at a low-volume Caribbean center: Value of tailored peri-pancreatectomy protocols  
*Cawich SO, Dixon E, Shukla PJ, Shrikhande SV, Deshpande RR, Mohammed F, Pearce NW, Francis W, Johnson S, Bujhawan J*
- 689 Comparison of prognosis and postoperative morbidities between standard pancreaticoduodenectomy and the TRIANGLE technique for resectable pancreatic ductal adenocarcinoma  
*Hang HX, Cai ZH, Yang YF, Fu X, Qiu YD, Cheng H*
- 700 Analysis of the impact of immunotherapy efficacy and safety in patients with gastric cancer and liver metastasis  
*Liu K, Wu CX, Liang H, Wang T, Zhang JY, Wang XT*

**Retrospective Study**

- 710 Clinical observation of extraction-site incisional hernia after laparoscopic colorectal surgery  
*Fan BH, Zhong KL, Zhu LJ, Chen Z, Li F, Wu WF*
- 717 Predicting short-term major postoperative complications in intestinal resection for Crohn's disease: A machine learning-based study  
*Wang FT, Lin Y, Yuan XQ, Gao RY, Wu XC, Xu WW, Wu TQ, Xia K, Jiao YR, Yin L, Chen CQ*

- 731 Analysis of factors impacting postoperative pain and quality of life in patients with mixed hemorrhoids: A retrospective study  
*Sun XW, Xu JY, Zhu CZ, Li SJ, Jin LJ, Zhu ZD*
- 740 Pre-operative visceral adipose tissue radiodensity is a potentially novel prognostic biomarker for early endoscopic post-operative recurrence in Crohn's disease  
*Gu P, Dube S, Gellada N, Choi SY, Win S, Lee YJ, Yang S, Haritunians T, Melmed GY, Vasiliauskas EA, Bonthala N, Syal G, Yarur AJ, Ziring D, Rabizadeh S, Fleshner P, Kallman C, Devkota S, Targan SR, Li D, McGovern DP*
- 751 Clinical study on the relationship between liver cirrhosis, ascites, and hyponatremia  
*Li XJ, Meng HH*
- 759 Comparison of the clinical effects of dual-modality endoscopy and traditional laparotomy for the treatment of intra- and extrahepatic bile duct stones  
*Wang W, Xia H, Dai B*
- 768 Role of ablation therapy in conjunction with surgical resection for neuroendocrine tumors involving the liver  
*Ostapenko A, Stroever S, Eyasu L, Kim M, Aploks K, Dong XD, Seshadri R*
- 777 Feasibility and safety of minimally invasive multivisceral resection for T4b rectal cancer: A 9-year review  
*Chan KS, Liu B, Tan MNA, How KY, Wong KY*
- 790 MH-STRALP: A scoring system for prognostication in patients with upper gastrointestinal bleeding  
*Hu JN, Xu F, Hao YR, Sun CY, Wu KM, Lin Y, Zhong L, Zeng X*
- Clinical Trials Study**
- 807 Early postoperative complications after transverse colostomy closure, a retrospective study  
*Liu F, Luo XJ, Li ZW, Liu XY, Liu XR, Lv Q, Shu XP, Zhang W, Peng D*
- 816 Clinical study of enhanced recovery after surgery in laparoscopic appendectomy for acute appendicitis  
*Li ZL, Ma HC, Yang Y, Chen JJ, Wang ZJ*
- Observational Study**
- 823 Reinforced tissue matrix to strengthen the abdominal wall following reversal of temporary ostomies or to treat incisional hernias  
*Lake SP, Deeken CR, Agarwal AK*
- Randomized Controlled Trial**
- 833 Whole-process case management effects on mental state and self-care ability in patients with liver cancer  
*Ju MD, Qin Q, Li M*
- Clinical and Translational Research**
- 842 Construction and validation of somatic mutation-derived long non-coding RNAs signatures of genomic instability to predict prognosis of hepatocellular carcinoma  
*Duan BT, Zhao XK, Cui YY, Liu DZ, Wang L, Zhou L, Zhang XY*

**Basic Study**

- 860 Influence of different magnetic forces on the effect of colonic anastomosis in rats  
*Tian BY, Zhang MM, Ma J, Lyu Y, Yan XP*
- 871 Inflammatory responses in esophageal mucosa before and after laparoscopic antireflux surgery  
*Ergun P, Kipcak S, Selvi Gunel N, Yildirim Sozmen E, Bor S*
- 882 Etanercept-synthesizing adipose-derived stem cell secretome: A promising therapeutic option for inflammatory bowel disease  
*Kim SJ, Kim OH, Hong HE, Ju JH, Lee DS*

**SYSTEMATIC REVIEWS**

- 893 Impact of frailty on short-term postoperative outcomes in patients undergoing colorectal cancer surgery: A systematic review and meta-analysis  
*Zhou Y, Zhang XL, Ni HX, Shao TJ, Wang P*

**META-ANALYSIS**

- 907 Endoscopic-ultrasound-guided biliary drainage with placement of electrocautery-enhanced lumen-apposing metal stent for palliation of malignant biliary obstruction: Updated meta-analysis  
*Peng ZX, Chen FF, Tang W, Zeng X, Du HJ, Pi RX, Liu HM, Lu XX*
- 921 Clinical efficacy and safety of erlotinib combined with chemotherapy in the treatment of advanced pancreatic cancer: A meta-analysis  
*Liu XY, Pan HN, Yu Y*

**CASE REPORT**

- 932 Link between mutations in *ACVRL1* and *PLA2G4A* genes and chronic intestinal ulcers: A case report and review of literature  
*Tang YJ, Zhang J, Wang J, Tian RD, Zhong WW, Yao BS, Hou BY, Chen YH, He W, He YH*
- 944 Mucinous neoplasm of the appendix: A case report and review of literature  
*Chang HC, Kang JC, Pu TW, Su RY, Chen CY, Hu JM*
- 955 Abdominal cocoon syndrome-a rare culprit behind small bowel ischemia and obstruction: Three case reports  
*Vipudhamorn W, Juthasilaparut T, Sutharat P, Sanmee S, Supatrakul E*
- 966 Endoscopic ultrasound-guided lauromacrogol injection for treatment of colorectal cavernous hemangioma: Two case reports  
*Zhu HT, Chen WG, Wang JJ, Guo JN, Zhang FM, Xu GQ, Chen HT*

**ABOUT COVER**

Editorial Board Member of *World Journal of Gastrointestinal Surgery*, Jia-Gang Han, MD, Professor, Department of General Surgery, Beijing Chaoyang Hospital, Capital Medical University, Beijing 100020, China. [hjg211@163.com](mailto:hjg211@163.com)

**AIMS AND SCOPE**

The primary aim of *World Journal of Gastrointestinal Surgery* (*WJGS, World J Gastrointest Surg*) is to provide scholars and readers from various fields of gastrointestinal surgery with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

*WJGS* mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal surgery and covering a wide range of topics including biliary tract surgical procedures, biliopancreatic diversion, colectomy, esophagectomy, esophagostomy, pancreas transplantation, and pancreatectomy, *etc.*

**INDEXING/ABSTRACTING**

The *WJGS* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJGS* as 2.0; IF without journal self cites: 1.9; 5-year IF: 2.2; Journal Citation Indicator: 0.52; Ranking: 113 among 212 journals in surgery; Quartile category: Q3; Ranking: 81 among 93 journals in gastroenterology and hepatology; and Quartile category: Q4.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Zi-Hang Xu, Production Department Director: Xiang Li, Editorial Office Director: Jia-Ru Fan.

**NAME OF JOURNAL**

*World Journal of Gastrointestinal Surgery*

**ISSN**

ISSN 1948-9366 (online)

**LAUNCH DATE**

November 30, 2009

**FREQUENCY**

Monthly

**EDITORS-IN-CHIEF**

Peter Schemmer

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/1948-9366/editorialboard.htm>

**PUBLICATION DATE**

March 27, 2024

**COPYRIGHT**

© 2024 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>



Randomized Controlled Trial

# Whole-process case management effects on mental state and self-care ability in patients with liver cancer

Man-Di Ju, Qin Qin, Meng Li

**Specialty type:** Gastroenterology and hepatology

**Provenance and peer review:** Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): 0  
Grade C (Good): C  
Grade D (Fair): 0  
Grade E (Poor): 0

**P-Reviewer:** Calvo H, Spain

**Received:** November 23, 2023

**Peer-review started:** November 23, 2023

**First decision:** December 18, 2023

**Revised:** December 22, 2023

**Accepted:** January 30, 2024

**Article in press:** January 30, 2024

**Published online:** March 27, 2024



**Man-Di Ju**, Department of Internal Medicine of Abdominal Oncology, Hubei Cancer Hospital, Wuhan 430070, Hubei Province, China

**Qin Qin**, Department of Hepatological Surgery, Wuhan No. 1 Hospital, Wuhan 430000, Hubei Province, China

**Meng Li**, Department of Nursing, General Hospital of the Yangtze River Shipping, Wuhan 430014, Hubei Province, China

**Corresponding author:** Meng Li, Nurse, Department of Nursing, General Hospital of the Yangtze River Shipping, No. 1 Huiji Road, Jiang'an District, Wuhan 430014, Hubei Province, China. [limeng333321@163.com](mailto:limeng333321@163.com)

## Abstract

### BACKGROUND

Regarding the incidence of malignant tumors in China, the incidence of liver cancer ranks fourth, second only to lung, gastric, and esophageal cancers. The case fatality rate ranks third after lung and cervical cancer. In a previous study, the whole-process management model was applied to patients with breast cancer, which effectively reduced their negative emotions and improved treatment adherence and nursing satisfaction.

### AIM

To explore Mental state and self-care ability in patients with liver cancer: effects of whole-process case management.

### METHODS

In this single-center, randomized, controlled study, 60 randomly selected patients with liver cancer who had been admitted to our hospital from January 2021 to January 2022 were randomly divided into an observation group ( $n = 30$ ), who received whole-process case management on the basis of routine nursing measures, and a control group ( $n = 30$ ), who were given routine nursing measures. We compared differences between the two groups in terms of anxiety, depression, the level of hope, self-care ability, symptom distress, sleep quality, and quality of life.

### RESULTS

Post-intervention, Hamilton anxiety scale, Hamilton depression scale, memory symptom assessment scale, and Pittsburgh sleep quality index scores in both

groups were lower than those pre-intervention, and the observation group had lower scores than the control group ( $P < 0.05$ ). Herth hope index, self-care ability assessment scale-revision in Chinese, and quality of life measurement scale for patients with liver cancer scores in both groups were higher than those pre-intervention, with higher scores in the observation group compared with the control group ( $P < 0.05$ ).

## CONCLUSION

Whole-process case management can effectively reduce anxiety and depression in patients with liver cancer, alleviate symptoms and problems, and improve the level of hope, self-care ability, sleep quality, and quality of life, as well as provide feasible nursing alternatives for patients with liver cancer.

**Key Words:** Liver cancer; Mental state; Self-care ability; Whole-process case management; Life quality; Nursing

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Patients with liver cancer are likely to face different dilemmas and have differing psychological states at various stages of the disease. A definitive diagnosis of liver cancer can be a significant trauma for patients, with patients and their families often unable to cope for a certain period of time. Patients with liver cancer are prone to negative psychological emotions such as anxiety, depression, and fear. The whole-process case management was shown to effectively reduce the anxiety and depression in patients with liver cancer, reduce symptom distress and improve the level of hope, self-care ability, sleep quality and life quality, as well as provide feasible nursing alternatives for patients with liver cancer.

**Citation:** Ju MD, Qin Q, Li M. Whole-process case management effects on mental state and self-care ability in patients with liver cancer. *World J Gastrointest Surg* 2024; 16(3): 833-841

**URL:** <https://www.wjgnet.com/1948-9366/full/v16/i3/833.htm>

**DOI:** <https://dx.doi.org/10.4240/wjgs.v16.i3.833>

## INTRODUCTION

In terms of global cancer rankings, liver cancer ranks sixth in incidence and second in terms of case fatality rate. In males, the incidence of liver cancer ranks fifth and the case fatality rate ranks second, while the incidence of liver cancer in women ranks ninth and the case fatality rate ranks sixth[1]. Regarding the incidence of malignant tumors in China, the incidence of liver cancer ranks fourth, second only to lung, gastric, and esophageal cancers. The case fatality rate ranks third after lung and cervical cancer. The incidence and mortality rates for liver cancer in males with malignant tumors are both ranked third. The incidence rate in females with malignant tumors is ranked sixth, and the case fatality rate is ranked fourth[2]. The incidence and mortality rates for liver cancer in China is higher than those of other countries, accounting for > 50% of the total number of patients with liver cancer globally[3,4].

Patients with liver cancer are likely to face different dilemmas and have differing psychological states at various stages of the disease. A definitive diagnosis of liver cancer can be a significant trauma for patients, with patients and their families often unable to cope for a certain period of time. Patients with liver cancer are prone to negative psychological emotions such as anxiety, depression, and fear. During treatment, patients are known to experience complex and diverse psychological dilemmas[5]. Moreover, physical pain, a long treatment process, medical treatment dilemmas, complications, and the threat of death may result in patients already physical and mental stress being unable to cope[6]. Furthermore, the cost of treatment is high, repeated hospitalizations are often required, and work and social life are affected. A patient's financial income is likely to be reduced, resulting in increased financial pressure on a family, and patients often have feelings of guilt. The incidence of liver cancer has been reported to peak between the ages of 40 and 60 years[7] and patients in this age group are frequently providing fundamental financial and emotional support within their families. Following the onset of liver cancer, the resulting stress may trigger psychological issues that are challenging to overcome, especially for male patients in China. Historically, clinical treatment for liver cancer has involved medical staff paying attention more often to surgery or drug treatment for patients with cancer, while overlooking or underestimating changes in the mental status of patients with cancer. It is important to investigate nursing intervention programs that are suitable for improving the mental status of patients with liver cancer.

The whole-process management model is a type of high-quality nursing service that continues throughout the entire diagnosis and treatment process of patients, that is, patients can enjoy continuous professional nursing services from admission to discharge and from discharge to review, as well as receive rehabilitation[8]. The United States and other Western countries have taken the lead in proposing whole-process nursing management and applying it to clinical nursing practice. Following promotion and development, whole-process case nursing has been found not to increase patients' medical costs, but rather to improve nursing efficiency and reduce medical costs[9]. In a previous study, the whole-process management model was applied to patients with breast cancer, which effectively reduced their negative emotions and improved treatment adherence and nursing satisfaction[8]. However, the application of this model to patients with liver cancer has rarely been reported. Therefore, this study aimed to explore the effects of the application of

whole-process case management on the mental state and self-care ability of patients with liver cancer.

## MATERIALS AND METHODS

### Study patients

In this single-center, randomized controlled study, 60 patients with liver cancer who had been admitted to our hospital from January 2021 to January 2022 were randomly selected. The patients were randomly divided into an observation group ( $n = 30$ ) and a control group ( $n = 30$ ). Inclusion criteria comprised patients: Aged  $\geq 18$  years, who met the clinical diagnostic criteria for liver cancer[10], and who were fully conscious, with good cognitive function, and fully able to communicate. Exclusion criteria comprised patients: With a prior history of mental illness, with other serious systemic diseases, with major negative events having occurred in recent life, who were unable to perform independent self-care, and with other malignant tumors. Included study patients or their family members signed a written informed consent form to participate in the study. All processes and procedures undertaken during this study were in accordance with the Declaration of Helsinki.

### Nursing programs

The control group was supported with routine nursing measures, including routine examination during hospitalization, vital signs testing, timely medical intervention, perioperative care, relevant medication information, routine health education, and bi-weekly telephone follow-up post-discharge.

The observation group received whole-process case management interventions in addition to routine nursing measures. Specifically, the interventions included full-time nurses organizing a seminar to broaden discussions concerning liver-cancer related information. The intervention steps were as follows:

A case management file was established for each patient to provide a continuous and dynamic medical record.

Patient-related issues were analyzed through communicating with patients in the observation group to become better informed concerning their physical and psychological issues, doubts, and confusions in relation to their daily care, and their health behaviors were analyzed to identify practical challenges for discussion.

A nursing plan was proposed. A consensus was reached following discussion and a standardized nursing plan was designed, and health education was conducted once a patient had been admitted. The nursing staff listened and replied to each patient's consultation content in a timely manner and provided evidence-based guidance. Intensive education was delivered concerning the ward environment, physical and psychological nursing was provided to reinforce the information provided by medical staff, and the care of patients with liver cancer was standardized during hospitalization, especially during the perioperative period, with a view to preventing common complications.

During hospitalization, a psychologist regularly evaluated each patient's mental status and adjusted psychological interventions in real time according to the assessment results.

Medical staff guided and educated patients while implementing nursing to strengthen patient self-management ability. Pathology-related information specific to liver cancer and perioperative self-intervention methods for patients were introduced. Adverse symptoms that a patient might encounter such as pain, hematemesis, and ascites were explained, and patients were informed of the corresponding countermeasures. Nursing staff positively confirmed patient self-management behaviors in relation to pain, discomfort, or complications.

At discharge, in accordance with the needs of each patient, the nursing staff formulated and implemented work plans for discharge preparation, including home self-care, referral to other hospitals, remote health management, and home follow-up. Personalized electronic files were established for patients throughout the course of the disease. A health management manual was issued to patients to facilitate recording their medication and psychological status outside the hospital. This manual also contained relevant patient information concerning liver cancer and frequently asked questions and responses. Patients were required to bring the manual to follow-up consultations to facilitate more accurate understanding of the current status of each patient by the medical team.

Post-discharge: Follow-up visits were conducted through telephone or Internet platforms such as WeChat, and personal nursing education after discharge was continued, as necessary. A specific WeChat forum was set up with a continuous service team and training was provided for team members. Through the establishment of a WeChat forum, timely team communication, discussion, and a timely response to patient-related questions could be realized. Through continuous and regular assessment of a patient's mental state *via* telephone or online platforms, and according to the assessment results, professional psychologists provided targeted psychological guidance to patients. Patients were regularly informed of new information concerning liver cancer *via* the WeChat platform. Each patient's weekly feedback information was collated and analyzed, and new content was re-introduced. The intervention was provided continuously for three months post-discharge.

### Evaluation indicators

**Anxiety and depression:** Anxiety was measured using the Hamilton anxiety scale (HAMA)[11] and depressive mood was measured using the Hamilton depression scale (HAMD)[12]. The HAMA has a total of 14 questions, involving a five-level scoring method ranging from 0 to 4 points, with a total score between 0 and 56 points. The higher the score, the more serious the anxiety. The HAMD has a total of 17 questions, involving a five-level scoring method ranging from 0 to 4 points, with a total score between 0 and 68 points. The higher the score, the more serious the depression.

**Level of hope:** The Herth hope index (HHI)[13] was used to assess a patient's level of hope. In total, 12 items involving three aspects, namely, attitude, action, and social relationships were assessed with each item ranged from 1 to 4 points, with a total possible score of 48 points. The higher the score, the higher the level of hope.

**Self-care ability:** Self-care ability was assessed using the self-care ability assessment scale- revision in Chinese (ASAS-R-C). The self-care ability assessment scale, first designed in 2014, was used to form the ASAS-R-C scale with a stable structure and clear dimensions[14]. The ASAS-R-C scale has three dimensions, namely, general self-care ability, developmental self-care ability, and self-care ability when personal health is poor. Using a Likert five-point scale scoring method, it includes 15 items (items 4, 11, 14, and 15 are reverse scored), with scores ranging from 15 to 75 points. The higher the score, the stronger the self-care ability.

**Symptom distress:** We used the memory symptom assessment scale (MSAS)[15], developed at the Memorial Sloan-Kettering Cancer Center in the United States, which is a multi-dimensional symptom self-rating scale consisting of three dimensions, namely, physiological symptoms, psychological symptoms and total distress index. It includes 32 items, of which the first 24 symptom-related items evaluate the symptoms experienced by patients with malignant tumors in terms of incidence, frequency, severity, and distress. Eight further symptom items are used to evaluate the frequency and severity of each symptom item, with scores represented in terms of points ranging from 1 to 4 (low to high), and the degree of distress represented in terms of points ranging from 0 to 4 (low to high). Frequency, severity, and life-distressing scores for each symptom item are added and divided by 3 to calculate the average score of the symptom. This scale had a Cronbach's  $\alpha$  coefficient of 0.79-0.87 and a content validity of 0.94, and the Chinese version of this scale had good reliability and validity.

**Sleep quality:** Patient sleep quality was scored using the Pittsburgh sleep quality index (PSQI)[16]. The scale has 24 questions divided into seven categories. The scores of each category range from 0 to 3 points, with 0 point indicating there are no problems and 3 points indicating major difficulties. The total score is obtained by adding the scores of each category and can range between 0 and 21 points. The higher the score, the poorer the sleep quality.

**Quality of life:** Quality of life was assessed using the quality of life measurement scale for patients with liver cancer (QLICP-LI)[17]. This scale, consisting of a general module (version 2.0) and a liver cancer-specific model, includes 32 items covering four areas, namely, physical function, psychological function, social function, and common symptoms and side-effects. It also includes eight items covering basic physiological function and mobility and activity ability; nine items covering cognition, emotion and will; and eight items covering interpersonal communication, social support and security, and social role. Common symptoms and side-effects are assessed using seven items. The liver cancer-specific module uses 12 items to assess liver-related signs, systemic manifestations, and specific psychological effects. In total, this scale comprises 13 aspects and 44 items across five areas, with each item ranging from 1 to 5 points and the total score ranging from 44 to 220 points. The higher the score, the better the quality of life.

### Statistical methods

SPSS22.0 statistical software was used to analyze the data. Following a normality test, the measurement data approximately followed a normal distribution (expressed as mean  $\pm$  SD). Counting data are represented as percentages. We conducted a *t*-test, a Pearson's chi-square test, a Fisher's exact test or a Mann-Whitney test, as required. *P* values < 0.05 were considered statistically significant.

## RESULTS

### Patient baseline data

In total, 60 patients with liver cancer (average age, 50.16  $\pm$  9.37 years; males, 66.67%; females, 33.33%; Child-Pugh grades: Grade A, 65%; grade B, 35.00%) were included in our study. In terms of education levels, 45% of the patients had at least completed junior high school, 36.67% had completed senior high school, and 18.33% had completed university. In terms of tumor-node-metastasis (TNM) stages, 8.33% of the patients had stage I liver cancer, 70.00% had stage II, and 21.67% had stage III. In addition, 15% of the patients had hepatocellular carcinoma, 68.33% had adenocarcinoma, 11.67% had squamous cell carcinoma, and 5.00% had liver cancer of other pathological types.

Compared with the control group, no statistically significant differences were observed in baseline data concerning age, sex, Child-Pugh grade, education level, pathological type, TNM stage, and other general data in the observation group (*P* > 0.05; Table 1).

### Comparisons between anxiety, depression, and level of hope

Pre-intervention, compared with the control group, no statistically significant differences were observed in the observation group concerning the HAMA, HAMD, and HHI scores (*P* > 0.05). Post-intervention, the HAMA and HAMD scores in both groups were significantly lower than those pre-intervention, and observation group scores were significantly lower than those in the control group (*P* < 0.05). Compared with pre-intervention scores, the HHI score in both groups was significantly higher, and observation group scores were significantly higher than those in the control group (*P* < 0.05; Table 2).

**Table 1 Comparison of baseline data between the two groups [n (%), mean ± SD]**

Item	Observation group (n = 30)	Control group (n = 30)	t/ $\chi^2$	P value
Age (yr)	49.48 ± 9.25	50.83 ± 9.60	0.555	0.581
Sex			0.300	0.583
Male	19 (63.33)	21 (70.00)		
Female	11 (36.67)	9 (30.00)		
Child-Pugh grade			0.073	0.786
Grade A	20 (66.67)	19 (63.33)		
Grade B	10 (33.33)	11 (36.67)		
Education level			0.584	0.559
Junior high school or lower	14 (43.34)	13 (46.67)		
Senior high school	12 (33.33)	10 (40.00)		
University or higher	4 (23.33)	7 (13.33)		
Pathological type			0.611	0.893
Hepatocellular carcinoma	4 (13.33)	5 (16.67)		
Adenocarcinoma	20 (66.67)	21 (70.00)		
Squamous cell carcinoma	4 (13.33)	3 (10.00)		
Others	2 (6.67)	1 (3.33)		
TNM stage			0.027	0.978
I	2 (6.67)	3 (10.00)		
II	22 (73.33)	20 (66.67)		
III	6 (20.00)	7 (23.33)		

TNM: Tumor-node-metastasis.

**Table 2 Comparison of anxiety and depression plus hope level between two groups (mean ± SD)**

Item	Observation group (n = 30)	Control group (n = 30)	t/ $\chi^2$	P value
HAMA				
Pre-intervention	20.44 ± 6.57	19.83 ± 5.82	0.384	0.702
Post-intervention	8.15 ± 2.38 <sup>a</sup>	10.09 ± 2.10 <sup>a</sup>	3.359	0.001
HAMD				
Pre-intervention	22.52 ± 6.63	22.98 ± 6.46	0.274	0.784
Post-intervention	9.45 ± 1.61 <sup>a</sup>	11.93 ± 2.33 <sup>a</sup>	4.776	< 0.001
HHI				
Pre-intervention	20.98 ± 5.51	21.31 ± 6.01	0.225	0.822
Post-intervention	34.07 ± 5.40 <sup>a</sup>	30.86 ± 5.39 <sup>a</sup>	2.304	0.024

<sup>a</sup>P < 0.05.

Note: Compared with pre-intervention. HAMA: Hamilton anxiety scale; HAMD: Hamilton depression scale; HHI: Herth hope index.

### Comparison of self-care ability

Pre-intervention, compared with the control group, no statistically significant difference was observed in the ASAS-R-C scores in the observation group ( $P > 0.05$ ). Post-intervention, the ASAS-R-C score was significantly higher in both groups compared with pre-intervention scores, and observation group scores were significantly higher than those in the control group ( $P < 0.05$ ; Table 3).

**Table 3 Comparison of self-care abilities assessment scale-revision in Chinese between two groups (mean ± SD)**

Item	n	Pre-intervention	Post-intervention	t	P value
Observation group	30	42.07 ± 8.42	58.55 ± 5.46	9.757	< 0.001
Control group	30	41.79 ± 7.12	53.94 ± 7.73	6.682	< 0.001
t		0.133	2.665		
P value		0.894	0.009		

### Comparison of symptom distress

Pre-intervention, compared with the control group, no statistically significant difference was observed in each dimension of the MSAS scores in the observation group ( $P > 0.05$ ). Post-intervention, the MSAS scores in each dimension in the two groups were significantly lower than those in each group pre-intervention, and observation group scores were significantly lower than those in the control group ( $P < 0.05$ ; Table 4).

### Comparison of sleep quality and quality of life

Pre-intervention, no statistically significant difference in PSQI scores was observed in the observation group compared with the control group ( $P > 0.05$ ). Post-intervention, the PSQI scores of both groups were significantly lower than those in each group pre-intervention, and observation group scores were significantly lower than those in the control group ( $P < 0.05$ ). Compared with each group pre-intervention, the QLICP-LI scores in both groups were significantly higher post-intervention, with observation group scores significantly higher post-intervention than those in the control group ( $P < 0.05$ ; Table 5).

## DISCUSSION

The onset age of liver cancer reaches a peak at 45 years, with males having a higher risk of developing liver cancer than females[18,19]. The clinical treatment for liver cancer currently mainly involves interventional treatment such as radiotherapy, chemotherapy, surgical resection, or Transcatheter arterial chemoembolization. While the above treatment methods have shown good efficacy, patients with liver cancer are under considerable physical and mental pressure during the treatment process. Some patients have severe depression, anxiety, and other negative emotions, and a poor mental state. Furthermore, the self-management ability of patients with cancer in China is of a medium level, and issues such as insufficient self-management awareness, lack of knowledge and information, and poor disease management ability have been reported[20-22]. These issues have an important effect on quality of life, treatment decisions, and a desire to survive.

The whole-process case management model provides a more comprehensive medical service compared with the general hospitalization mode. The main highlight of the whole-process case management model is managed participation in the entire treatment course, which can also be regarded as an extension of the hospitalization routine treatment outside the hospital[23,24].

In this study, the HAMA, HAMD, MSAS, and PSQI scores of both groups post-intervention were lower than pre-intervention scores, with the observation group having lower scores than the control group ( $P < 0.05$ ). The HHI, ASAS-R-C, and QLICP-LI scores were higher in both groups compared to their respective baseline values. However, the observation group had significantly higher scores than the control group ( $P < 0.05$ ). Our findings indicate that the whole-process case management model could effectively reduce the anxiety and depression in patients with liver cancer, reduce symptom distress, and improve the level of hope, self-care ability, sleep quality and life quality.

The whole-process case management model established a relationship of mutual trust with patients in the early stage of treatment, with regular communication undertaken through various channels such as telephone and the WeChat platform, which provided patients with continuous confidence in treatment, encouragement when encountering difficulties, and facilitated responses to their questions. Patients could receive advice when they were facing life pressures, which helped them to overcome negative thoughts. The patients were asked to regularly record their information concerning their physical and psychological status, identify issues that were mostly causing them anxiety, and to set out recently completed treatment goals. These recordings allowed them to gain confidence in the treatment process and to improve their level of hope[25].

In addition, in the observation group, the whole-process case management service was implemented *via* the WeChat platform, which was convenient for medical staff to fully and dynamically learn about the self-care situation of patients outside of the hospital, and to provide timely adjustments to strengthen the content and direction of continuous care. For patients who could not use WeChat, nurses could guide family members to enter the WeChat platform and join the whole-process case management. In the process of home treatment, patients could complete remote health consultations *via* the network, doctors and nursing staff could communicate with patients in real time, and nursing staff could provide relevant knowledge in relation to liver cancer, precautions when taking medication, and preliminary treatment when discomfort or complications occurred to patients *via* the network platform. Patients carried out self-health management under professional supervision and guidance. This method has been shown to effectively reduce the effect of various

**Table 4 Comparison of symptom distress between the two groups (mean ± SD)**

Item	Observation group (n = 30)	Control group (n = 30)	t/χ <sup>2</sup>	P value
Symptom distress in physiology				
Pre-intervention	1.22 ± 0.53	1.25 ± 0.32	0.278	0.781
Post-intervention	0.82 ± 0.37 <sup>a</sup>	1.05 ± 0.28 <sup>a</sup>	2.668	0.009
Symptom distress in psychology				
Pre-intervention	1.67 ± 0.39	1.78 ± 0.30	1.280	0.205
Post-intervention	1.09 ± 0.32 <sup>a</sup>	1.29 ± 0.27 <sup>a</sup>	2.630	0.011
Distress index				
Pre-intervention	1.35 ± 0.42	1.44 ± 0.35	0.869	0.388
Post-intervention	0.96 ± 0.25 <sup>a</sup>	1.21 ± 0.23 <sup>a</sup>	3.863	< 0.003

<sup>a</sup>P < 0.05.

Note: Compared with pre-intervention.

**Table 5 Comparison of sleep quality and life quality (mean ± SD)**

Item	PSQI		QLICP-LI	
	Pre-intervention	Post-intervention	Pre-intervention	Post-intervention
Observation group (n = 30)	13.35 ± 4.51	7.25 ± 2.27 <sup>a</sup>	126.65 ± 17.07	165.89 ± 15.25 <sup>a</sup>
Control group (n = 30)	13.27 ± 4.59	8.72 ± 2.79 <sup>a</sup>	122.40 ± 16.67	145.05 ± 13.80 <sup>a</sup>
t/χ <sup>2</sup>	0.063	2.239	0.977	5.550
P value	0.949	0.029	0.332	< 0.001

<sup>a</sup>P < 0.05.

Note: Compared with pre-intervention. PSQI: Pittsburgh sleep quality index; QLICP-LI: Quality of life measurement scale for patients with liver cancer.

symptoms during hospitalization and following discharge in relation to the physical and mental health of patients, ensuring patient safety post-discharge[26].

This study had some limitations. First, this single-center clinical study involved only a small number of patients from the same hospital and within the same region, which may have led to selection bias. Second, this study only followed the patients for three months and long-term effects of the intervention were not observed. Third, nurses collaborate, coordinate, and execute whole nursing, which is key to ensuring the safety of nursing and improving the quality of nursing. While the nurses involved in the whole-process case management in this study were specialist nurses, there can be deficiencies in managing all aspects of the nursing process and implementing nursing intervention strategies, and the theoretical knowledge and operation technology may not be able to quickly meet the requirements of the whole-process case management model. The next step is to continue to expand the sample size, improve the study design, and encourage nursing staff to continuously improve their nursing skills to ensure effective implementation of the whole-process case management model in future studies.

## CONCLUSION

In summary, the whole-process case management was shown to effectively reduce the anxiety and depression in patients with liver cancer, reduce symptom distress and improve the level of hope, self-care ability, sleep quality and life quality, as well as provide feasible nursing alternatives for patients with liver cancer.

## ARTICLE HIGHLIGHTS

### **Research background**

In China, liver cancer incidence is fourth, following lung, gastric, and esophageal cancers. Case fatality rate is third after lung and cervical cancer. A previous study applied whole-process management model to breast cancer patients, effectively reducing negative emotions, improving treatment adherence, and nursing satisfaction.

### **Research motivation**

Improve the hope level, self-care ability, sleep quality and quality of life, and provide feasible care plan for liver cancer patients.

### **Research objectives**

To study the psychological state and self-care ability of liver cancer patients: the effect of the whole process of case management

### **Research methods**

We compared anxiety, depression, hope level, self-care ability, symptom distress, sleep quality, and quality of life between the two groups.

### **Research results**

Whole-course case management can effectively reduce anxiety and depression, relieve symptoms and problems.

### **Research conclusions**

Improve patient self-care ability and provide feasible nursing plan for liver cancer patients.

### **Research perspectives**

This is a research perspective of the whole-process case management.

---

## FOOTNOTES

**Co-first authors:** Man-Di Ju and Qin Qin.

**Author contributions:** Ju MD and Qin Q designed the research; Li M, Ju MD and Qin Q performed the research; Li M, Ju MD and Qin Q contributed new reagents/analytic tools; Li M, Ju MD and Qin Q analyzed the data; Ju MD and Qin Q wrote the paper; All authors were involved in the critical review of the results and have contributed to, read, and approved the final manuscript. Ju MD and Qin Q contributed equally to this work as co-first authors equally to this work. Ju MD and Qin Q contributed efforts of equal substance throughout the research process. The choice of these researchers as co-first authors acknowledges and respects this equal contribution, while recognizing the spirit of teamwork and collaboration of this study. In summary, we believe that designating Ju MD and Qin Q as co-first authors of are fitting for our manuscript as it accurately reflects our team's collaborative spirit, equal contributions, and diversity.

**Institutional review board statement:** This study protocol was approved by the General Hospital of the Yangtze River Shipping, and all the families have voluntarily participated in the study and have signed informed consent forms.

**Clinical trial registration statement:** We have not registered URL number. We have the ethical certificate approved by the unit.

**Informed consent statement:** All the families have voluntarily participated in the study and have signed informed consent forms.

**Conflict-of-interest statement:** All the authors report no relevant conflicts of interest for this article.

**Data sharing statement:** Data generated from this investigation are available upon reasonable request from the corresponding author at [limeng333321@163.com](mailto:limeng333321@163.com).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** China

**ORCID number:** Meng Li 0009-0005-1170-6134.

**S-Editor:** Li L

**L-Editor:** A

**P-Editor:** Yuan YY

## REFERENCES

- 1 **Sung H**, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, Bray F. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin* 2021; **71**: 209-249 [PMID: 33538338 DOI: 10.3322/caac.21660]
- 2 **Chen W**, Zheng R, Baade PD, Zhang S, Zeng H, Bray F, Jemal A, Yu XQ, He J. Cancer statistics in China, 2015. *CA Cancer J Clin* 2016; **66**: 115-132 [PMID: 26808342 DOI: 10.3322/caac.21338]
- 3 **Xia C**, Dong X, Li H, Cao M, Sun D, He S, Yang F, Yan X, Zhang S, Li N, Chen W. Cancer statistics in China and United States, 2022: profiles, trends, and determinants. *Chin Med J (Engl)* 2022; **135**: 584-590 [PMID: 35143424 DOI: 10.1097/CM9.0000000000002108]
- 4 **Cao W**, Chen HD, Yu YW, Li N, Chen WQ. Changing profiles of cancer burden worldwide and in China: a secondary analysis of the global cancer statistics 2020. *Chin Med J (Engl)* 2021; **134**: 783-791 [PMID: 33734139 DOI: 10.1097/CM9.0000000000001474]
- 5 **Zhang J**, Liu Y, Xu Y. Soothing liver-qi stagnation method for cancer-related depression: A protocol for systematic review and meta-analysis. *Medicine (Baltimore)* 2020; **99**: e22797 [PMID: 33120798 DOI: 10.1097/MD.00000000000022797]
- 6 **Kosciusko R**, Geller D, Kucinski B, Shadel W, Vanegas Y, Tsung A, Marsh W, Steel J. Substance Use, Depression, and Illness Perception Among Cancer Patients. *Am J Health Behav* 2019; **43**: 287-299 [PMID: 30808469 DOI: 10.5993/AJHB.43.2.6]
- 7 **Lin J**, Zhang H, Yu H, Bi X, Zhang W, Yin J, Zhao P, Liang X, Qu C, Wang M, Hu M, Liu K, Wang Y, Zhou Z, Wang J, Tan X, Liu W, Shao Z, Cai J, Tang W, Cao G. Epidemiological Characteristics of Primary Liver Cancer in Mainland China From 2003 to 2020: A Representative Multicenter Study. *Front Oncol* 2022; **12**: 906778 [PMID: 35800051 DOI: 10.3389/fonc.2022.906778]
- 8 **Jin L**, Zhao Y, Wang P, Zhu R, Bai J, Li J, Jia X, Wang Z. Efficacy of the Whole-Course Case Management Model on Compliance and Satisfaction of Breast Cancer Patients with Whole-Course Standardized Treatment. *J Oncol* 2022; **2022**: 2003324 [PMID: 35783153 DOI: 10.1155/2022/2003324]
- 9 **Chen J**, Bai H. Evaluation of Implementation Effect of Cervical Cancer Comprehensive Treatment Patients With Whole-Course High-Quality Care Combined With Network Continuation Care. *Front Surg* 2022; **9**: 838848 [PMID: 35174206 DOI: 10.3389/fsurg.2022.838848]
- 10 **Anwanwan D**, Singh SK, Singh S, Saikam V, Singh R. Challenges in liver cancer and possible treatment approaches. *Biochim Biophys Acta Rev Cancer* 2020; **1873**: 188314 [PMID: 31682895 DOI: 10.1016/j.bbcan.2019.188314]
- 11 **Ye Z**, Zhang Y, Du M, Lu S, Zhao Q, Yang S. The Correlation Between Probiotics and Anxiety and Depression Levels in Cancer Patients: A Retrospective Cohort Study. *Front Psychiatry* 2022; **13**: 830081 [PMID: 35432048 DOI: 10.3389/fpsy.2022.830081]
- 12 **Zhang S**, Chen H, Zhang M, Sun X, Liu X. Reduction of depression symptoms in laryngeal cancer patients receiving psychology services. *Am J Transl Res* 2020; **12**: 6637-6645 [PMID: 33194060]
- 13 **Li Y**, Ni N, Zhou Z, Dong J, Fu Y, Li J, Luan Z, Peng X. Hope and symptom burden of women with breast cancer undergoing chemotherapy: A cross-sectional study. *J Clin Nurs* 2021; **30**: 2293-2300 [PMID: 33756013 DOI: 10.1111/jocn.15759]
- 14 **Guo L**, Zauszniewski JA, Ding X, Zhang L, Gao H, Guo Q, Liu K. The Appraisal of Self-Care Agency Scale-Revised (ASAS-R): Reliability and Validity Among Older Chinese People. *West J Nurs Res* 2017; **39**: 1459-1476 [PMID: 27703077 DOI: 10.1177/0193945916672821]
- 15 **Llamas-Ramos I**, Llamas-Ramos R, Carrillo-González GM, Sepúlveda-Ramírez J, Vargas-Rosero E. Symptom Prevalence in Spanish and Colombian Oncology Patients Measured with the MSAS. *Cancers (Basel)* 2022; **14** [PMID: 35406396 DOI: 10.3390/cancers14071624]
- 16 **Hamzeh S**, Safari-Faramani R, Khatony A. Effects of Aromatherapy with Lavender and Peppermint Essential Oils on the Sleep Quality of Cancer Patients: A Randomized Controlled Trial. *Evid Based Complement Alternat Med* 2020; **2020**: 7480204 [PMID: 32308715 DOI: 10.1155/2020/7480204]
- 17 **Wang L**, Huang YE, Zhang H, Sun Y, Liu Y, Zhao L, Zhao Y, Xue L. Individualized whole-course management in prostatic puncture biopsy: Application and efficiency. *Zhonghua Nan Ke Xue* 2022; **28**: 608-611 [PMID: 37556218]
- 18 **Mattuzzi C**, Lippi G. Current Cancer Epidemiology. *J Epidemiol Glob Health* 2019; **9**: 217-222 [PMID: 31854162 DOI: 10.2991/jegh.k.191008.001]
- 19 **Liu Y**, Zheng J, Hao J, Wang RR, Liu X, Gu P, Yu H, Yu Y, Wu C, Ou B, Peng Z. Global burden of primary liver cancer by five etiologies and global prediction by 2035 based on global burden of disease study 2019. *Cancer Med* 2022; **11**: 1310-1323 [PMID: 35118819 DOI: 10.1002/cam4.4551]
- 20 **Allegrante JP**, Wells MT, Peterson JC. Interventions to Support Behavioral Self-Management of Chronic Diseases. *Annu Rev Public Health* 2019; **40**: 127-146 [PMID: 30601717 DOI: 10.1146/annurev-publhealth-040218-044008]
- 21 **Yin L**, Zhang W, Liu L, Guo L, Guo M, He X, Zhu L. Application of nursing intervention based on the IKAP model in self-management of patients with gastric cancer. *Am J Transl Res* 2022; **14**: 6389-6398 [PMID: 36247277]
- 22 **Zhu X**, Han S, Chu H, Wang M, Chen S. Influence of self-management exercise intervention on the cancer related fatigue severity and self-management efficacy of patients with non-small cell lung cancer after operation. *J Pak Med Assoc* 2020; **70**: 88-93 [PMID: 33177734]
- 23 **Pang L**, Wang Y, Xing Y, Zhao C. Application Effects of Whole Course High-Quality Nursing on Patients with Liver Cancer during Radiotherapy. *Iran J Public Health* 2019; **48**: 1777-1785 [PMID: 31850254]
- 24 **Qian X**, Gong L, Zhou F, Zhang Y, Wang H. High-Quality Nursing Combined with the Whole-Course Responsibility Nursing Intervention Reduces the Incidence of Complications in Severe Aneurysmal Subarachnoid Hemorrhage. *Evid Based Complement Alternat Med* 2022; **2022**: 3252718 [PMID: 35880108 DOI: 10.1155/2022/3252718]
- 25 **Zhang Q**, Wan R, Liu C. The impact of intense nursing care in improving anxiety, depression, and quality of life in patients with liver cancer: A systematic review and meta-analysis. *Medicine (Baltimore)* 2020; **99**: e21677 [PMID: 32846784 DOI: 10.1097/MD.00000000000021677]
- 26 **Chai Y**, Li L, Wu YL, Wang T, Jia YM, Lin XL, Chen X, Zhong H, Liu LX, Tao LD. The effects of case management for breast cancer patients: A protocol for systematic review and meta-analysis. *Medicine (Baltimore)* 2022; **101**: e28960 [PMID: 35244058 DOI: 10.1097/MD.00000000000028960]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA  
**Telephone:** +1-925-3991568  
**E-mail:** [office@baishideng.com](mailto:office@baishideng.com)  
**Help Desk:** <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>

