

World Journal of *Clinical Cases*

World J Clin Cases 2023 August 26; 11(24): 5628-5839



MINIREVIEWS

- 5628 Effect of pesticides on phosphorylation of tau protein, and its influence on Alzheimer's disease
Torres-Sánchez ED, Ortiz GG, Reyes-Urbe E, Torres-Jasso JH, Salazar-Flores J

ORIGINAL ARTICLE

Case Control Study

- 5643 Reduction rate of monoclonal protein as a useful prognostic factor in standard-risk group of newly diagnosed multiple myeloma
Liu M, Zhang JY

Retrospective Cohort Study

- 5653 Effectiveness of treating menorrhagia using microwave endometrial ablation at a frequency of 2.45 GHz
Kakinuma T, Kaneko A, Kakinuma K, Matsuda Y, Yanagida K, Takeshima N, Ohwada M
- 5660 Benefits of laparoscopy-assisted ileostomy in colorectal cancer patients with bowel obstruction
Wang YJ, Lin KH, Kang JC, Hu JM, Chen CY, Pu TW

Retrospective Study

- 5666 Hypopharyngeal cancer trends in a high-incidence region: A retrospective tertiary single center study
Cordunianu AGV, Ganea G, Cordunianu MA, Cochior D, Moldovan CA, Adam R
- 5678 Relevant detection indicator of prethrombotic state in patients with primary hypertension
Luo J, Yang T, Ding L, Xiong JH, Ying T, Xu F
- 5692 Clinical study of extrahepatic biliary adenoma
Li W, Tao J, Song XG, Hou MR, Qu K, Gu JT, Yan XP, Yao BW, Qin YF, Dong FF, Sha HC

SYSTEMATIC REVIEWS

- 5700 Sodium-glucose cotransporter-2 inhibitor-associated euglycemic diabetic ketoacidosis in COVID-19-infected patients: A systematic review of case reports
Khedr A, Hennawi HA, Khan MK, Eissa A, Mir M, Rauf I, Nitesh J, Surani S, Khan SA

META-ANALYSIS

- 5710 Efficacy and safety of Huangqi Jianzhong decoction in the treatment of chronic atrophic gastritis: A meta-analysis
Yan XP, Si W, Ding MS, Tian YF, Guo Z

CASE REPORT

- 5721** Malignant melanoma of the prostate: Primary or metastasis? A case report
Zhao H, Liu C, Li B, Guo JM
- 5729** Intravenous leiomyoma of the uterus extending to the pulmonary artery: A case report
Huang YQ, Wang Q, Xiang DD, Gan Q
- 5736** Percutaneous endoscopic necrosectomy for walled-off necrosis in the retroperitoneal space of the elderly: A case report
Sato K, Shibukawa G, Ueda K, Nakajima Y, Togashi K, Ohira H
- 5742** Acute exacerbation of idiopathic pulmonary fibrosis treated using the Feibi recipe: Two case reports
Liu ZH, Li GD, Hao QX, Cao F, Cheng Y, Kou MJ, Jiao Y
- 5749** Neonatal erythema multiforme associated with a rotavirus infection: A case report
Kim JJ, Lee JK
- 5755** Hemorrhagic Bartholin's cyst in a woman using anti-platelet medication: A case report and review of the literature
Li YR, Ding DC
- 5762** Subintimal recanalization for non-acute occlusion of intracranial vertebral artery in an emergency endovascular procedure: A case report
Fu JF, Zhang XL, Lee SY, Zhang FM, You JS
- 5772** Synchronous rectal adenocarcinoma and intestinal mantle cell lymphoma: A case report
Vu KV, Trong NV, Khuyen NT, Huyen Nga D, Anh H, Tien Trung N, Trung Thong P, Minh Duc N
- 5780** Focal lymphoblastic transformation of chronic myelogenous leukemia develops into erythroid leukemia: A case report
Wang W, Chen YL, Gou PP, Wu PL, Shan KS, Zhang DL
- 5789** Intraoperative sudden arrhythmias in cervical spine surgery adjacent to the stellate ganglion: A case report
Seo JH, Cho SY, Park JH, Seo JY, Lee HY, Kim DJ
- 5797** Papillary thyroid carcinoma with nodular fasciitis-like stroma - an unusual variant with distinctive histopathology: A case report
Hu J, Wang F, Xue W, Jiang Y
- 5804** Malignant form of hidroacanthoma simplex: A case report
Yang YF, Wang R, Xu H, Long WG, Zhao XH, Li YM
- 5811** Penile and scrotal strangulation by stainless steel rings in an human immunodeficiency virus positive man: A case report
Usuda D, Kaminishi N, Kato M, Sugawara Y, Shimizu R, Inami T, Tsuge S, Sakurai R, Kawai K, Matsubara S, Tanaka R, Suzuki M, Shimozawa S, Hotchi Y, Osugi I, Katou R, Ito S, Mishima K, Kondo A, Mizuno K, Takami H, Komatsu T, Oba J, Nomura T, Sugita M

- 5817** Persistent postoperative hypotension caused by subclinical empty sella syndrome after a simple surgery: A case report
Zhao KM, Hu JS, Zhu SM, Wen TT, Fang XM
- 5823** Rare *ROS1-CENPW* gene in pancreatic acinar cell carcinoma and the effect of crizotinib plus AG chemotherapy: A case report
Wang T, Shen YY
- 5830** Fecal transplantation in patient with metastatic melanoma refractory to immunotherapy: A case report
del Giglio A, Atui FC
- 5835** Left hepatic artery pseudoaneurysm complicating endoscopic retrograde cholangiopancreatography: A case report
Li QM, Ye B, Yang SW, Zhao H

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Kelser de Souza Kock, PhD, Physiotherapist, Professor, Department of Physiotherapy/Medicine, University of South of Santa Catarina, Tubarão 88700000, SC, Brazil. kelserkock@yahoo.com.br

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJCC* as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xu Guo*; Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

August 26, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Sodium-glucose cotransporter-2 inhibitor-associated euglycemic diabetic ketoacidosis in COVID-19-infected patients: A systematic review of case reports

Anwar Khedr, Hussam Al Hennawi, Muhammed Khuzzaim Khan, Aalaa Eissa, Mikael Mir, Ibtisam Rauf, Jain Nitesh, Salim Surani, Syed Anjum Khan

Specialty type: Medicine, general and internal

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C, C, C
Grade D (Fair): 0
Grade E (Poor): E

P-Reviewer: Feizi A, Iran;
Martinez-Castelao A, Spain; Shao JQ, China; Su G, China

Received: May 29, 2023

Peer-review started: May 29, 2023

First decision: June 19, 2023

Revised: July 13, 2023

Accepted: July 28, 2023

Article in press: July 28, 2023

Published online: August 26, 2023



Anwar Khedr, Department of Medicine, Bronx Care Health System, Bronx, NY 10457, United States

Hussam Al Hennawi, Department of Internal Medicine, Jefferson Abington Hospital, Abington, PA 19001, United States

Muhammed Khuzzaim Khan, Department of Internal Medicine, Dow University of Health Science, Karachi 74200, Pakistan

Aalaa Eissa, Department of Medicine, Kafrelsheikh University, KFS 33511, Egypt

Mikael Mir, Department of Medicine, University of Minnesota, Minneapolis, MN 55455, United States

Ibtisam Rauf, Department of Medicine, St. George's University, School of Medicine, St. George SW17 0RE, Grenada

Jain Nitesh, Department of Critical Care, Mayo Clinic Health System, Mankato, MN 56001, United States

Salim Surani, Department of Medicine & Pharmacology, Texas A&M University, College Station, TX 77843, United States

Syed Anjum Khan, Department of Critical Care Medicine, Mayo Clinic Health System, Mankato, MN 56001, United States

Corresponding author: Salim Surani, FCCP, MD, MHSc, Academic Editor, Professor, Department of Medicine & Pharmacology, Texas A&M University, 400 Bizzell St, College Station, TX 77843, United States. srsurani@hotmail.com

Abstract

BACKGROUND

Diabetic ketoacidosis (DKA) manifests as hyperglycemia, metabolic acidosis, and ketosis. However, euglycemic DKA (eu-DKA) conceals severe DKA with glucose levels below 200 mg/dL. Sodium-glucose cotransporter-2 (SGLT2) inhibitors can induce eu-DKA in diabetic patients. Notably, coronavirus disease 2019 (COVID-19)-infected individuals with diabetes using SGLT2 inhibitors face an augmented

risk of eu-DKA due to the direct toxic impact of the virus on pancreatic islets. This study aims to comprehensively investigate the association between SGLT2 inhibitors and eu-DKA in COVID-19 patients through meticulous case report analysis. Additionally, we endeavor to examine the outcomes and treatment approaches for COVID-19-infected diabetics receiving SGLT2 inhibitors, providing indispensable insights for healthcare professionals managing this specific patient population.

AIM

To investigate the connection between SGLT2 inhibitors and euglycemic DKA in COVID-19 patients through a meticulous analysis of case reports.

METHODS

We conducted an exhaustive search across prominent electronic databases, including PubMed, SCOPUS, Web of Science, and Google Scholar. This search encompassed the period from December 2019 to May 2022, incorporating published studies and pre-prints. The search terms employed encompassed “SGLT2 inhibitors”, “euglycemic DKA”, “COVID-19”, and related variations. By incorporating these diverse sources, our objective was to ensure a thorough exploration of the existing literature on this subject, thereby augmenting the validity and robustness of our findings.

RESULTS

Our search yielded a total of seven case reports and one case series, collectively comprising a cohort of twelve patients. These reports detailed instances of eu-DKA in individuals with COVID-19. Crucially, all twelve patients were utilizing SGLT2 as their primary anti-diabetic medication. Upon admission, all oral medications were promptly discontinued, and the patients were initiated on intravenous insulin therapy to effectively manage the DKA. Encouragingly, eleven patients demonstrated a favorable outcome, while regrettably, one patient succumbed to the condition. Subsequently, SGLT2 were discontinued for all patients upon their discharge from the hospital. These findings provide valuable insights into the clinical management and outcomes of eu-DKA cases associated with COVID-19 and SGLT2, underscoring the critical importance of prompt intervention and vigilant medication adjustments.

CONCLUSION

Our study sheds light on the possibility of diabetic patients developing both drug-related and unrelated DKA, as well as encountering adverse outcomes in the context of COVID-19, despite maintaining satisfactory glycemic control. The relationship between glycemic control and clinical outcomes in COVID-19 remains ambiguous. Consequently, this systematic review proposes that COVID-19-infected diabetic patients using SGLT2 should contemplate alternative treatment protocols until their recovery from the disease.

Key Words: Sodium-glucose transporter 2 inhibitors; COVID-19; SARS-CoV-2; Diabetic ketoacidosis; Euglycemic diabetic ketoacidosis; Diabetes mellitus

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: This systematic review provides a comprehensive analysis of the relationship between sodium-glucose cotransporter-2 (SGLT2) inhibitors, euglycemic diabetic ketoacidosis (eu-DKA), and coronavirus disease 2019 (COVID-19) in patients with diabetes. Despite maintaining optimal glycemic control, individuals using SGLT2 inhibitors are still susceptible to both drug-induced and unrelated diabetic ketoacidosis (DKA), with potential adverse consequences during COVID-19 infection. Clinicians should exercise caution when prescribing SGLT2 inhibitors to diabetic patients affected by COVID-19 and carefully consider alternative treatment strategies. A thorough understanding of the intricate interplay between SGLT2 inhibitors, euglycemic DKA, and COVID-19 is crucial for optimizing patient management and achieving favorable clinical outcomes.

Citation: Khedr A, Hennawi HA, Khan MK, Eissa A, Mir M, Rauf I, Nitesh J, Surani S, Khan SA. Sodium-glucose cotransporter-2 inhibitor-associated euglycemic diabetic ketoacidosis in COVID-19-infected patients: A systematic review of case reports. *World J Clin Cases* 2023; 11(24): 5700-5709

URL: <https://www.wjgnet.com/2307-8960/full/v11/i24/5700.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i24.5700>

INTRODUCTION

Understanding the signs, symptoms, risk factors, and outcomes of severe acute respiratory syndrome coronavirus 2

(SARS-CoV-2) infection remains an ongoing endeavor for clinicians and scientists. While coronavirus disease 2019 (COVID-19) research progresses rapidly, recent discoveries call for rigorous investigation to establish new hypotheses and evidence-based conclusions. Patients with diabetes mellitus face a higher likelihood of developing severe COVID-19 and experiencing adverse outcomes[1]. Notably, SGLT2 inhibitors exhibit cardiovascular benefits that extend beyond glycemic control[2]. Diabetic ketoacidosis (DKA) represents a life-threatening complication of diabetes mellitus (DM) characterized by elevated serum glucose (> 250 mg/dL), high anion gap metabolic acidosis, and plasma ketone levels. However, DKA can occur even with mild to moderate serum glucose elevation, known as euglycemic DKA (eu-DKA)[3].

The Food and Drug Administration has issued warnings about the increased risk of eu-DKA associated with sodium-glucose cotransporter-2 (SGLT2) inhibitors[4]. However, findings from the CANVAS trial suggest a low incidence of eu-DKA in patients using SGLT2 inhibitors (0.6 events per 1000 patient years)[5]. It is important to note that COVID-19 infection may amplify this risk due to the potential pancreato-toxic effects of the virus. Although three studies have examined the relationship between SGLT2 inhibitors and eu-DKA, methodological variations exist[6-8]. The prevalence of eu-DKA among type 2 diabetes mellitus (T2DM) patients treated with SGLT2 inhibitors is less than 0.1%[9]. Eu-DKA is a critical crisis that often goes unnoticed due to the absence of evident hyperglycemic symptoms. Pathophysiological studies have elucidated the mechanisms underlying this effect[10]. SGLT2 inhibitors induce glucosuria by inhibiting the sodium-glucose cotransporter in the convoluted proximal tubule, leading to decreased serum glucose levels and subsequent insulinopenia. Insulinopenia triggers an increase in counter-regulatory hormones (catecholamines and glucagon), which stimulates lipolysis and enhances fatty acid production, resulting in ketosis in the euglycemic state given the prior reduction in blood glucose by SGLT2 inhibitors.

The combined effect of COVID-19's pancreatic toxicity and SGLT2 inhibitors' glucosuria induces severe insulinopenia, potentially elevating the risk of eu-DKA in affected patients. However, the understanding of this mechanism and the management of eu-DKA in individuals with concurrent COVID-19 infection remains limited in the existing literature. In this systematic review, we have included a comprehensive analysis of relevant case report studies that provide important insights into the association between SGLT2 inhibitors, eu-DKA, and COVID-19 in patients with diabetes mellitus. These case reports shed light on the clinical manifestations, treatment approaches, and outcomes of eu-DKA in individuals with COVID-19 who are receiving SGLT2 inhibitor therapy.

MATERIALS AND METHODS

Data sources and search strategy

This systematic review adhered to the guidelines outlined in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA)[11]. A comprehensive electronic search was conducted from December 2019 to May 2022 using the SCOPUS, Web of Science, MEDLINE, and Google Scholar databases. The search was limited to English language publications, and the detailed search string employed is provided in the [Supplementary material](#). In addition, we included the pharmaceutical, generic, and trade names of SGLT2 inhibitors in our search. To ensure thoroughness, we manually examined the reference lists of the included studies to identify any potentially relevant articles that might have been overlooked. Detailed search strategies used for all databases are available in the [Supplementary material](#). The research protocol for this review was registered in the International Prospective Register of Systematic Reviews under registration number CRD42022341562.

Study selection

We included published case reports, case series, and pre-print articles. The selection of studies was based on specific eligibility criteria, which included the following: (1) Reports describing patients with euglycemic diabetic ketoacidosis (eu-DKA) who were admitted and treated following the guidelines set by the Association of British Clinical Diabetologists[12]; (2) inclusion of adult patients with either type 1 diabetes mellitus (T1DM) or T2DM; (3) utilization of SGLT2 inhibitors as the primary intervention for glucose control; and (4) confirmation of eu-DKA cases. Exclusion criteria were as follows: (1) Studies that did not focus on the association between SGLT2 inhibitors and eu-DKA in COVID-19 patients; (2) studies involving pediatric patients; (3) non-English language publications; (4) studies without clear documentation of eu-DKA cases or treatment approaches; and (5) review articles, editorials, and conference abstracts. The study selection process followed specific eligibility criteria based on the PICO framework:

Participants (P): Adult patients with either T1DM or T2DM diagnosed with eu-DKA.

Intervention (I): Utilization of SGLT2 inhibitors as the primary intervention for glucose control.

Comparator (C): Not applicable as this review focused on the association between SGLT2 inhibitors and eu-DKA in COVID-19 patients.

Outcomes (O): Clinical manifestations, treatment approaches, and outcomes of eu-DKA in patients with concurrent COVID-19 infection. The favorable outcome was defined as survival and hospital discharge.

Data extraction and assessment of study quality

The systematic search yielded articles that were imported into the Endnote Reference Library software, where duplicates were identified and removed. Initially, two independent reviewers (Khedr A and Hennawi HA) screened the articles based on titles and abstracts. Subsequently, two additional independent reviewers (Mir M and Rauf I) performed full-text screening of the filtered articles. The entire content of each article was thoroughly reviewed, with two authors (Khan MK and Eissa A) ensuring its relevance and piloting the process with one report. In case of any discrepancies, a third investigator (Nitesh J) was consulted for resolution. Information extracted from the articles included demographic background,

comorbidities, disease onset, initial symptoms, laboratory tests, diabetes mellitus type, SGLT2 inhibitor type, study date, study design, treatment intervention, and case outcomes. The quality of the included studies was assessed using the CARE (case report) guidelines[13].

Synthesis of results

The extracted information was qualitatively synthesized using a narrative approach due to the limitations of the included case reports, which presented small effect sizes and precluded quantitative analysis for calculating effect estimates.

RESULTS

Study results, study characteristics, and baseline demographics

The study selection process was summarized using the PRISMA flow chart (Figure 1), which illustrates the search and selection of studies. Initially, a total of 3370 studies were identified from the databases, and after removing duplicates ($n = 1823$), eight studies met the inclusion criteria and were included in this review. These studies consisted of seven case reports and one case series that investigated the incidence of eu-DKA in COVID-19 patients receiving SGLT2 inhibitors [14-21]. The combined study population across these studies comprised 12 patients.

Quality assessment

All the included case reports adhered to the CARE guidelines, ensuring a standardized reporting of case details. A comprehensive quality assessment table can be found in Supplementary Table 1, providing a detailed evaluation of the methodological quality of each included study.

Evidence synthesis

The patients included in the studies were from various countries, including the United States ($n = 3$)[14,18,19], United Kingdom ($n = 5$)[15], Brazil ($n = 1$)[16], Malaysia ($n = 1$)[20], and Belgium ($n = 2$)[17,21]. The reported cases of type 1 diabetes mellitus (T1DM) were observed in the studies conducted in the United Kingdom and Belgium[17,21].

The age range of the patients included in the studies varied from 40 to 79 years, with a higher proportion of male patients (75%) compared to female patients. Among the patients, T2DM was the most prevalent form, accounting for 83.33% of the cases. The most commonly observed comorbidities among the patients were hypertension (50%), hypothyroidism (16.66%), obstructive sleep apnea (16.7%), and hyperlipidemia (8.3%). Empagliflozin was the predominant SGLT2 inhibitor used by the patients, accounting for 83.3% of the cases.

Common symptoms reported by the patients included tachypnea, dyspnea, and tachycardia. The diagnostic criteria for eu-DKA frequently involved a combination of arterial blood gas (ABG) analysis, serum tests, and urine analysis. Serum glucose levels ranged from 113 to 286 mg/dL. ABG analysis revealed deviations from normal levels, with pCO₂ ranging from 13 to 43 mmHg, bicarbonate ranging from 3 to 20 mEq/L, and pH ranging from 6.94 to 7.48.

COVID-19 was primarily diagnosed using reverse transcription-polymerase chain reaction, and the symptoms of eu-DKA appeared 2 to 9 d after the identification of COVID-19. Treatment for COVID-19 included intubation and oxygen therapy for the majority of patients, except for one patient with T1DM who also required invasive mechanical ventilation. Another patient received High Flow Nasal Cannula. Empagliflozin use was associated with a shorter duration between COVID-19 infection and the onset of eu-DKA symptoms. All patients received intravenous fluids and intravenous insulin as part of the eu-DKA treatment protocol.

Out of the 12 patients included in the studies, 11 patients successfully survived, recovered, and were discharged from the hospital, while unfortunately, one patient died. A favorable outcome was defined as not requiring oxygen or life support and having blood pH within the normal range of 7.35-7.45. The patients with T1DM also experienced successful recoveries. SGLT2 inhibitors were discontinued until the resolution of COVID-19, and subcutaneous insulin was initiated after recovery from eu-DKA. A summary of the results can be found in Table 1.

DISCUSSION

The use of SGLT2 inhibitors in adults with COVID-19 infection remains a controversial topic, and this systematic review represents the first attempt to define the outcomes of this particular patient cohort. However, the availability of data is unfortunately limited. Nonetheless, the available data suggest that SGLT2 inhibitors may contribute to the development of eu-DKA in COVID-19-infected patients.

We included 8 studies comprising a total of 12 patients. The survival rate among these patients exceeded 90%. Most patients presented with symptoms of COVID-19 within 2-3 d before the onset of eu-DKA. Significant deviations in arterial blood pH levels were observed, with the lowest reported pH being 6.87. In response, all patients had their SGLT2 inhibitors immediately discontinued and received treatment with fluids and intravenous insulin.

The term "eu-DKA" was first defined by Munro *et al*[22] in 1973. It is distinct from classic DKA in that it is characterized by severe metabolic acidosis despite normal blood glucose levels. Diagnosis of eu-DKA is confirmed through direct measurement of beta-hydroxybutyrate levels in the blood and assessment of arterial blood pH levels[23].

Table 1 Data summarized from included studies

Ref.	Study Origin	Number of patients	Sex	Age	Type of DM	Previous co-morbidities	SGLT2 inhibitors used before	COVID-19 symptoms onset	Eu-DKA diagnosis	Method of COVID-19 diagnosis	Autonomic symptoms	ABG count	Serum Analysis	Urinalysis	Eu-DKA management	COVID-19 Management	Outcome	Treatment intervention after resolution
Dass <i>et al</i> [14]	United States	1	Female	59	T2DM	COVID-19, Community acquired pneumonia	Empagliflozin, sitagliptin	9 d before eu-DKA onset	ABG analysis, serum analysis, urine analysis	Not reported	Tachypnea and tachycardia	pH: 6.94; PaCO ₂ : 13; PaO ₂ : 99; HCO ₃ : 3	Lactate: 0.9; glucose: 154; confirmed bicarb: < 10; serum osmolality: 346; anion gap: 30	3+ glucose and 2+ ketones; negative UDS and normal salicylate levels	insulin drip and IV fluid	Not reported	Discharged after complete recovery	Sitagliptin and metformin continued; empagliflozin discontinued; started with 20 units of insulin glargine
Vitale <i>et al</i> [15]	United Kingdom	5	3 males, 2 females	52-79	T2DM	Covid-19, Hypertension	Empagliflozin, Canagliflozin	3-8 d before eu-DKA onset	ABG analysis, serum analysis	RT-PCR	Tachypnea, dyspnea, nausea, anorexia, abdominal pain	pH: 7.09-7.31; PaCO ₂ : 19-43; HCO ₃ : 5-20	Lactate: 1.1-2.4; glucose: 146-286; anion gap: 20-40	Not analyzed	IV insulin, intubation	Not reported	1 male died; rest of the patients recovered	Not reported
Batista <i>et al</i> [16]	Brazil	1	Male	56	T2DM	COVID-19	Empagliflozin	5 d before eu-DKA onset	ABG analysis, serum analysis, urine analysis	RT-PCR	Tachypnea, tachycardia	pH: 7.28; pCO ₂ : 19 mmHg; HCO ₃ : 8.9; base excess: 15.7	Sodium: 132; potassium: 5.6; chloride: 99; glucose: 118; hemoglobin A1c (HbA1c) 7.2%	ketones	Glucose, insulin, and KCl	Oxygen therapy, Azithromycin 500 mg	Discharged after complete recovery	Not reported
Philippe <i>et al</i> [17]	Belgium	1	Male	60	T1DM	COVID-19, hypothyroidism, obstructive sleep apnea	Empagliflozin	2 d before eu-DKA onset	ABG analysis, serum analysis, urine analysis	RT-PCR	Polypnea, myalgia, diarrhea. And fever	pH: 7.48; pO ₂ : 17; pCO ₂ : 27; HCO ₃ : 19; anion gap: 17	Lactate: 1.4; glucose: 234; HbA1c: 7.4	3+ ketones	IV fluids, IV insulin	Oxygen therapy, invasive mechanical ventilation	Discharged after complete recovery	Empagliflozin was discontinued and subcutaneous glargine was started
Morrison <i>et al</i> [18]	United States	1	Male	40	T2DM	COVID-19	Empagliflozin	3 d before eu-DKA onset	ABG analysis, serum	RT-PCR	Tachypnea, tachycardia, diaphoretic	pH: 7.06; pCO ₂ :	Sodium level: 133; carbon	Glucose (> 1000 mg/dL)	IV fluids, IV insulin	No intervention due to mild	Discharged after complete	Subcutaneous glargine started

									analysis, urine analysis			37; pO2: 31; HCO3: 10.0; lactate: 2.3	dioxide: 11; glucose:177; anion gap: 25; HbA1c: 10.6%	and ketones (> 80 mg/dL)		symptoms	recovery	
Fang <i>et al</i> [19]	United States	1	Male	52	T2DM	COVID-19, hypertension, hyperlipidemia	Empagliflozin	2 d before eu-DKA onset	ABG analysis, serum analysis, urine analysis	RT-PCR	Hypoxia, dyspnea. Fever, anorexia	pH: 7.30; pCO2: 37	glucose:113; anion gap: 18	Glucose (> 500 mg/ dL) and ketones (> 80 mg/dL)	IV fluids, IV insulin	Intubation, oxygen therapy	Discharged after complete recovery	Not reported
Yii ESS <i>et al</i> [28]	Malaysia	1	Male	37	T2DM	COVID-19	Empagliflozin	3 d before eu-DKA onset	ABG analysis, serum analysis, urine analysis	Not reported	Dyspnea, Tachypnea, Stable heart rate	pH: 6.87; pCO2: 17; pO2: 37; HCO3: 3.1; lactate: 1.7	Glucose: 11.9; urea 11.4; sodium 136; potassium 4.5; chloride 106; creatinine 105	Ketone: 3.0	18 h of renal replacement therapy, noradrenaline infusion	Intubation, ICU	Discharged after complete recovery	Not reported
Oriot <i>et al</i> [29]	Belgium	1	Male	52	T1DM	COVID-19, hypothyroidism, obstructive sleep apnea syndrome	Empagliflozin	2 d before	ABG analysis, serum analysis	RT-PCR	Polypnea, myalgia, diarrhea, and fever	pH: 7.48; pCO2: 27; pO2: 47; HCO3: 19; lactate: 1.4; anion gap: 17	HbA1c: 7.4	Ketone: 3+	IC fluids, IV insulin	Not reported	Discharged after complete recovery	Empagliflozin was discontinued

Reporting units: Glucose (mg/dL), ketones (mg/dL), PaCO₂ and PaO₂ (mmHg), HCO₃ (mEq/L), Lactate (mmol/L), Chloride measuring unit (mEq/L), Osmolarity (mOsm/Kg), CO₂ (mEq/L), Ketone: mmol/L.

ABG: Arterial blood gases; COVID-19: Coronavirus disease 2019; ICU: Intensive care unit; RT-PCR: Real-time polymerase chain reaction; HB: Hemoglobin; DM: Diabetes mellitus; Eu-DKA: Diabetic ketoacidosis with glucose.

The occurrence of eu-DKA due to SGLT2 inhibitors involves decreased insulin production (insulinopenia) and increased glucagon secretion. Multiple factors influence increased glucagon secretion, which occurs through both direct and indirect mechanisms. The inhibitory effects of SGLT2 inhibitors on the SGLT2 transporters in the glucagon-secreting pancreatic alpha cells of the Langerhans islets directly contribute to increased glucagon secretion. Indirectly, increased glucose excretion leads to lower insulin levels and decreased insulin to glucagon ratio. The resultant decrease in insulin stimulates the synthesis of free fatty acids and ketone bodies, leading to excessive catabolism of fatty acids and subsequent ketosis[24]. Ketosis is further exacerbated in the presence of SGLT2 inhibitors, as these medications inhibit the




Figure 1 PRISMA chart.

reabsorption of glucose in the proximal renal tubules, leading to glucosuria and promoting a state of starvation[25].

Eu-DKA has emerged as a prevalent condition observed in diabetic patients with COVID-19 who are also using SGLT2 inhibitors. The higher incidence of eu-DKA in male diabetic patients with COVID-19 aligns with reports indicating a greater propensity for presentation in this subgroup. The clinical manifestation of eu-DKA in COVID-19-infected diabetic patients taking SGLT2 inhibitors is diverse, including tachypnea and tachycardia during clinical examination, along with an anion gap metabolic acidosis and normal serum glucose levels observed in laboratory investigations.

In patients infected with COVID-19, eu-DKA differs from the non-infected population due to the pancreatic toxicity induced by the virus, leading to severe insulinopenia. Consequently, infected patients require insulin therapy in addition to the discontinuation of SGLT2 inhibitors to prevent the recurrence of eu-DKA. Conversely, in patients without COVID-19 infection, resolution and prevention of eu-DKA can typically be achieved by discontinuing SGLT2 inhibitors, allowing insulin levels to return to normal[18]. Unlike individual case reports, the accumulation of evidence from multiple cases supports the occurrence of eu-DKA in diabetic patients following the onset of COVID-19 and concurrent use of SGLT2 inhibitors.

The severe toxic effects of COVID-19 on the pancreas have been well-documented. When these toxic effects combine with the dehydrating and glycosuric effects of SGLT2 inhibitors, it can lead to the development of eu-DKA. Recent studies have discussed the occurrence of pancreatic injury in patients with COVID-19, highlighting the link between COVID-19-related pancreatic toxicity, enzyme elevation, and insulinopenia[26,27]. The angiotensin converting enzyme 2 (ACE2) receptor, which is expressed in the beta cells of Langerhans islets, plays a role in both COVID-19 and SARS-CoV-1 infections. These viruses utilize the ACE2 receptor to enter the cell, triggering immune system activation and the release of cytokines and chemokines that lead to cell death[28,29]. The destructive effect on beta cells results in insulinopenia and subsequent ketoacidosis. Moreover, the acidic environment of eu-DKA is known to facilitate the growth of COVID-19.

Given the widespread use of SGLT2 inhibitors for their cardiovascular and renal benefits, clinicians must familiarize themselves with eu-DKA to enable timely diagnosis and treatment, particularly in the context of the ongoing COVID-19 pandemic.

Limitations

This study has limitations, including a small number of studies with small effect sizes, all of which were case reports or case series. The non-randomized allocation of interventions and the absence of a standardized protocol for diagnosing eu-

DKA in COVID-19 patients introduce selection and bias risks. The diverse treatment settings and limited data availability further hinder comprehensive analysis. Further research is needed to address these limitations and provide more robust evidence on the topic.

CONCLUSION

Our systematic review suggests that SGLT2 inhibitors may increase the risk of euglycemic diabetic ketoacidosis in COVID-19-infected diabetic patients. The pancreatic toxicity associated with SARS-CoV-2 infection may contribute to this effect. While this narrative synthesis of case reports provides valuable insights, further research with larger sample sizes and rigorous designs, such as retrospective cohorts, is needed to investigate the association between eu-DKA development in COVID-19 patients and SGLT2 inhibitors. Studies with larger effect sizes and randomization would help elucidate clinically relevant endpoints and enable more effective management of individuals at higher risk.

ARTICLE HIGHLIGHTS

Research background

The coexistence of coronavirus disease 2019 (COVID-19) infection and the use of sodium-glucose cotransporter-2 (SGLT2) inhibitors have generated debate due to the potential risk of euglycemic diabetic ketoacidosis (eu-DKA) development. Limited information regarding this specific patient population is available, necessitating a systematic review to investigate the outcomes and characteristics associated with eu-DKA in COVID-19-infected diabetic patients treated with SGLT2 inhibitors.

Research motivation

Given the controversy and limited data surrounding the association between SGLT2 inhibitors, COVID-19 infection, and eu-DKA, there is a pressing need to investigate this topic to enhance our understanding of the potential risks and outcomes.

Research objectives

The primary objectives of this study are to examine the association between SGLT2 inhibitors and the development of eu-DKA in COVID-19-infected diabetic patients, explore the potential mechanisms underlying this relationship, and assess the clinical outcomes and management strategies for this patient population.

Research methods

We conducted a comprehensive search of relevant databases to identify studies reporting on the association between SGLT2 inhibitors and eu-DKA in COVID-19-infected diabetic patients. We followed the PRISMA guidelines for study selection and data extraction. The extracted data were qualitatively synthesized to provide a narrative overview of the findings.

Research results

The systematic review included eight studies comprising 12 patients, investigating the association between SGLT2 inhibitors and eu-DKA in COVID-19-infected diabetic patients. The majority of patients presented with eu-DKA symptoms 2-3 d after the onset of COVID-19 symptoms. The survival rate was over 90%, with one reported fatality. Significant pH deviations were observed, with the lowest reported pH being 6.87. All patients discontinued SGLT2 inhibitors and received treatment with fluids and IV insulin. The results highlight the potential risk of developing eu-DKA in this patient population.

Research conclusions

This systematic review concludes that the use of SGLT2 inhibitors in COVID-19-infected diabetic patients may increase the risk of eu-DKA. The pancreatic toxicity induced by the severe acute respiratory syndrome coronavirus 2 virus is believed to contribute to this phenomenon. The analysis of case reports provides evidence supporting the association between SGLT2 inhibitors and eu-DKA in this patient population. Further studies with larger sample sizes and robust designs are necessary to enhance our understanding and inform clinical decision-making for high-risk individuals.

Research perspectives

Further research is needed to investigate the mechanisms of eu-DKA in COVID-19 patients on SGLT2 inhibitors. Larger randomized studies are necessary to establish a causal relationship and identify risk factors. Standardized protocols for diagnosis and management should be developed to improve patient outcomes. These research perspectives will enhance understanding and guide evidence-based approaches in the future.

FOOTNOTES

Author contributions: Khedr A, Hennawi HA, Khan MK, Eissa A, Mir M, and Rauf I contributed to the conception and design of the manuscript; Khan MK, Eissa A, Mir M, Rauf I and Nitesh J contributed to the collection of the data; Khedr A, Hennawi HA, Mir M, Rauf I, and Nitesh J contributed to the interpretation of the analysis; Khedr A, Hennawi HA, Khan MK, Eissa A, and Nitesh J drafted the manuscript; Surani S and Khan SA provided the oversight; All the authors critically revised the manuscript, gave final approval, and agreed to be accountable for all aspects of the work, ensuring integrity and accuracy.

Conflict-of-interest statement: None of the authors have any conflict to disclose.

PRISMA 2009 Checklist statement: The authors have read the PRISMA 2009 Checklist, and the manuscript was prepared and revised according to the PRISMA 2009 Checklist.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: United States

ORCID number: Anwar Khedr 0000-0002-2730-3031; Hussam Al Hennawi 0000-0002-1968-4962; Muhammed Khuzzaim Khan 0000-0003-4586-8607; Aalaa Eissa 0000-0002-3814-6510; Mikael Mir 0000-0002-4984-4621; Ibtisam Rauf 0000-0002-0160-5592; Jain Nitesh 0000-0003-0139-3515; Salim Surani 0000-0001-7105-4266; Syed Anjum Khan 0000-0002-2452-2079.

Corresponding Author's Membership in Professional Societies: American College of Chest Physician; American College of Physician; Society of Critical Care Medicine.

S-Editor: Liu JH

L-Editor: A

P-Editor: Yuan YY

REFERENCES

- Peña JE, Rascón-Pacheco RA, Ascencio-Montiel IJ, González-Figueroa E, Fernández-Gárate JE, Medina-Gómez OS, Borja-Bustamante P, Santillán-Oropeza JA, Borja-Aburto VH. Hypertension, Diabetes and Obesity, Major Risk Factors for Death in Patients with COVID-19 in Mexico. *Arch Med Res* 2021; **52**: 443-449 [PMID: 33380361 DOI: 10.1016/j.arcmed.2020.12.002]
- Usman MS, Siddiqi TJ, Memon MM, Khan MS, Rawasia WF, Talha Ayub M, Sreenivasan J, Golzar Y. Sodium-glucose co-transporter 2 inhibitors and cardiovascular outcomes: A systematic review and meta-analysis. *Eur J Prev Cardiol* 2018; **25**: 495-502 [PMID: 29372664 DOI: 10.1177/2047487318755531]
- Modi A, Agrawal A, Morgan F. Euglycemic Diabetic Ketoacidosis: A Review. *Curr Diabetes Rev* 2017; **13**: 315-321 [PMID: 27097605 DOI: 10.2174/1573399812666160421121307]
- Food and Drug Administration. The FDA warns that SGLT2 inhibitors for diabetes can cause a serious condition of too much acid in the blood. FDA gov, 2015 [DOI: 10.2165/00128415-201013130-00009]
- Neal B, Perkovic V, Mahaffey KW, de Zeeuw D, Fulcher G, Erond N, Shaw W, Law G, Desai M, Matthews DR; CANVAS Program Collaborative Group. Canagliflozin and Cardiovascular and Renal Events in Type 2 Diabetes. *N Engl J Med* 2017; **377**: 644-657 [PMID: 28605608 DOI: 10.1056/NEJMoa1611925]
- Barski L, Eshkoli T, Brandstaetter E, Jotkowitz A. Euglycemic diabetic ketoacidosis. *Eur J Intern Med* 2019; **63**: 9-14 [PMID: 30910328 DOI: 10.1016/j.ejim.2019.03.014]
- Peters AL, Buschur EO, Buse JB, Cohan P, Diner JC, Hirsch IB. Euglycemic Diabetic Ketoacidosis: A Potential Complication of Treatment With Sodium-Glucose Cotransporter 2 Inhibition. *Diabetes Care* 2015; **38**: 1687-1693 [PMID: 26078479 DOI: 10.2337/dc15-0843]
- Meyer EJ, Gabb G, Jesudason D. SGLT2 Inhibitor-Associated Euglycemic Diabetic Ketoacidosis: A South Australian Clinical Case Series and Australian Spontaneous Adverse Event Notifications. *Diabetes Care* 2018; **41**: e47-e49 [PMID: 29440112 DOI: 10.2337/dc17-1721]
- Tang H, Li D, Wang T, Zhai S, Song Y. Effect of Sodium-Glucose Cotransporter 2 Inhibitors on Diabetic Ketoacidosis Among Patients With Type 2 Diabetes: A Meta-analysis of Randomized Controlled Trials. *Diabetes Care* 2016; **39**: e123-e124 [PMID: 27311492 DOI: 10.2337/dc16-0885]
- Ogawa W, Sakaguchi K. Euglycemic diabetic ketoacidosis induced by SGLT2 inhibitors: possible mechanism and contributing factors. *J Diabetes Investig* 2016; **7**: 135-138 [PMID: 27042263 DOI: 10.1111/jdi.12401]
- Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009; **6**: e1000097 [PMID: 19621072 DOI: 10.1371/journal.pmed.1000097]
- Winocour PH, Ford M, Ainsworth A; Association of British Clinical Diabetologists. Association of British Clinical Diabetologists (ABCD): survey of specialist diabetes care services in the UK, 2000. 2. Workforce issues, roles and responsibilities of diabetes specialist nurses. *Diabet Med* 2002; **19** Suppl 4: 27-31 [PMID: 12121334 DOI: 10.1046/j.1464-5491.19.s4.6.x]
- Gagnier JJ, Kienle G, Altman DG, Moher D, Sox H, Riley D; CARE Group. The CARE guidelines: consensus-based clinical case report guideline development. *J Diet Suppl* 2013; **10**: 381-390 [PMID: 24237192 DOI: 10.3109/19390211.2013.830679]
- Dass B, Beck A, Holmes C, Morton G. Euglycemic DKA (euDKA) as a presentation of COVID-19. *Clin Case Rep* 2021; **9**: 395-398 [PMID: 33362928 DOI: 10.1002/ccr3.3540]
- Vitale RJ, Valtis YK, McDonnell ME, Palermo NE, Fisher NDL. Euglycemic Diabetic Ketoacidosis With COVID-19 Infection in Patients

- With Type 2 Diabetes Taking SGLT2 Inhibitors. *AACE Clin Case Rep* 2021; **7**: 10-13 [PMID: 33521255 DOI: 10.1016/j.aace.2020.11.019]
- 16 **Batista DV**, Vieira CAFA, Costa TA, Lima EG. COVID-19-associated euglycemic diabetic ketoacidosis in a patient with type 2 diabetes on SGLT2 inhibitor: a case report. *Diabetol Int* 2021; **12**: 313-316 [PMID: 33133998 DOI: 10.1007/s13340-020-00473-3]
 - 17 **Oriot P**, Hermans MP. Euglycemic diabetic ketoacidosis in a patient with type 1 diabetes and SARS-CoV-2 pneumonia: case-report and review of the literature. *Acta Clin Belg* 2022; **77**: 113-117 [PMID: 32544373 DOI: 10.1080/17843286.2020.1780390]
 - 18 **Morrison N**, Barnett K, Tantum J, Morrison HK, Whalen M. A Case of Euglycemic Diabetic Ketoacidosis in a Patient With Type 2 Diabetes Mellitus and COVID-19. *Cureus* 2020; **12**: e12029 [DOI: 10.7759/cureus.12029]
 - 19 **Fang J**, Genco M, Caskey RN. COVID-19 Precipitating Euglycaemic Diabetic Ketoacidosis with SGLT2 Inhibitor Use. *Eur J Case Rep Intern Med* 2020; **7**: 001943 [PMID: 33194873 DOI: 10.12890/2020_001943]
 - 20 **Qiu H**, Novikov A, Vallon V. Ketosis and diabetic ketoacidosis in response to SGLT2 inhibitors: Basic mechanisms and therapeutic perspectives. *Diabetes Metab Res Rev* 2017; **33** [PMID: 28099783 DOI: 10.1002/dmrr.2886]
 - 21 **Yii ESS**, Azli AW, Sitaram PN. Sodium-glucose cotransporter 2 inhibitor-induced euglycemic diabetic ketoacidosis in a patient with coronavirus disease 2019: a case report. *J Med Case Rep* 2022; **16**: 17 [PMID: 34983625 DOI: 10.1186/s13256-021-03232-3]
 - 22 **Munro JF**, Campbell IW, McCuish AC, Duncan LJ. Euglycaemic diabetic ketoacidosis. *Br Med J* 1973; **2**: 578-580 [PMID: 4197425 DOI: 10.1136/bmj.2.5866.578]
 - 23 **Vanelli M**, Chiari G, Capuano C, Iovane B, Bernardini A, Giacalone T. The direct measurement of 3-beta-hydroxy butyrate enhances the management of diabetic ketoacidosis in children and reduces time and costs of treatment. *Diabetes Nutr Metab* 2003; **16**: 312-316 [PMID: 15000443]
 - 24 **Taylor SI**, Blau JE, Rother KI. SGLT2 Inhibitors May Predispose to Ketoacidosis. *J Clin Endocrinol Metab* 2015; **100**: 2849-2852 [PMID: 26086329 DOI: 10.1210/jc.2015-1884]
 - 25 **Yang L**, Han Y, Nilsson-Payant BE, Gupta V, Wang P, Duan X, Tang X, Zhu J, Zhao Z, Jaffré F, Zhang T, Kim TW, Harschnitz O, Redmond D, Houghton S, Liu C, Naji A, Cicci G, Guttikonda S, Bram Y, Nguyen DT, Cioffi M, Chandar V, Hoagland DA, Huang Y, Xiang J, Wang H, Lyden D, Borczuk A, Chen HJ, Studer L, Pan FC, Ho DD, tenOever BR, Evans T, Schwartz RE, Chen S. A Human Pluripotent Stem Cell-based Platform to Study SARS-CoV-2 Tropism and Model Virus Infection in Human Cells and Organoids. *Cell Stem Cell* 2020; **27**: 125-136.e7 [PMID: 32579880 DOI: 10.1016/j.stem.2020.06.015]
 - 26 **Wang F**, Wang H, Fan J, Zhang Y, Zhao Q. Pancreatic Injury Patterns in Patients With Coronavirus Disease 19 Pneumonia. *Gastroenterology* 2020; **159**: 367-370 [PMID: 32247022 DOI: 10.1053/j.gastro.2020.03.055]
 - 27 **Ashok A**, Faghhi M, Singh VK. Mild Pancreatic Enzyme Elevations in COVID-19 Pneumonia: Synonymous With Injury or Noise? *Gastroenterology* 2021; **160**: 1872 [PMID: 32544396 DOI: 10.1053/j.gastro.2020.05.086]
 - 28 **Yang JK**, Lin SS, Ji XJ, Guo LM. Binding of SARS coronavirus to its receptor damages islets and causes acute diabetes. *Acta Diabetol* 2010; **47**: 193-199 [PMID: 19333547 DOI: 10.1007/s00592-009-0109-4]
 - 29 **Fukuda M**, Nabeta M, Muta T, Fukami K, Takasu O. Euglycemic diabetic ketoacidosis caused by canagliflozin: a case report. *Int J Emerg Med* 2020; **13**: 2 [PMID: 31969112 DOI: 10.1186/s12245-020-0261-8]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

