

Gastrointestinal Motility Problems in Patients with ...

<https://link.springer.com/article/10.2165/00002512-199710040-00002> ▾

Oct 17, 2012 - Gastrointestinal (GI) motility disorders are frequent in patients with Parkinson's disease, manifesting mainly as **dysphagia, disorders of gastric emptying and constipation**. The most likely causes of these disorders are cerebral degeneration and degeneration of the my enteric plexus.

Cited by: 230 Author: Wolfgang H. Jost
Publish Year: 1997

GI Complications of Parkinson's Disease - UCLA Digestive ...

<https://www.uclahealth.org/gastro/motility/parkinsons> ▾

While Obtaining Clues About the Disease's Pathogenesis. Schedule a Parkinson's GI motility clinic appointment. This specialized clinic sees patients with **gastrointestinal** complications of Parkinson's disease. They will also discuss motility testing and refer you for the appropriate **study** if necessary. Call the clinic at 310-208-5400.

PEOPLE ALSO ASK

- Can Parkinson's disease cause constipation? ▾
- Is constipation a side effect of PD? ▾
- What are the symptoms of Parkinson's disease? ▾
- How does Parkinson's affect the digestive system? ▾

Feedback

Gastric emptying time and gastric motility in patients ...

<https://pubmed.ncbi.nlm.nih.gov/11748735>

Gastrointestinal symptoms such as nausea, abdominal pain, and bloating are frequent complaints of patients with Parkinson's disease (PD). It has been postulated that impaired gastrointestinal function may contribute to the development of motor fluctuations such as delay on and no on in patients with PD.

Advanced Parkinson's Disease: Bowel and Bladder Issues | ...

<https://www.andanarkinson.org/article/bowel-and> ▾

Constipation in Parkinson's Disease - ScienceDirect

<https://www.sciencedirect.com/science/article/pii/S0074774217300971>

Jan 01, 2017 · Astarloa, Mena, Sanchez, de la Vega, and de Yébenes (1992) observed that a significant improvement of constipation in PD patients determined an improvement in extrapyramidal function, which was probably related to an increase in the levodopa absorption and bioavailability due to the augmentation in gastric emptying and in gastrointestinal motility. This could explain the beneficial ...

Cited by: 21 Author: Fabrizio Stocchi, Margherita Torti Publish Year: 2017

ENT-01 Restored Bowel Motility in Constipated Parkinson's ...

<https://parkinsonsnewstoday.com/2019/09/04/enterin...>

Sep 04, 2019 · "We demonstrate in this study that ENT-01 can restore gastrointestinal motility in patients with PD, suggesting that a major division of the nervous system and the largest sensory organ in the body (the GI tract) is not irreversibly damaged in patients with PD, despite the long-standing constipation that might suggest otherwise," Denise Barbut, MD, said in a press release. Barbut is the study's senior ...

User rating: 5/5

PEOPLE ALSO ASK

Parkinson's Disease

Medical Condition

A chronic and progressive movement disorder.

- Common (More than 200,000 cases per year in US)
- Often requires lab test or imaging
- Treatments can help manage condition, no known cure
- Can last several years or be lifelong

Parkinson's disease is caused by a loss of nerve cells in substantia nigra of the brain. This leads to a reduction in a chemical called dopamine in the brain. Many different symptoms are associated with Parkinson's disease and the more common symptoms are slowness in movement and muscle stiffness. Currently there is no cure for Parkinson's disease. But treatments are available to help relieve the symptoms and maintain quality of life. These treatments include: - Supportive therapies - such as physiotherapy - Medication - Surgery

Symptoms

激活 Windows



Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 62653

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Study on the characteristics of intestinal motility of constipation in patients with Parkinson's disease

Intestinal motility in Parkinson's disease

Mei Zhang, Shuang Yang, Xiao-Cui Li, Hong-Ming Zhu, Ding Peng, Bang-Yi Li, Tian-Xu Jia, Chen Tian

Match Overview

1	Crossref 40 words Wenshu Jiao, Song Wang, Jiqi Guan, Jialu Shi et al. "Mil... ermented with <i>Lactococcus lactis</i> KLD54 0325 alleviates tol	1%
2	Crossref 24 words Anibal Sanchez Moura, Cristiano Cosme Nascimento Franc o de Sá, Celia Lopez de Cerna, Luciana Leão Vicente et al.	1%
3	Internet 47 words crawled on 21-Feb-2020 journals.nw.com	<1%
4	Internet 14 words crawled on 06-Apr-2020 academic.hep.com.cn	<1%
5	Internet 14 words crawled on 26-Jul-2020 tesesara.spandidos-publications.com	<1%

激活 Windows
转到设置以激活 Windows。



A Study of Gastrointestinal Impairment in Parkinson's Disease

<https://www.michaeljfox.org/grant/study...> ▾

A Study of Gastrointestinal Impairment in Parkinson's Disease. Gastrointestinal (GI) symptoms, such as constipation, nausea and vomiting, are common in Parkinson's disease (PD), but the cause of these symptoms is not entirely clear. Slow movement of food through the stomach, intestine and colon is one possibility.

Constipation in chronic kidney disease: it is time to ...

<https://rrtjournal.biomedcentral.com/articles/10.1186/s41100-019-0246-3> ▾

Dec 12, 2019 · Constipation is highly prevalent in patients with chronic kidney disease (CKD) and is primarily characterized by decreased intestinal motility. This chronic disorder affects the quality of life of patients. However, nephrologist and dialysis clinicians have long had a disproportionately limited understanding of constipation. Accumulating evidence has revealed a relationship between ...

Cited by: 1

Author: Ryota Ikee, Kazuhiro Yano, Tomomi Tsuru

Publish Year: 2019

PEOPLE ALSO ASK

Does "Parkinson s" cause constipation? ▾

What is the relationship between constipation and Parkinson? ▾

What are the gastrointestinal effects of Parkinsons Disease? ▾

Is it common for dialysis patients to have constipation? ▾

Feedback

ENT-01 Restored Bowel Motility in Constipated Parkinson's ...

<https://parkinsonsnewstoday.com/2019/09/04/enterin...> ▾

Sep 04, 2019 · "We demonstrate in this study that ENT-01 can restore gastrointestinal motility in patients with PD, suggesting that a major division of the nervous system and the largest sensory organ in the body (the GI tract) is not irreversibly damaged in patients with PD, despite the long-standing constipation that might suggest otherwise," Denise Barbut, MD, said in a press release. Barbut is the ...

User rating: 5/5