

## Pharmacological management of neuropathic pain in patients with vestibular schwannomas: Experience of the Atlantic Lateral Skull Base Clinic

Andrea LO Hebb, Jana Sawynok, Manohar Bance, Simon Walling, Ken Chisholm, David P Morris

Andrea LO Hebb, Atlantic Lateral Skull Base Clinic, Division of Neurosurgery, Dalhousie University, QEII Halifax Infirmary Site, Halifax NS B3H 3A7, Canada

Andrea LO Hebb, Capital District Health Authority, QEII Halifax Infirmary Site, 3<sup>rd</sup> Floor, Division of Neurosurgery, Halifax NS B3H 3A7, Canada

Jana Sawynok, Department of Pharmacology, Dalhousie University, Nova Scotia B3H 4R2, Canada

Manohar Bance, David P Morris, Division Otolaryngology, Dalhousie University, QEII Health Sciences Centre, Halifax NS B3H 2Y9, Canada

Simon Walling, Division of Neurosurgery, Dalhousie University, Halifax Infirmary Site, QEII Health Sciences Centre, Halifax NS B3H 3A7, Canada

Ken Chisholm, Department of Anesthesia and Pain Management, Dalhousie University, Halifax NS B3H 2Y9, Canada

**Author contributions:** All authors had contributed substantially to conception and design, or acquisition of data, or analysis and interpretation of data; drafted the article or revised it critically for important intellectual content and gave final approval of the version to be published.

**Correspondence to:** Dr. Andrea LO Hebb, MSc, PhD, RN, Capital District Health Authority, QEII Halifax Infirmary Site, 3<sup>rd</sup> Floor, Division of Neurosurgery, 1796 Summer Street, Halifax NS B3H 3A7, Canada. [andrea.hebb@cdha.nshealth.ca](mailto:andrea.hebb@cdha.nshealth.ca)

Telephone: +1-902-473-4824 Fax: +1-902-425-4176

Received: April 29, 2014 Revised: May 31, 2014

Accepted: June 5, 2014

Published online: June 9, 2014

### Abstract

Neuropathic pain is chronic pain generated by disorders of the peripheral and central nervous system, including skull base tumours. A skull base tumour can be any type of tumour that forms in the skull base, and this includes vestibular schwannomas which arise from the sheath of the inner ear vestibulocochlear nerve (eighth cranial nerve). Growth of the tumour, surgical resection, and/or stereotactic radiotherapy may result in

compression and/or irritation of the fifth cranial nerve (trigeminal nerve) resulting in facial pain and/or numbness. Non-trigeminal afferent input may contribute to the wide constellation of symptoms seen in orofacial pain patients. The purpose of this report was to develop a decision tool to guide the recognition and treatment of neuropathic pain in this specialized population. Recommendations for treatment are based on evidence presented in Canadian and international neuropathic treatment guidelines. Algorithms are included for assessment and treatment of adult patients with agents that are recognized to have analgesic efficacy within the broad context of neuropathic pain.

© 2014 Baishideng Publishing Group Inc. All rights reserved.

**Key words:** Acoustic neuroma; Stereotactic radiotherapy; Tricyclic antidepressants; Serotonin-norepinephrine reuptake inhibitors; Calcium channel modulators; Tramadol; Opioids

**Core tip:** The complexity of managing trigeminal neuralgia and neuropathic pain conditions among patients with skull base tumors requires a simple albeit comprehensive treatment algorithm that can be employed effectively by general practitioners, surgeons and other primary care prescribers in acute care or ambulatory clinical settings. We describe a simple treatment algorithm formulated on recommended best practice and based on clinical experience. It is intended to guide treatment, facilitate management and evaluation of outcome data (self-reported pain, quality of life measures) to elucidate the use of standardized approaches to pain management in patients with skull base etiology.

Hebb ALO, Sawynok J, Bance M, Walling S, Chisholm K, Morris DP. Pharmacological management of neuropathic pain in patients with vestibular schwannomas: Experience of the Atlantic

Lateral Skull Base Clinic. *World J Pharmacol* 2014; 3(2): 24-32  
Available from: URL: <http://www.wjgnet.com/2220-3192/full/v3/i2/24.htm> DOI: <http://dx.doi.org/10.5497/wjp.v3.i2.24>

## INTRODUCTION

This report considers pharmacological approaches to be used within the skull base tumour health community. Its intent is to offer assistance in treating patients who present with neuropathic pain (NP) in the context of orofacial, head and neck pain, including trigeminal neuralgia (TN), which is the most common craniofacial pain syndrome that is neuropathic in origin. The impetus for assembling the information stems from the experiences of practitioners within an interdisciplinary clinic who consistently rely on formularies and consultation with colleagues for advice in treating neuralgia and NP. The document offers suggested algorithms for assessment and treatment of patients with agents that are recognized to have analgesic efficacy within the broader context of NP. It makes the distinction between generalized NP<sup>[1]</sup> and NP due to skull base tumours affecting the head, neck, face and glossopharyngeal region<sup>[2,3]</sup>, but also recognizes commonalities in mechanisms underlying various forms of NP<sup>[1]</sup>.

### The Atlantic Lateral Skull Base Clinic

Vestibular schwannomas (VS) are slow growing, benign neoplasms that may be life threatening due to compression of central structures. Extra-axial tumours arising from the Schwann cell sheath of the vestibular or cochlear nerve (8<sup>th</sup> cranial nerve) are referred to as acoustic neuromas or, more appropriately, VS. Clinical diagnosis of a VS suggests an incidence of 0.7-1 per 100000 people, although incidental (*i.e.*, non-clinically relevant) discovery of VS suggests an incidence as high as 2 per 10000 people<sup>[4]</sup>. In some cases, VS are associated with NP and TN, often secondary to tumour compression or stereotactic radiation treatment affecting cranial nerves within the cerebellopontine angle (CPA).

The Atlantic Lateral Skull Base Clinic provides coordinated care through Neurotology (Division of Otolaryngology), Neurosurgery and the Stereotactic Radiotherapy Group to patients with unilateral or bilateral VS, other CPA tumours, as well as lesions of the petrous apex and jugular foramen. This program is unique in Canada, offering a single centre, multidisciplinary approach for lateral skull base lesions. The Atlantic Lateral Skull Base Clinic serves a population of more than 2 million people in a catchment area that includes Newfoundland, Prince Edward Island, New Brunswick and Nova Scotia<sup>[5]</sup>. Treatment options for skull base tumours include monitoring clinical progression, surgery, stereotactic radiation therapy (SRT), balance and hearing rehabilitation. An interdisciplinary clinic provides an ideal environment in which to identify and intervene in the treatment and management of NP affecting the head and neck region. The focus

of this work is on clinical experiences within our clinic. With low incidence, it will be very unlikely there will ever be randomized clinical trials that provide direct guidance for pain management in patients with VS. In that event, one must rely on extrapolation, and what we report is implementation of that approach and what it looks like in clinic.

### Disease background

NP is a chronic pain state that is initiated by peripheral and central nervous system injury caused by trauma, inflammation, infection or metabolic disease, and includes conditions such as distal polyneuropathy due to diabetes [diabetic neuropathy (DN)] and post-herpetic neuralgia (PHN)<sup>[1]</sup>. The Canadian Pain Society (CPS) estimates (based on 8.2% chronic NP prevalence in the general population) that 1 million Canadians live with NP<sup>[6-8]</sup>. Neuropathic pain interferes with activities of daily living and work performance, impairs mood, decreases quality of living and generates three-fold increases in health care costs relative to matched controls<sup>[9]</sup>. NP conditions involve spontaneous (paroxysmal or ongoing) and stimulus-evoked (*e.g.*, mechanical and thermal) symptoms; continuous or intermittent spontaneous pain is frequently described as burning, stabbing, shooting or shock-like; stimulus-evoked pain includes allodynia (pain in response to non-painful stimulation, extreme sensitivity to touch) and hyperalgesia (enhanced response to painful stimuli); NP can also involve tingling and numbness<sup>[1,6,9]</sup>.

TN involves irritation or compression of the 5<sup>th</sup> cranial nerve (trigeminal nerve), which evokes paroxysmal episodic stabbing pain of the facial area. Classically, pain is described as a sharp, shooting, electric shock-like, unilateral pain with acute onset and termination in distribution of the trigeminal nerve; this usually involves the V2 (maxillary) and V3 (mandibular) divisions but is rare in the V1 (ocular) division<sup>[10]</sup>. TN has an incidence of 4-28 per 100000 person years<sup>[11,12]</sup>. It can arise due to vascular alterations, non-vascular lesions, or tumours and other skull base abnormalities which exert pressure on the trigeminal nerve located in the CPA. Trigeminal NP is more continuous, and is characterized as burning, aching, throbbing<sup>[10]</sup>. VS, or acoustic neuromas, are the most frequent CPA tumour to cause TN-like symptoms<sup>[13]</sup>. Non-trigeminal nociceptive input, concurrent with induced masticatory responses (*i.e.*, hypoglossal, spinal accessory, facial, glossopharyngeal and vagal motor centers), may contribute to the wide constellation of symptoms seen in orofacial pain patients<sup>[14]</sup>. The distinction between TN and trigeminal NP is important, as there are different treatment recommendations for each<sup>[1,10]</sup>.

### Current algorithms

Treatment guidelines and decision rules improve patient outcomes. Recent literature providing strategies for the treatment of NP include the consensus statement and guidelines from the CPS<sup>[6]</sup>, as well as the Neuropathic Pain Special Interest Group of the International Associa-

tion for the Study of Pain<sup>[15,16]</sup>. It needs to be emphasized that most recommendations have been derived from studies on PHN and DN. The European Federation of Neurological Societies (EFNS) Task Force included an approach to the management of NP pain associated with damage to the trigeminal nerve<sup>[17]</sup>.

There remains a paucity of information on how to assess, diagnose and treat pain in patients with VS or other skull base tumours, pain that appears to be of neuropathic origin and typically resides within the head, face, neck and glossopharyngeal area. In order to provide guidance relating to this specialized patient population, we constructed an algorithm for NP and craniofacial pain, including TN, based on Canadian contemporary standards of care and existing NP treatment algorithms. Recommendations consider pharmaceutical agents with evidence of efficacy in neuropathic pain, patient tolerability of the dose range expected to be needed, and actual therapeutic efficacy observed within our clinical practice. To date, there has not been a published tool that provides a clearly-defined algorithm for the assessment and treatment of NP in patients with skull base tumours. It is intended that this report provides guidance to primary care practitioners treating NP in patients with skull base tumours, using specific drugs or combinations of drugs, to improve outcomes in clinical practice with respect to patient self-reports of pain and quality of life.

## CHARACTERIZATION OF PAIN ASSOCIATED WITH SKULL BASE TUMOURS

Skull base tumours involve the proliferation of abnormal cells in the part of the brain that meets the base of the skull. Symptoms of skull base tumours may include twitching, paralysis or facial pain. Craniofacial NP disorders include neuropathies, neuromas and neuralgias. Although significant interpractitioner and institutional variability exists, facial neuropathy and trigeminal nerve disturbances are relatively uncommon in comparison with unilateral hearing loss, tinnitus and vertigo or even idiopathic headache<sup>[18,19]</sup>. Even if uncommon, when a patient presents with what appears to be NP, the origin of the pain and appropriate treatment actions may be more difficult to determine than the existence of pain. At the very least, the challenge is to identify where the pain is coming from and to distinguish it from idiopathic headaches. Headaches are present in 50%-60% of patients with unilateral VS at the time of diagnosis<sup>[4,20]</sup>. Clinically, headaches that are unresponsive to over-the-counter analgesics may be a subtle cue that the pain originates from compression of the cranial nerves by the tumour<sup>[21]</sup>. It has been reported that 3% to 45% of patients with CPA VS experience facial paresis, facial neuropathy and trigeminal nerve disturbances (hypoesthesia, paresthesia, and neuralgia) due to compression by the tumour on the ipsilateral<sup>[13,22]</sup>, and less commonly, on contralateral<sup>[23]</sup> cra-

nial nerves; up to 93% are at risk secondary to irradiation treatment<sup>[24-27]</sup>.


The incidence of pain following SRT treatment is common. It has been estimated that 93% of lesions treated with SRT leave the patient at risk of radiation-induced TN<sup>[26]</sup>. Other centers report that trigeminal symptoms occur in  $\geq 3\%$  of patients whose tumours approach the level of the trigeminal nerve<sup>[25]</sup>. It is also important to distinguish NP from pain associated with hydrocephalus which may occur following tumour irradiation<sup>[19]</sup>. As such, patients presenting with pain should be referred for a magnetic resonance imaging (MRI) or computerized tomography scan. It is evident, given the diversity of pain mechanisms and individual patient responses that no single drug works for all NP states. In this respect, successful management of pain syndromes first necessitates accurate assessment.

## PHARMACOLOGICAL MANAGEMENT OF NEUROPATHIC PAIN


Assessment of pain and associated symptoms is necessary for diagnosis and management of NP. The self-report version of the Leeds Assessment of Neuropathic Symptoms and Signs scale and the French Neuropathic Pain Group clinician-administered questionnaire called DN4 may be used, based on their high sensitivity and selectivity<sup>[1,28,29]</sup>. The use of these tools is meant to complement but not replace clinical judgment.

Current treatment guidelines provide an evidence-based approach to the treatment of NP. Treatment guidelines have been developed based on data collected from randomized controlled clinical trials of anticonvulsants (carbamazepine, oxcarbazepine), tricyclic antidepressants (TCAs) (amitriptyline, nortriptyline), serotonin-norepinephrine reuptake inhibitors (SNRIs) (duloxetine, venlafaxine), calcium channel ligands (gabapentin, pregabalin), local anesthetics (5% lidocaine patch), opioids (morphine, methadone) and opioid-like hybrid drugs (tramadol)<sup>[6,16,17]</sup>. It should be mentioned that if the 5% lidocaine patch is unavailable, a lidocaine gel formulation or compounded cream may be substituted, although limited efficacy in non-post-herpetic pain has been reported<sup>[30]</sup>. A general overview of our suggested algorithm for the management of NP in our clinic is presented in Figures 1 and 2.

The algorithms presented below are based on published clinical guidelines to simplify the management and evaluation of NP in patients with lateral skull base tumours. Suggested first, second and third line agents [determined by efficacy, indicated by number-needed-to-treat (NNT), and patient tolerability as indicated by number-needed-to-harm (NNH), or side-effects] are listed in Figure 3. NNT and NNH vary according to the etiology of the pain and reference consulted<sup>[31,32]</sup>. Algorithms for individual pharmacological agents include initial starting doses, titration doses, temporal intervals and maximal dosing schedules<sup>[33]</sup>. In addition, TCAs, SNRIs, gabapentinoids, opioid analgesics and tramadol must all be used



**Figure 1** Assessment and management of neuropathic pain. NP: Neuropathic pain; PTSS: Pain treatment satisfaction scale.



**Figure 2** Management of neuropathic pain or trigeminal neuralgia in the Atlantic Lateral Skull Base Clinic. NP: Neuropathic pain; TN: Trigeminal neuralgia; SNRI: Serotonin-norepinephrine reuptake inhibitor.

with caution in elderly patients because of the risk of falls and cognitive impairment<sup>[15,16]</sup>. Recommendations are also made regarding the duration of adequate trials at maximum tolerated dosages to evaluate the impact on self-reported pain.

### Initial and subsequent agents

For management of classical TN (characterized by paroxysmal, unilateral pain), carbamazepine is the first choice, but otherwise it is not used; this agent has a NNT around


1.9 with virtually complete pain relief<sup>[6,10,34-36]</sup> (Table 1). Oxcarbazepine can, and should, be substituted for carbamazepine if there is an unacceptable side effect profile<sup>[10,34-36]</sup> (Table 1).

For management of NP, there are several TCAs available, but amitriptyline and nortriptyline are commonly used, and exhibit NNT values of 1.3-3.6<sup>[32,37-39]</sup>; NNH values are 3-6 for minor, and 14-28 for major, harm. The analgesic properties of TCAs are independent of their antidepressant effects, and several mechanisms, in

**Table 1 Summary of drugs used for management of trigeminal neuralgia and neuropathic pain**

Drug	Action	Dosing	Common side effects <sup>1</sup>
CBZ	Blocks Na <sup>+</sup> and Ca <sup>2+</sup> channels	300-1000 mg; 100 mg BID initially, increase by 200 mg weekly Adequate trial 8-12 wk, 2 wk at maximal dose	Drowsiness, ataxia, headaches, nausea, vomiting, constipation, blurred vision, rash Drug interactions Taper doses when discontinuing
OXC	Keto derivative of CBZ, same actions	Equivalent efficacy to CBZ; 300-2400 mg; 300 mg BID initially, increase by 600 mg weekly Adequate trial 8-12 wk, 2 wk at maximal dose	Improved tolerability compared to CBZ Vertigo, fatigue, dizziness, nausea, hyponatraemia in high doses No major drug interactions Taper doses when discontinuing
TCAs or tricyclic antidepressants: nortriptyline, amitriptyline	Block NA and 5-HT reuptake, block Na <sup>+</sup> channels, interact with several neurotransmitter systems	Nortriptyline 10 mg (elderly) or 25 mg (adult) at bedtime; increase dose by 10 or 25 mg every 3-7 d; up to 75-100 mg daily Adequate trial 6-8 wk, 2 wk at maximal dose Amitriptyline doses similar	Nortriptyline is better tolerated than amitriptyline Dry mouth, constipation, blurred vision, sedation, orthostatic hypotension Taper doses when discontinuing
SNRIs or serotonin-noradrenaline reuptake inhibitors: Duloxetine, Venlafaxine	Similar actions to TCAs, but fewer interactions with receptor systems	Duloxetine: 60 mg/d, increase to 120 mg after 1 wk Venlafaxine: 75 mg/d, increase to 225 mg over 3 wk Adequate trial 4-6 wk, 2 wk at maximum dose	Headache, nausea, dry mouth, sleepiness, fatigue, constipation, dizziness, decreased appetite, and increased sweating. Taper doses when discontinuing Drowsiness, dizziness, weakness, feeling nervous, tinnitus, increased sweating, blurred vision, dry mouth, changes in appetite or weight, facial flushing, mild nausea, constipation, sexual side-effects Taper doses over 7-10 d when discontinuing
Gabapentin	Ca <sup>2+</sup> channel modulator	100-300 mg TID, increase dose every 1-7 d; maximum dose 3600 mg daily Adequate trial 3-8 wk titration, 2-8 wk at maximal dose	Dizziness, sedation, weight gain, weakness, tiredness, nausea, diarrhea, constipation, blurred vision, headache, breast swelling, dry mouth, fatigue, myalgia, loss of balance or coordination Taper doses when discontinuing
Pregabalin	Ca <sup>2+</sup> channel modulator	50 mg OD or 25 mg BID, double dose each week; maximum daily dose 600 mg Adequate trial is 4 wk at maximal dose	Dizziness, drowsiness, loss of balance or coordination, problems with memory or concentration, anxiety, depersonalization, hypertonia, hypesthesia, decreased libido, nystagmus, paresthesia, twitching, breast swelling, tremors, dry mouth, constipation Taper doses when discontinuing (minimum of one week)
Tramadol	Inhibits NA and 5-HT reuptake, binds opioid receptors	50 mg BID, increase by 50-100 mg daily in divided doses over 3-7 d as tolerated; 400 mg is maximum dose (300 mg in elderly) Adequate trial is 4 wk at maximum dose	Dizziness, spinning sensation, constipation, upset stomach, headache, drowsiness, feeling nervous or anxious
Morphine	Interacts with mu opioid receptors in spinal cord and brain, regulates synaptic activity in pain pathways	10-15 mg q4h or prn (equianalgesic doses for other opioids); after 1-2 wk, convert to long-acting opioid (e.g., CR hydromorphone) Titration allows for dose escalation Adequate trial is 4-6 wk	Sedation, pruritus, constipation, diarrhea, weight loss, nausea, vomiting, stomach pain, loss of appetite, flushing (warmth, redness, tingling), headache, dizziness, spinning sensation, memory problems, sleep problems (insomnia), strange dreams Taper doses when discontinuing

<sup>1</sup>These are not a complete list of side effects and others may occur. CBZ: Carbamazepine; 5-HT: Serotonin; NA: Noradrenaline; OXC: Oxcarbazepine; q4h: Every four hours; BID: Twice a day; TCA: Tricyclic antidepressant; SNRI: Serotonin-norepinephrine reuptake inhibitors; prn: Pro re nata; TID: Three times a day; OD: Once daily; CR: Continuous release.



**Figure 3 Treatment algorithm for trigeminal neuralgia or neuropathic craniofacial pain.** TN: Trigeminal neuralgia; SNRIs: Serotonin-norepinephrine reuptake inhibitors; TCA: Tricyclic antidepressant.

addition to blockage of serotonin and norepinephrine reuptake, are involved in their actions<sup>[40]</sup>. It should be emphasized that caution must be employed with the use of TCAs in older patients because of anticholinergic adverse effects, sedation, risk of falls, and risk of cardiac toxicity. Nortriptyline, a metabolite of amitriptyline with similar pharmacological effectiveness reflected in similar NNT values<sup>[6,37,39]</sup>, has a lower incidence of adverse effects compared to amitriptyline<sup>[32,41]</sup>. The lowest effective dose of TCA should be used in NP patients, avoiding patients with ischemic heart disease or increased risk of cardiac death<sup>[15]</sup> (Table 1). Where available, lidocaine medicated plaster (NNT = 4.4, NNH = 29), a topical formulation, can be considered first-line treatment if the pain is focal and there are tolerability issues for oral for-

mulations<sup>[16,37]</sup>.

The EFNS and Neuropathic Pain Special Interest Group (NeuPSIG) recommend the two SNRIs, duloxetine and venlafaxine, as first-line options, while Canadian guidelines consider these second-line options for treatment of NP. Duloxetine (NNT = 5-6; NNH = 7-9 minor, 13-15 major) has primarily been examined in DN, while venlafaxine (NNT = 3-5, NNH = 7-9 minor, 16 major) has been examined in a broader range of NP conditions<sup>[31,37-39]</sup> (Table 1). Venlafaxine may be employed in combination with gabapentin to increase its efficacy<sup>[16,30,42,43]</sup>.

The Consensus Statement and Guidelines from the Canadian Pain Society list gabapentin and pregabalin as first-line agents in the treatment of NP<sup>[6]</sup>. Most of the literature and guidelines for NP are based on PHN and DN and may not be applicable to all NP conditions<sup>[9,15]</sup>. Gabapentin (NNT = 4.1-6.4, NNH = 3.7) and pregabalin (NNT = 3.3-6.0, NNH = 3.7-8.8) (Table 1)<sup>[32,37,38]</sup> can be considered as first- or second-line agents for management of orofacial NP associated with skull base tumours. The EFNS Task Force has identified the usefulness of combination therapy, including TCA-gabapentin for the management of NP and TN<sup>[17]</sup> (Figure 2).

Gabapentin and pregabalin decrease the release of several neurotransmitters involved in pain through binding to the  $\alpha 2\text{-}\delta$  subunit of voltage-gated calcium channels, synaptic  $\gamma$ -aminobutyric acid modulation and synaptogenesis<sup>[44]</sup>. Side effects of gabapentin include dry mouth, dizziness, gastrointestinal disturbances and cognitive impairment. There is some evidence that the efficacy of gabapentin is increased in painful DN, when combined with venlafaxine or morphine<sup>[31,45]</sup>. Combination of gabapentin with nortriptyline was superior to either drug alone, and combination of gabapentin with opioids has shown increased efficacy for the treatment of TN and NP<sup>[17]</sup> (Figure 2).

### Third-line agents

Tramadol, a hybrid drug with SNRI and  $\mu$ -opioid agonist properties, has a NNT = 3.4-4.9, and a NNH = 7.7; it should not be combined with TCAs due to increased risk of serotonin syndrome<sup>[32,37]</sup> (Table 1). Opioids are generally safe if titrated slowly<sup>[46]</sup>. While there is some evidence to support opioids in NP<sup>[15,32,37,39]</sup>, opioids as primary therapy are not always effective, and combination therapy may be needed<sup>[31,42,44]</sup>. The analgesic efficacy of a combination of morphine and gabapentin was increased compared to each individual agent in patients with PHN and painful DN; however, while the maximal tolerated doses were lower in combination therapy, there was report of increased adverse effects<sup>[16]</sup>.

There are many liabilities and controversies surrounding the use of opioids in those with chronic non-cancer pain<sup>[28,46]</sup>. For treating NP with opioids we follow recommendations of the 2010 National Opioid Use Guideline Group<sup>[46]</sup>; referral to a chronic pain service is indicated when combination therapies involve primary and adjunct treatment beyond the common agents listed.

### Other agents

Although beyond the scope of this document and better reserved for pain specialty clinics, interventional procedures, compounded drugs (such as carbamazepine, gabapentin, antidepressants, lidocaine) delivered as topical formulations for the orofacial region<sup>[47]</sup>, invasive techniques (may include intravenous lidocaine)<sup>[6]</sup> or intradermal botulinum toxin<sup>[16]</sup> may be considered. Several recent case series reports and a controlled study support the efficacy and safety of botulinum toxin for TN<sup>[48-50]</sup>. Future treatment guidelines will position these options within current schemes.

## CASE EXAMPLES

Two case examples of the presentation and treatment of neuropathic pain resulting from VS are outlined below.

### Case 1

Presentation and diagnosis: RG, a 72-year-old retired fireman, presented to the Atlantic Lateral Skull Base Clinic in August 2010 with progressive right-sided hearing loss and tinnitus over 1-2 years with a new presentation of numbness and tingling on the right side of his face including the tip of his tongue. MRI revealed a right-sided cerebellar pontine angle tumour (25 mm  $\times$  27 mm) in the axial plane with mass effect on the brainstem consistent with a vestibular schwannoma/acoustic neuroma. RG opted for stereotactic radiation therapy (SRT) over surgery or a more conservative "wait and scan" approach.

Symptoms prior to treatment: In December 2010, prior to SRT therapy, lidocaine hydrochloride swish and spit was ineffective in controlling symptoms. In January 2011, he reported exacerbation of symptoms on the right side of his face, dysesthesia rather than numbness, constant burning involving the tip of the tongue and the bottom of the tongue on the right hand side; symptoms were exacerbated by eating. He had lost 30 pounds due to decreased food intake. Pregabalin 75 mg twice a day and dexamethasone 4 mg every morning with breakfast was prescribed. Over the next month, pregabalin was increased to 150 mg twice a day as he noted some early improvement. The pregabalin dose was again increased to 300 mg and RG gained 3 pounds.

Treatment and follow up: SRT began in February 2011. RG noticed an immediate improvement in his tinnitus. In March 2011, gabapentin replaced pregabalin and was increased to 1800 mg/d while dexamethasone weaning began. RG experienced hypotension in response to the increase in gabapentin and the dose was decreased to 300 mg three times a day. SRT was completed in April 2011. At this time, RG reported increased facial pain, so gabapentin was increased to 1200 and then 1500 mg/d. RG was not reporting any relief from symptoms, and numbness and right-sided trigeminal neuralgia persisted. In October 2011, he was weaned off gabapentin, and amitriptyline was prescribed for facial pain 10 mg every night, and increased to 20 mg after 1 wk. In November

2011, he reported side effects which included sleepiness and nausea. His family doctor changed the prescription to nortriptyline 10 mg for 5 d, then 20 mg/d for 5 d, and then 30 mg/d, with a plan to increase this again in the next 5 d to 50 mg/d, his current dose in February 2012. He has improved facial pain and reports improvements in eating and tasting food. He still has some paresthesia and dysesthesia in the right side of his face in the trigeminal distribution in all 3 branches. His vestibular schwannoma has shrunk by 0.5 cm.

## Case 2

**Presentation and diagnosis:** DB aged 71, presented to the Atlantic Lateral Skull Base Clinic in March 2008 reporting a recent history of loud noises prompting headaches. She subsequently had an audiogram which showed slight decrease in useful hearing on the right side compared to the left. An MRI (November 2007) revealed a small right-sided acoustic neuroma (18 mm × 9 mm in the axial plane). General ears, nose and throat exam was normal, other than the slight decrease in useful hearing; no tinnitus or vertigo was reported. The Unterberger stepping test revealed a moderate pulling to the right; cerebellar and oculomotor function was normal. Her facial nerve was completely normal with no facial weakness, facial twitching or facial numbness. She remained stable for a year.

**Symptoms prior to treatment:** In July 2009, her hearing deteriorated, to 60% of normal, and she was experiencing disequilibrium. MRI revealed slight growth in the neuroma (20 mm in longest dimension, 13 mm perpendicular to the petrous apex). A subsequent MRI in November 2009 showed further growth (20 mm × 15 mm).

**Treatment and follow up:** DB was referred for SRT, and this was completed June 2010. MRI in January 2011 revealed central cystic changes in tumour composition consistent with SRT. DB reported no new clinical symptoms other than further decrease in hearing on the right side.

In March 2011, new neurological symptoms emerged. Initially she described the right side of her tongue as heavy. She began to have some right-sided painful secondary paresthesia on her face in cheek area, upper and lower lips and tongue [maxillary (V2) and mandibular (V3) divisions of the trigeminal nerve] that she describes as a persistent burning sensation that is made worse by eating. She also described discomfort in her right eye and right-sided facial twitching. MRI in March 2011 showed swelling of the tumour in the right CPA consistent with SRT-induced changes. She was started on gabapentin 300 mg once daily for one week, increasing the dose to twice daily the second week, and thrice daily for the third week if tolerated.

In November of 2011 she returned. Gabapentin 300 mg once daily reduced her symptoms; however, it was associated with intolerable constipation and myalgia in the upper arms. She was weaned off gabapentin and her constipation and myalgia dissipated. The right-sided acoustic neuroma had further decreased in size, approximately 18 mm × 12 mm (May 2013) compared to 22 mm × 16 mm

(April 2012), yet she continued to experience persistent chronic orofacial pain. While it is possible that DB would be refractory to other pharmacologic interventions, she has declined further interventions, despite the pain and effect on her quality of life.

## CONCLUSION

The present report describes evidence for application of pain management strategies in patients with VS. A decision tool and treatment algorithm is presented to facilitate evaluation and management of patients with NP resulting from skull base disorders. A pharmacological algorithm, with primary and adjuvant treatment, summarizes pharmaceutical choice and management of NP and TN in patients with VS and can be made available in the clinic for quick review by the treating physicians. This instrument is intended to guide treatment of neuropathic/trigeminal pain in patients in acute care or ambulatory clinical settings. Essential to the comprehensive management of patients is evaluation of the effects of the intervention on quality-of-life and patient satisfaction with pain management. Patient satisfaction with pain management, using the Pain Treatment Satisfaction Scale<sup>[51]</sup>, as well as the American Pain Society Satisfaction Survey, was influenced by effectiveness of medication on pain severity, independent of initial pain intensity, and by communication<sup>[52]</sup>. Comparison of outcome data (self-reported pain and quality of life) will elucidate the use of standardized approaches to managing pain among patients with specific skull base etiology. The purpose of the present treatment algorithm was to develop a common scheme that may be utilized by beginning practitioners for treating this relatively uncommon, but clinically challenging, condition.

## REFERENCES

- 1 **Baron R**, Binder A, Wasner G. Neuropathic pain: diagnosis, pathophysiological mechanisms, and treatment. *Lancet Neurol* 2010; **9**: 807-819 [PMID: 20650402 DOI: 10.1016/S1474-4422(10)70143-5]
- 2 **Elias WJ**, Burchiel KJ. Trigeminal neuralgia and other neuropathic pain syndromes of the head and face. *Curr Pain Headache Rep* 2002; **6**: 115-124 [PMID: 11872182 DOI: 10.1007/s11916-002-0007-8]
- 3 **Zakrzewska JM**. Diagnosis and differential diagnosis of trigeminal neuralgia. *Clin J Pain* 2002; **18**: 14-21 [PMID: 11803298 DOI: 10.1097/00002508-200201000-00003]
- 4 **Kutz JW**. Acoustic neuroma. 2011. Available from: URL: <http://emedicine.medscape.com/article/882876-overview>
- 5 **Statistics Canada**. Canada's population estimates. 2011. Available from: URL: <http://www.statcan.gc.ca/daily-quotidien/100929/t100929b3-eng.htm>
- 6 **Moulin DE**, Clark AJ, Gilron I, Ware MA, Watson CP, Sessle BJ, Coderre T, Morley-Forster PK, Stinson J, Boulanger A, Peng P, Finley GA, Taenzer P, Squire P, Dion D, Chokan A, Gilani A, Gordon A, Henry J, Jovey R, Lynch M, Mailis-Gagnon A, Panju A, Rollman GB, Velly A. Pharmacological management of chronic neuropathic pain-consensus statement and guidelines from the Canadian Pain Society. *Pain Res Manag* 2007; **12**: 13-21 [PMID: 17372630]
- 7 **Torrance N**, Smith BH, Bennett MI, Lee AJ. The epidemiol-

- ogy of chronic pain of predominantly neuropathic origin. Results from a general population survey. *J Pain* 2006; **7**: 281-289 [PMID: 16618472 DOI: 10.1016/j.jpain.2005.11.008]
- 8 **Smith BH**, Torrance N. Epidemiology of neuropathic pain and its impact on quality of life. *Curr Pain Headache Rep* 2012; **16**: 191-198 [PMID: 22395856 DOI: 10.1007/s11916-012-0256-0]
- 9 **Gilron I**, Watson CP, Cahill CM, Moulin DE. Neuropathic pain: a practical guide for the clinician. *CMAJ* 2006; **175**: 265-275 [PMID: 16880448 DOI: 10.1503/cmaj.060146]
- 10 **Zakrzewska JM**. Medical management of trigeminal neuropathic pains. *Expert Opin Pharmacother* 2010; **11**: 1239-1254 [PMID: 20426709 DOI: 10.1517/14656561003767449]
- 11 **Koopman JS**, Deilman JP, Huygen FJ, de Mos M, Martin CG, Sturkenboom MC. Incidence of facial pain in the general population. *Pain* 2009; **147**: 122-127 [PMID: 19783099 DOI: 10.1016/j.pain.2009.08.023]
- 12 **van Hecke O**, Austin SK, Khan RA, Smith BH, Torrance N. Neuropathic pain in the general population: a systematic review of epidemiological studies. *Pain* 2014; **155**: 654-662 [PMID: 24291734 DOI: 10.1016/j.pain.2013.11.013]
- 13 **Matsuka Y**, Fort ET, Merrill RL. Trigeminal neuralgia due to an acoustic neuroma in the cerebellopontine angle. *J Orofac Pain* 2000; **14**: 147-151 [PMID: 11203749]
- 14 **Türp JC**, Kowalski CJ, O'Leary N, Stohler CS. Pain maps from facial pain patients indicate a broad pain geography. *J Dent Res* 1998; **77**: 1465-1472 [PMID: 9649175 DOI: 10.1177/00220345980770061101]
- 15 **Dworkin RH**, O'Connor AB, Backonja M, Farrar JT, Finnerup NB, Jensen TS, Kalso EA, Loeser JD, Miaskowski C, Nurmikko TJ, Portenoy RK, Rice AS, Stacey BR, Treede RD, Turk DC, Wallace MS. Pharmacologic management of neuropathic pain: evidence-based recommendations. *Pain* 2007; **132**: 237-251 [PMID: 17920770 DOI: 10.1016/j.pain.2007.08.033]
- 16 **Dworkin RH**, O'Connor AB, Audette J, Baron R, Gourlay GK, Haanpää ML, Kent JL, Krane EJ, LeBel AA, Levy RM, Mackey SC, Mayer DC, Miaskowski C, Raja SN, Rice AS, Schmäder KE, Stacey B, Stanos S, Treede RD, Turk DC, Walco GA, Wells CD. Recommendations for the pharmacological management of neuropathic pain: an overview and literature update. *Mayo Clin Proc* 2010; **85**: S3-S14 [PMID: 20194146 DOI: 10.4065/mcp.2009.0649]
- 17 **Attal N**, Cruccu G, Baron R, Haanpää M, Hansson P, Jensen TS, Nurmikko T. EFNS guidelines on the pharmacological treatment of neuropathic pain: 2010 revision. *Eur J Neurol* 2010; **17**: 1113-e88 [PMID: 20402746 DOI: 10.1111/j.1468-1331.2010.02999.x]
- 18 **Morrison GA**, Sterkers JM. Unusual presentations of acoustic tumours. *Clin Otolaryngol Allied Sci* 1996; **21**: 80-83 [PMID: 8674229 DOI: 10.1111/j.1365-2273.1996.tb01030.x]
- 19 **Shin YJ**, Lapeyre-Mestre M, Gafsi I, Cognard C, Deguine O, Tremoulet M, Frayssé B. Neurotological complications after radiosurgery versus conservative management in acoustic neuromas: a systematic review-based study. *Acta Otolaryngol* 2003; **123**: 59-64 [PMID: 12625575 DOI: 10.1081/003655402100028084]
- 20 **Walter J**, Roland PS, Isaacson B. Acoustic Neuroma. Medscape 2011. Available from: URL: <http://emedicine.medscape.com/article/882876-overview#showall>
- 21 **Hankinson T**, Haque R, Kellner CP, Bruce JN, Sinson GP, Reiter GT. Skull Base Tumours. Medscape 2011. Available from: URL: <http://emedicine.medscape.com/article/250237-overview>
- 22 **Matthies C**, Samii M. Management of 1000 vestibular schwannomas (acoustic neuromas): clinical presentation. *Neurosurgery* 1997; **40**: 1-10 [DOI: 10.1097/00006123-199701000-00001]
- 23 **Chamadoira C**, Cerejo A, Duarte F, Vaz R. [Trigeminal neuralgia caused by contra lateral cerebellopontine angle tumor. A case report]. *Neurocirugia (Astur)* 2010; **21**: 50-52 [PMID: 20186375]
- 24 **Chung WY**, Liu KD, Shiao CY, Wu HM, Wang LW, Guo WY, Ho DM, Pan DH. Gamma knife surgery for vestibular schwannoma: 10-year experience of 195 cases. *J Neurosurg* 2005; **102** Suppl: 87-96 [PMID: 15662787 DOI: 10.3171/jns.2005.102.s\_supplement.0087]
- 25 **Lunsford LD**, Niranjan A, Flickinger JC, Maitz A, Kondziolka D. Radiosurgery of vestibular schwannomas: summary of experience in 829 cases. *J Neurosurg* 2005; **102** Suppl: 195-199 [PMID: 15662809 DOI: 10.3171/jns.2005.102.s\_supplement.0195]
- 26 **Combs SE**, Thilmann C, Debus J, Schulz-Ertner D. Long-term outcome of stereotactic radiosurgery (SRS) in patients with acoustic neuromas. *Int J Radiat Oncol Biol Phys* 2006; **64**: 1341-1347 [PMID: 16464537 DOI: 10.1016/j.ijrobp.2005.10.024]
- 27 **Sughrue ME**, Yang I, Han SJ, Aranda D, Kane AJ, Amols M, Smith ZA, Parsa AT. Non-audiofacial morbidity after Gamma Knife surgery for vestibular schwannoma. *Neurosurg Focus* 2009; **27**: E4 [PMID: 19951057 DOI: 10.3171/2009.9.FOCUS09198]
- 28 **Bennett MI**, Smith BH, Torrance N, Potter J. The S-LANSS score for identifying pain of predominantly neuropathic origin: validation for use in clinical and postal research. *J Pain* 2005; **6**: 149-158 [PMID: 15772908 DOI: 10.1016/j.jpain.2004.11.007]
- 29 **Bouhassira D**, Attal N, Alchaar H, Boureau F, Brochet B, Bruxelle J, Cunin G, Fermanian J, Ginies P, Grun-Overdyking A, Jafari-Schluep H, Lanteri-Minet M, Laurent B, Mick G, Serrie A, Valade D, Vicaute E. Comparison of pain syndromes associated with nervous or somatic lesions and development of a new neuropathic pain diagnostic questionnaire (DN4). *Pain* 2005; **114**: 29-36 [PMID: 15733628 DOI: 10.1016/j.pain.2004.12.010]
- 30 **O'Connor AB**, Dworkin RH. Treatment of neuropathic pain: an overview of recent guidelines. *Am J Med* 2009; **122**: S22-S32 [PMID: 19801049]
- 31 **Finnerup NB**, Otto M, McQuay HJ, Jensen TS, Sindrup SH. Algorithm for neuropathic pain treatment: an evidence based proposal. *Pain* 2005; **118**: 289-305 [PMID: 16213659 DOI: 10.1016/j.pain.2005.08.013]
- 32 **Jefferies K**. Treatment of neuropathic pain. *Semin Neurol* 2010; **30**: 425-432 [PMID: 20941675 DOI: 10.1055/s-0030-1267286]
- 33 **E-CPS**. Compendium of pharmaceuticals and specialties. 2012 (online). Available from: URL: <http://www.pharmacists.ca/index.cfm/function/store/PublicationDetail.cfm?pub=5>
- 34 **Cruccu G**, Gronseth G, Alksne J, Argoff C, Brainin M, Burchiel K, Nurmikko T, Zakrzewska JM. AAN-EFNS guidelines on trigeminal neuralgia management. *Eur J Neurol* 2008; **15**: 1013-1028 [PMID: 18721143 DOI: 10.1111/j.1468-1331.2008.02185.x]
- 35 **Cruccu G**, Truini A. Refractory trigeminal neuralgia. Non-surgical treatment options. *CNS Drugs* 2013; **27**: 91-96 [PMID: 23225488 DOI: 10.1007/s40263-012-0023-0]
- 36 **Jorns TP**, Zakrzewska JM. Evidence-based approach to the medical management of trigeminal neuralgia. *Br J Neurosurg* 2007; **21**: 253-261 [PMID: 17612914 DOI: 10.1080/02688690701219175]
- 37 **Lindsay TJ**, Rodgers BC, Savath V, Hettinger K. Treating diabetic peripheral neuropathic pain. *Am Fam Physician* 2010; **82**: 151-158 [PMID: 20642268]
- 38 **Moore RA**, Derry S, Aldington D, Cole P, Wiffen PJ. Amitriptyline for neuropathic pain and fibromyalgia in adults. *Cochrane Database Syst Rev* 2012; **12**: CD008242 [DOI: 10.1002/14651858]
- 39 **Watson CP**, Gilron I, Sawynok J, Lynch ME. Nontricyclic antidepressant analgesics and pain: are serotonin norepinephrine reuptake inhibitors (SNRIs) any better? *Pain* 2011; **152**: 2206-2210 [PMID: 21723037 DOI: 10.1016/j.pain.2011.05.032]
- 40 **Verdu B**, Decosterd I, Buclin T, Stiefel F, Berner A. Antidepressants for the treatment of chronic pain. *Drugs* 2008; **68**: 2611-2632 [PMID: 19093703 DOI: 10.2165/0003495-200868180-00007]
- 41 **Watson CP**, Vernich L, Chipman M, Reed K. Nortriptyline

- versus amitriptyline in postherpetic neuralgia: a randomized trial. *Neurology* 1998; **51**: 1166-1171 [PMID: 9781549 DOI: 10.1212/WNL.51.4.1166]
- 42 **Backonja MM**, Irving G, Argoff C. Rational multidrug therapy in the treatment of neuropathic pain. *Curr Pain Headache Rep* 2006; **10**: 34-38 [PMID: 16499828 DOI: 10.1007/s11916-006-0007-1]
  - 43 **Vorobeychik Y**, Gordin V, Mao J, Chen L. Combination therapy for neuropathic pain: a review of current evidence. *CNS Drugs* 2011; **25**: 1023-1034 [PMID: 22133325 DOI: 10.2165/11596280-000000000-00000]
  - 44 **Taylor CP**. Mechanisms of analgesia by gabapentin and pregabalin--calcium channel alpha2-delta [Cavalpha2-delta] ligands. *Pain* 2009; **142**: 13-16 [PMID: 19128880 DOI: 10.1016/j.pain.2008.11.019]
  - 45 **Gilron I**, Bailey JM, Tu D, Holden RR, Weaver DF, Houlton RL. Morphine, gabapentin, or their combination for neuropathic pain. *N Engl J Med* 2005; **352**: 1324-1334 [PMID: 15800228 DOI: 10.1056/NEJMoa042580]
  - 46 **NOUGG (National Opioid Use Guideline Group)**. Canadian Guideline for Safe and Effective Use of Opioids for Chronic Non-Cancer Pain. 2010. Available from: URL: <http://nationalpaincentre.mcmaster.ca/opioid/>
  - 47 **Heir G**, Karolchek S, Kalladka M, Vishwanath A, Gomes J, Khatri R, Nasri C, Eliav E, Ananthan S. Use of topical medication in orofacial neuropathic pain: a retrospective study. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2008; **105**: 466-469 [PMID: 18329583 DOI: 10.1016/j.tripleo.2007.09.030]
  - 48 **Zúñiga C**, Díaz S, Piedimonte F, Micheli F. Beneficial effects of botulinum toxin type A in trigeminal neuralgia. *Arq Neuropsiquiatr* 2008; **66**: 500-503 [PMID: 18813708 DOI: 10.1590/S0004-282X2008000400012]
  - 49 **Bohluli B**, Motamedi MH, Bagheri SC, Bayat M, Lassemi E, Navi F, Moharamnejad N. Use of botulinum toxin A for drug-refractory trigeminal neuralgia: preliminary report. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2011; **111**: 47-50 [PMID: 20674409 DOI: 10.1016/j.tripleo.2010.04.043]
  - 50 **Wu CJ**, Lian YJ, Zheng YK, Zhang HF, Chen Y, Xie NC, Wang LJ. Botulinum toxin type A for the treatment of trigeminal neuralgia: results from a randomized, double-blind, placebo-controlled trial. *Cephalalgia* 2012; **32**: 443-450 [PMID: 22492424 DOI: 10.1177/0333102412441721]
  - 51 **Evans CJ**, Trudeau E, Mertzanis P, Marquis P, Peña BM, Wong J, Mayne T. Development and validation of the Pain Treatment Satisfaction Scale (PTSS): a patient satisfaction questionnaire for use in patients with chronic or acute pain. *Pain* 2004; **112**: 254-266 [PMID: 15561380 DOI: 10.1016/j.pain.2004.09.005]
  - 52 **Carlson J**, Youngblood R, Dalton JA, Blau W, Lindley C. Is patient satisfaction a legitimate outcome of pain management? *J Pain Symptom Manage* 2003; **25**: 264-275 [PMID: 12614961 DOI: 10.1016/S0885-3924(02)00677-2]

**P- Reviewer:** Gottschalk A, Noll-Hussong M, Tao F

**S- Editor:** Ji FF **L- Editor:** A **E- Editor:** Liu SQ





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>

