


# World Journal of *Gastrointestinal Pharmacology and Therapeutics*

*World J Gastrointest Pharmacol Ther* 2017 May 6; 8(2): 90-154





**Contents**

Quarterly Volume 8 Number 2 May 6, 2017

**EDITORIAL**

- 90 Management of esophageal caustic injury  
*De Lusong MAA, Timbol ABG, Tuazon DJS*

**FRONTIER**

- 99 5-Aminosalicylates to maintain remission in Crohn's disease: Interpreting conflicting systematic review evidence  
*Gordon M*

**MINIREVIEWS**

- 103 Combination therapy for inflammatory bowel disease  
*Sultan KS, Berkowitz JC, Khan S*
- 114 Inflammatory bowel disease: Efficient remission maintenance is crucial for cost containment  
*Actis GC, Pellicano R*

**ORIGINAL ARTICLE**

**Case Control Study**

- 120 Thiol/disulphide homeostasis in celiac disease  
*Kaplan M, Ates I, Yuksel M, Ozderin Ozin Y, Alisik M, Erel O, Kayacetin E*

**Observational Study**

- 127 Correlation of rapid point-of-care vs send-out fecal calprotectin monitoring in pediatric inflammatory bowel disease  
*Rodriguez A, Yokomizo L, Christofferson M, Barnes D, Khavari N, Park KT*
- 131 Clinical and economic impact of infliximab one-hour infusion protocol in patients with inflammatory bowel diseases: A multicenter study  
*Viola A, Costantino G, Privitera AC, Bossa F, Lauria A, Grossi L, Principi MB, Della Valle N, Cappello M*

- 137 Interferon-free treatments in patients with hepatitis C genotype 1-4 infections in a real-world setting  
*Ramos H, Linares P, Badia E, Martín I, Gómez J, Almohalla C, Jorquera F, Calvo S, García I, Conde P, Álvarez B, Karpman G, Lorenzo S, Gozalo V, Vásquez M, Joao D, de Benito M, Ruiz L, Jiménez F, Sáez-Royuela F; Asociación Castellano y Leonesa de Hepatología (ACyLHE)*

**Randomized Controlled Trial**

- 147 Low dose oral curcumin is not effective in induction of remission in mild to moderate ulcerative colitis: Results from a randomized double blind placebo controlled trial  
*Kedia S, Bhatia V, Thareja S, Garg S, Mouli VP, Bopanna S, Tiwari V, Makharja G, Ahuja V*

## Contents

*World Journal of Gastrointestinal Pharmacology and Therapeutics*

Volume 8 Number 2 May 6, 2017

### ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Pharmacology and Therapeutics*, Dong Joon Kim, MD, PhD, Professor, Research Fellow, Department of Internal Medicine Center for Liver and Digestive Diseases, Hallym University Chuncheon Sacred Heart Hospital, Hallym University Hospital, Gangwon-do 200-704, South Korea

### AIM AND SCOPE

*World Journal of Gastrointestinal Pharmacology and Therapeutics* (*World J Gastrointest Pharmacol Ther*, *WJGPT*, online ISSN 2150-5349, DOI: 10.4292), is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

*WJGPT* covers topics concerning: (1) Clinical pharmacological research articles on specific drugs, concerning with pharmacodynamics, pharmacokinetics, toxicology, clinical trial, drug reactions, drug metabolism and adverse reaction monitoring, etc.; (2) Research progress of clinical pharmacology; (3) Introduction and evaluation of new drugs; (4) Experiences and problems in applied therapeutics; (5) Research and introductions of methodology in clinical pharmacology; and (6) Guidelines of clinical trial.

We encourage authors to submit their manuscripts to *WJGPT*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great basic and clinical significance.

### INDEXING/ABSTRACTING

*World Journal of Gastrointestinal Pharmacology and Therapeutics* is now indexed in PubMed, PubMed Central.

### FLYLEAF

#### I-IV Editorial Board

### EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*  
Responsible Electronic Editor: *Ya-Jing Lu*  
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Fang-Fang Ji*  
Proofing Editorial Office Director: *Xin-Xia Song*

#### NAME OF JOURNAL

*World Journal of Gastrointestinal Pharmacology and Therapeutics*

#### ISSN

ISSN 2150-5349 (online)

#### LAUNCH DATE

May 6, 2010

#### FREQUENCY

Quarterly

#### EDITOR-IN-CHIEF

**Hugh J Freeman, MD, FRCPC, FACP, Professor,**  
Department of Medicine (Gastroenterology), University of British Columbia, Hospital, 2211 Wesbrook Mall, Vancouver, BC V6T1W5, Canada

#### EDITORIAL BOARD MEMBERS

All editorial board members resources online at <http://www.wjgnet.com/2150-5349/editorialboard.htm>

#### EDITORIAL OFFICE

Xiu-Xia Song, Director  
*World Journal of Gastrointestinal Pharmacology and Therapeutics*  
Baishideng Publishing Group Inc  
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA  
Telephone: +1-925-2238242  
Fax: +1-925-2238243  
E-mail: [editorialoffice@wjgnet.com](mailto:editorialoffice@wjgnet.com)  
Help Desk: <http://www.f6publishing.com/helpdesk>  
<http://www.wjgnet.com>

#### PUBLISHER

Baishideng Publishing Group Inc  
7901 Stoneridge Drive, Suite 501,  
Pleasanton, CA 94588, USA  
Telephone: +1-925-2238242  
Fax: +1-925-2238243  
E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
Help Desk: <http://www.f6publishing.com/helpdesk>  
<http://www.wjgnet.com>

#### PUBLICATION DATE

May 6, 2017

#### COPYRIGHT

© 2017 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

#### SPECIAL STATEMENT

All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

#### INSTRUCTIONS TO AUTHORS

<http://www.wjgnet.com/bpg/gerinfo/204>

#### ONLINE SUBMISSION

<http://www.f6publishing.com>

## Case Control Study

# Thiol/disulphide homeostasis in celiac disease

Mustafa Kaplan, Ihsan Ates, Mahmut Yuksel, Yasemin Ozderin Ozin, Murat Alisik, Ozcan Erel, Ertugrul Kayacetin

Mustafa Kaplan, Mahmut Yuksel, Yasemin Ozderin Ozin, Ertugrul Kayacetin, Department of Gastroenterology, Turkey Yuksek Ihtisas Training and Research Hospital, 06500 Ankara, Turkey

Ihsan Ates, Department of Internal Medicine, Ankara Numune Training and Research Hospital, 06500 Ankara, Turkey

Murat Alisik, Ozcan Erel, Department of Biochemistry, Yildirim Beyazıt University Medical Faculty, 06500 Ankara, Turkey

**Author contributions:** Kaplan M and Ates I contributed equally to this work; Kaplan M, Ates I and Yuksel M collected and analyzed the data, and drafted the manuscript; Alisik M and Erel O provided analytical oversight; Ozderin Ozin Y and Kayacetin E designed and supervised the study; Kaplan M, Ates I, Alisik M and Erel O revised the manuscript for important intellectual content; Ozderin Ozin Y and Kayacetin E offered the technical or material support; Kaplan M and Ates I provided administrative support; all authors have read and approved the final version to be published.

**Institutional review board statement:** The study was designed around the 2013 Brazil version of the Declaration of Helsinki. It was approved by the Turkey Yuksek Ihtisas Training and Research Hospital Ethical Board Research Commission.

**Informed consent statement:** Written consent was taken from all participants who were included in the study.

**Conflict-of-interest statement:** The authors declared that there is no conflict of interest.

**Data sharing statement:** Technical appendix, statistical code, and dataset available from the corresponding author at [dr.ihsanates@hotmail.com](mailto:dr.ihsanates@hotmail.com). No additional data are available.

**Open-Access:** This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

[licenses/by-nc/4.0/](http://creativecommons.org/licenses/by-nc/4.0/)

Manuscript source: Invited manuscript

Correspondence to: Ihsan Ates, MD, Department of Internal Medicine, Ankara Numune Training and Research Hospital, 06500 Ankara, Turkey. [dr.ihsanates@hotmail.com](mailto:dr.ihsanates@hotmail.com)  
Telephone: +90-312-5084675  
Fax: +90-312-3569003

Received: November 30, 2016

Peer-review started: December 2, 2016

First decision: January 5, 2017

Revised: January 25, 2017

Accepted: March 12, 2017

Article in press: March 14, 2017

Published online: May 6, 2016

## Abstract

### AIM

To determine dynamic thiol/disulphide homeostasis in celiac disease and to examine the associate with celiac autoantibodies and gluten-free diet.

### METHODS

Seventy three patients with celiac disease and 73 healthy volunteers were enrolled in the study. In both groups, thiol/disulphide homeostasis was examined with a new colorimetric method recently developed by Erel and Neselioglu.

### RESULTS

In patients with celiac disease, native thiol ( $P = 0.027$ ) and total thiol ( $P = 0.031$ ) levels were lower, while disulphide ( $P < 0.001$ ) level, disulphide/native thiol ( $P < 0.001$ ) and disulphide/total thiol ( $P < 0.001$ ) ratios were higher compared to the control group. In patients who do not comply with a gluten-free diet, disulphide/native thiol ratio was found higher compared to the patients who comply with the diet ( $P < 0.001$ ). In patients with



any autoantibody-positive, disulphide/native thiol ratio was observed higher compared to the patients with autoantibody-negative ( $P < 0.05$ ). It is found that there is a negative correlation between celiac autoantibodies, and native thiol, total thiol levels and native thiol/total thiol ratio, while a positive correlation is observed between disulphide, disulphide/native thiol and disulphide/total thiol levels.

### CONCLUSION

This study is first in the literature which found that the patients with celiac disease the dynamic thiol/disulphide balance shifts through disulphide form compared to the control group.

**Key words:** Anti-gliadin antibodies; Anti-tissue transglutaminase antibody; Gluten-free diet; Oxidative stress; Thiol oxidation

© The Author(s) 2017. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core tip:** To the best of our knowledge, for the first time in this study, total and native thiol levels in celiac patients were found lower compared to the control group while disulphide level, disulphide/total thiol and disulphide/native thiol ratios were found to be higher. Also, this study is first in which a negative correlation between celiac autoantibodies and native thiol, and total thiol levels and native thiol/total thiol ratio is observed while there is a positive correlation between disulphide level and disulphide/native thiol and disulphide/total thiol ratios.

Kaplan M, Ates I, Yuksel M, Ozderin Ozin Y, Alisik M, Erel O, Kayacetin E. Thiol/disulphide homeostasis in celiac disease. *World J Gastrointest Pharmacol Ther* 2017; 8(2): 120-126 Available from: URL: <http://www.wjgnet.com/2150-5349/full/v8/i2/120.htm> DOI: <http://dx.doi.org/10.4292/wjgpt.v8.i2.120>

### INTRODUCTION

Celiac disease (CD), observed in genetically predisposed individuals, is a chronic/autoimmune disease of the small intestine, characterized by symptoms such as mucosal damage induced by gliadin, malabsorption, anemia, diarrhea and growth retardation<sup>[1,2]</sup>. While environmental, genetic and immunological factors have a role in etio-pathogenesis of the disease, oxidative stress has also been proved to play a critical role in development of disease<sup>[3,4]</sup>.

With the development of diagnostic methods, gliadin and related toxic effects of prolamines on small intestine are understood better. Gluten peptides in enterocytes, particularly p31-43  $\alpha$ -gliadin peptides, induce certain signal transduction pathways by accumulating in lysosomes and increase the levels of oxidant radicals<sup>[5]</sup>. Based on increased free radicals, a deterioration occurs in the oxidation redox equilibrium<sup>[6]</sup>. At the first stage of

oxidative damage in cellular level based upon free radicals, disulphide (-S-S) linkages are formed by thiol groups (-SH) of amino acids such as sulfur containing cysteine and methionine being oxidized as well as the thiol/disulphide balance collapses in favor of disulphide<sup>[7-12]</sup>. The resulting disulphide bonds are reduced to thiol groups and thiol reserves increase again. Through these reactions at the cellular level, dynamic thiol/disulphide homeostatic status is maintained<sup>[13]</sup>. This dynamic equilibrium is considered to be effective in many cellular processes such as cell death and proliferation, and especially antioxidant balance<sup>[14,15]</sup>. Due to these effects, there are certain studies showing that dynamic equilibrium collapses in cardiovascular diseases and cancer, the oxidative stress of which is particularly evident<sup>[16-18]</sup>.

Dynamic thiol/disulphide homeostasis began to be measured in an easy and repeatable way in 2014 by a new method developed by Erel *et al.*<sup>[19]</sup> with high accuracy and sensitivity. In the literature review, we have not found any study examining dynamic thiol/disulphide homeostatic status in celiac patients using this new method.

In this study, we aimed to measure native thiol, total thiol, disulphide, disulphide/native thiol, disulphide/total thiol and native thiol/total thiol levels in celiac patients with a new and fully automated method of analysis and determine dynamic thiol/disulphide homeostasis.

### MATERIALS AND METHODS

#### Study population

This study was conducted in Turkey Yuksek Ihtisas Training and Research Hospital Gastroenterology Clinic and Ankara Numune Training and Research Hospital Internal Medicine Clinic between January and June 2015.

The study included a total of 146 participants including 73 celiac patients and 73 healthy volunteers. Celiac group is composed of first 73 celiac patients, over the age of 18, who admitted to the polyclinic for routine control. Patients that were diagnosed with CD *via* endoscopic biopsy and subject to regular follow-up in our clinic were included in the patient group in order of their applications. The healthy control group consisted of healthy volunteers, who have applied to our hospital for a check-up, without a chronic disease and drug use and those with similar demographic characteristics to the patient group.

Patients with known diabetes mellitus, kidney failure, malignancy, liver disease, thyroid disease, rheumatic disease, cardiovascular and cerebrovascular disease; smoking, alcohol consumption, vitamin supplements and unfollowed patients were excluded from the study.

In our clinic, the diagnosis of celiac disease in routine is made endoscopically with 2<sup>nd</sup> part duodenal biopsy and anti-gliadin antibody IgA-G or anti-tissue transglutaminase IgA-D positivity. Crypt hyperplasia, villus atrophy and submucosal lymphocytic infiltration is considered significant in the biopsy.

We created two subgroups (GCD: Patients non-

compliant with gluten free diet, GFD: Patients compliant with gluten free diet) to understand the effect of diet in oxidative stress in CD. Poor compliance to diet is defined as taking any kind of gluten containing materials. Patients' compliance to gluten diet is obtained from patient files and applied questionnaires. Patients that are compliant to gluten free diet from the beginning and for at least 5 years are included in GFD group.

Body mass index (BMI) was calculated by dividing body dry weight to the square of tall stature in meters ( $\text{BMI} = \text{kg/m}^2$ ).

The study was conducted in accordance with the Declaration of Helsinki 2013 Brasil version and was approved by the Local Ethics Research Committee. All subjects provided written informed consent prior to participation in the study.

### Biochemical parameters

For thiol/disulphide hemostasis tests, venous blood samples were drawn from patient and control groups after overnight fasting. Blood samples were swiftly centrifuged at 4000 rpm for 10 min, then plasma and serum samples were separated and stored at  $-80^\circ\text{C}$ . Then all parameters were studied in the same session and in the same serum sample.

Laboratory parameters other than thiol/disulphide hemostasis parameters of the participants were their routine parameters at the time they were included in the study and those were recorded from patient files.

### Thiol/disulphide homeostasis

Thiol/disulphide levels were measured by a newly developed, fully-automated and colorimetric method by Erel and Neselioglu<sup>[19]</sup>. When disulphide levels were divided to native thiol and total thiol levels: Disulphide/native thiol and disulphide/total thiol ratios were obtained. When native thiol level was divided to total thiol level, native thiol/total thiol ratio was obtained as a result.

### Statistical analysis

Statistical Package for Social Sciences (SPSS) for Windows 20 (IBM SPSS Inc., Chicago, United States) program was employed for statistical assessments. Kolmogorov-Smirnov test was utilized to determine the distribution of data. Continuous variables with normal distribution were expressed as mean  $\pm$  SD, and continuous variables without normal distribution were expressed as median (min-max). Categorical variables were presented in numbers and percentage. Continuous variables were compared to independent sample *t*-test or Mann-Whitney *U* test where necessary. The relationship between the numeric parameters was analyzed by Pearson and Spearman correlation analysis. In the examination of the relation between thiol/disulphide homeostasis parameters and celiac antibodies, the effects of demographic and clinical factors were adjusted by partial correlation. A  $P < 0.05$  was considered significant for statistical

analyses.

## RESULTS

The demographic characteristics and laboratory findings of all groups are summarized in Table 1. The study population consisted of a total of 146 patients, including 73 celiac patients (female/male: 58/15; age:  $44.1 \pm 13$  years, BMI:  $24.5 \pm 4.7 \text{ kg/m}^2$ ) and 73 controls (female/male: 55/18; age:  $43.7 \pm 13.6$  years, BMI:  $24.9 \pm 4.7 \text{ kg/m}^2$ ). There were no significant difference between two groups in terms of sex, age and BMI levels ( $P > 0.05$ ). The median disease duration of followed celiac patients was determined as 6 years (min: 1 years, max: 25 years).

The mean total protein levels in celiac patients were determined similar to the control group ( $P > 0.05$ ). In celiac patients, mean albumin ( $4.4 \pm 0.3 \text{ g/L}$  vs  $4.1 \pm 0.3 \text{ g/L}$ , respectively;  $P = 0.002$ ), median alanine aminotransferase ( $23 \text{ IU/L}$  vs  $20 \text{ IU/L}$ ,  $P = 0.035$ ), aspartate aminotransferase ( $20 \text{ IU/L}$ , etc.,  $17 \text{ IU/L}$ , respectively;  $P = 0.002$ ) and c-reactive protein ( $3.5 \text{ mg/L}$  vs  $1.2 \text{ mg/L}$ , respectively,  $P = 0.008$ ) levels were higher, compared to the control group.

Mean native thiol ( $322.7 \pm 39.7 \text{ mmol/L}$  vs  $339.6 \pm 51.3 \text{ mmol/L}$ , respectively,  $P = 0.027$ ) and total thiol levels ( $343.7 \pm 41.9 \text{ mmol/L}$  vs  $360.4 \pm 50.3 \text{ mmol/L}$ ,  $P = 0.031$ ) were determined lower in celiac patients, compared to the control group, while native thiol/total thiol ratio did not differ significantly between the groups ( $P > 0.05$ ). Mean disulphide level ( $13.0 \pm 3.7 \text{ mmol/L}$  vs  $10.2 \pm 3.9 \text{ mmol/L}$ , respectively;  $P < 0.001$ ), disulphide/native thiol ( $3.8\% \pm 1.2\%$  vs  $2.8\% \pm 1.1\%$ , respectively,  $P < 0.001$ ) and disulphide/total thiol ratio ( $4.1\% \pm 1.4\%$  vs  $2.9\% \pm 1.1\%$ , respectively,  $P < 0.001$ ) were determined higher in celiac patients compared to the control group.

According to dietary compliance and antibody positivity in the celiac group, the distribution of native thiol, total disulphide and disulphide/native thiol ratios were shown in detail in Table 2. According to this; in patients with dietary compliance, mean native thiol was higher ( $327.9 \pm 35.6 \text{ mmol/L}$  vs  $310.6 \pm 46.8 \text{ mmol/L}$ ,  $P = 0.013$ ), mean disulphide ( $12.0 \pm 3.3 \text{ mmol/L}$  vs  $13.8 \pm 3.2 \text{ mmol/L}$ , respectively;  $P = 0.034$ ) level and disulphide/native thiol ratio ( $3.5\% \pm 1.7\%$  vs  $5.2\% \pm 1.4\%$ , respectively,  $P < 0.001$ ) were much lower compared to patients without dietary compliance. As for antibody positive patients, native thiol level was determined low, disulphide level and disulphide/native thiol ratio were determined higher.

The correlation analysis of native thiol, total thiol, disulphide, disulphide/native thiol, disulphide/total thiol and native thiol/total thiol with demographic and clinical findings is shown in Table 3 in detail. Celiac autoantibodies displayed a negative correlation with native thiol, total thiol and native thiol/total thiol levels and a positive correlation with disulphide, disulphide/native thiol and disulphide/total thiol ratio. The relation

**Table 1** Demographic characteristics and laboratory findings of study population

Variables	Celiac (n = 73)	Control (n = 73)	P value
Gender (male), n (%)	15 (20.5)	18 (24.7)	0.553
Age (yr)	44.1 ± 13	43.7 ± 13.6	0.866
BMI (kg/m <sup>2</sup> )	24.5 ± 4.7	24.9 ± 4.7	0.867
Smoking, n (%)			
Non-smokers	49 (67.1)	50 (68.5)	
Smokers	18 (24.7)	17 (23.3)	0.981
Quit smoking	6 (8.2)	6 (8.2)	
Alcohol, n (%)	2 (2.7)	-	-
Duration of disease (yr)	6 (1-25)	-	-
Total protein (g/L)	7.4 ± 0.6	7.4 ± 0.5	0.964
Albumin (g/L)	4.4 ± 0.3	4.1 ± 0.3	0.002 <sup>a</sup>
ALT (IU/L)	23 (8)	20 (7)	0.035 <sup>a</sup>
AST (IU/L)	20 (12)	17 (9)	0.002 <sup>a</sup>
CRP (mg/L)	3.5 (8.4)	1.2 (2)	0.008 <sup>a</sup>
Native thiol (μmol/L)	322.7 ± 39.7	339.6 ± 51.3	0.027 <sup>a</sup>
Total thiol (μmol/L)	343.7 ± 41.9	360.4 ± 50.3	0.031 <sup>a</sup>
Disulphide (μmol/L)	13.0 ± 3.7	10.2 ± 3.9	< 0.001 <sup>a</sup>
Disulphide/native thiol (%)	3.8 ± 1.2	2.8 ± 1.1	< 0.001 <sup>a</sup>
Disulphide/total thiol (%)	4.1 ± 1.4	2.9 ± 1.1	< 0.001 <sup>a</sup>
Native thiol/total thiol (%)	93.8 ± 2.6	94.3 ± 5.7	0.553

<sup>a</sup>P < 0.05. BMI: Body mass index; ALT: Alanine aminotransferase; AST: Aspartate aminotransferase; CRP: C-reactive protein.

**Table 2** Dietary compliance of native thiol, disulphide and disulphide/native thiol and its distribution regarding antibody positivity

Variables	Native thiol (μmol/L)	P value	Disulphide (μmol/L)	P value	Disulphide/native thiol (μmol/L)	P value
Diet						
GCD (n = 22)	310.6 ± 46.8	0.013 <sup>a</sup>	13.8 ± 3.2	0.034 <sup>a</sup>	5.2 ± 1.4	< 0.001 <sup>a</sup>
GFD (n = 51)	327.9 ± 35.6		12.0 ± 3.3		3.5 ± 1.7	
AGA-IgA						
(-) (n = 33)	310.5 ± 39.5	0.015 <sup>a</sup>	14.1 ± 3.7	0.020 <sup>a</sup>	4.7 ± 1.9	0.005 <sup>a</sup>
(+) (n = 40)	333.8 ± 40.1		12.0 ± 3.8		3.6 ± 1.4	
AGA-IgG						
(-) (n = 47)	308.5 ± 36.8	0.027 <sup>a</sup>	13.9 ± 3.1	0.001 <sup>a</sup>	4.6 ± 1.7	< 0.001 <sup>a</sup>
(+) (n = 26)	330.5 ± 45.2		11.1 ± 3.9		3.0 ± 1.1	
Anti-t TGA						
(-) (n = 29)	315.5 ± 38.2	0.001 <sup>a</sup>	13.9 ± 3.9	0.019 <sup>a</sup>	4.5 ± 1.9	0.007 <sup>a</sup>
(+) (n = 44)	346.2 ± 40.5		11.7 ± 3.8		3.4 ± 1.5	
Anti-t TGG						
(-) (n = 61)	318.1 ± 37.6	0.002 <sup>a</sup>	13.5 ± 3.9	0.009 <sup>a</sup>		0.035 <sup>a</sup>
(+) (n = 12)	345.3 ± 51.5		10.3 ± 3.0			

<sup>a</sup>P < 0.05. GCD: Gluten-containing diet; GFD: Gluten-free diet; AGA-IgA: Anti gliadin antibodies IgA; AGA-IgG: Anti gliadin antibodies IgG; Anti-t TGA: Anti-tissue Transglutaminase IgA antibodies; Anti-t TGG: Anti-tissue Transglutaminase IgA antibodies.

between thiol/disulphide homeostasis parameters and celiac autoantibodies were observed to continue even when the effects of demographic and clinical findings were removed.

C-reactive protein level displayed a negative correlation with native thiol, total thiol and native thiol/total thiol levels, and a positive correlation between disulphide, disulphide/native thiol, disulphide/total thiol levels.

## DISCUSSION

To our knowledge, for the first time in this study, total and native thiol levels in celiac patients were

determined lower compared to the control group while disulphide level, disulphide/total thiol and disulphide/native thiol ratios were found to be higher. Also this is the first study to determine a negative correlation of celiac autoantibodies with native thiol and total thiol levels with native thiol/total thiol ratio, and a positive correlation of disulphide level with disulphide/native thiol and disulphide/total thiol ratios.

Celiac disease is a disease characterized by the inflammatory response created by intestinal mucosa based upon gliadin peptides taken with gluten containing food and consequently mucosal inflammation, crypt hyperplasia and villus atrophy<sup>[20]</sup>. The most important factor in etiopathogenesis of the disease is considered

**Table 3 Findings related to thiol/disulphide hemostasis parameters in celiac patient group**

Variables	Native thiol		Total thiol		Disulphide		Disulphide/total thiol		Disulphide/native thiol		Native thiol/total thiol	
	<i>r</i>	<i>P</i> value	<i>r</i>	<i>P</i> value	<i>r</i>	<i>P</i> value	<i>r</i>	<i>P</i> value	<i>r</i>	<i>P</i> value	<i>r</i>	<i>P</i> value
Age	0.261	0.026 <sup>a</sup>	0.262	0.025 <sup>a</sup>	-0.041	0.729	-0.132	0.264	-0.122	0.302	0.021	0.861
BMI	0.19	0.107	0.187	0.114	-0.033	0.783	-0.096	0.421	-0.113	0.34	0.045	0.702
Total protein	-0.032	0.786	0.005	0.97	0.059	0.618	0.032	0.787	0.086	0.472	-0.156	0.187
Albumin	0.009	0.937	0.025	0.835	-0.037	0.754	-0.065	0.582	-0.040	0.735	-0.071	0.548
AST	0.051	0.666	0.046	0.702	-0.03	0.801	-0.057	0.631	-0.031	0.796	0.033	0.781
ALT	-0.091	0.444	-0.116	0.329	-0.063	0.598	-0.012	0.919	0.002	0.985	0.094	0.427
CRP	-0.327	0.018 <sup>a</sup>	-0.304	0.038 <sup>a</sup>	0.369	0.015 <sup>a</sup>	0.287	0.043 <sup>a</sup>	0.332	0.029 <sup>a</sup>	-0.244	0.038 <sup>a</sup>
AGA-IgA	-0.325	0.009 <sup>a</sup>	-0.266	0.035 <sup>a</sup>	0.384	0.024 <sup>a</sup>	0.324	0.010 <sup>a</sup>	0.325	0.009 <sup>a</sup>	-0.266	0.035 <sup>a</sup>
AGA-IgG	-0.332	0.008 <sup>a</sup>	-0.271	0.022 <sup>a</sup>	0.298	0.010 <sup>a</sup>	0.297	0.011 <sup>a</sup>	0.253	0.031 <sup>a</sup>	-0.279	0.017 <sup>a</sup>
Anti-t TGA	-0.342	0.007 <sup>a</sup>	-0.28	0.017 <sup>a</sup>	0.305	0.009 <sup>a</sup>	0.311	0.004 <sup>a</sup>	0.280	0.035 <sup>a</sup>	-0.294	0.028 <sup>a</sup>
Anti-t TGG	-0.35	0.00 <sup>a</sup>	-0.316	0.004 <sup>a</sup>	0.315	0.007 <sup>a</sup>	0.34	0.001 <sup>a</sup>	0.302	0.014 <sup>a</sup>	-0.304	0.019 <sup>a</sup>
AGA-IgA <sup>1</sup>	-0.313	0.006 <sup>a</sup>	-0.301	0.026 <sup>a</sup>	0.334	0.004 <sup>a</sup>	0.353	< 0.001 <sup>a</sup>	0.315	0.011 <sup>a</sup>	-0.319	0.013 <sup>a</sup>
AGA-IgG <sup>1</sup>	-0.335	0.004 <sup>a</sup>	-0.324	0.005 <sup>a</sup>	0.309	0.018 <sup>a</sup>	0.366	0.023 <sup>a</sup>	0.349	0.033 <sup>a</sup>	-0.288	0.045 <sup>a</sup>
Anti-t TGA <sup>1</sup>	-0.261	0.043 <sup>a</sup>	-0.243	0.035 <sup>a</sup>	0.376	0.036 <sup>a</sup>	0.322	0.011 <sup>a</sup>	0.335	0.009 <sup>a</sup>	-0.273	0.038 <sup>a</sup>
Anti-t TGG <sup>1</sup>	-0.333	0.032 <sup>a</sup>	-0.282	0.037 <sup>a</sup>	0.294	0.031 <sup>a</sup>	0.305	0.026 <sup>a</sup>	0.336	0.036 <sup>a</sup>	-0.314	0.013 <sup>a</sup>

<sup>a</sup>*P* < 0.05. <sup>1</sup>Demographic characteristics and laboratory parameters are adjusted. BMI: Body mass index; ALT: Alanine aminotransferase; AST: Aspartate aminotransferase; CRP: C-reactive protein; AGA-IgA: Anti gliadin antibodies IgA; AGA-IgG: Anti gliadin antibodies IgG; Anti-t TGA: Anti-tissue Transglutaminase IgA antibodies; Anti-t TGG: Anti-tissue Transglutaminase IgA antibodies.

to be environmental factors such as gliadin and the autoimmune response against them. Although the role of oxidative stress in the etiopathogenesis of the disease is unknown, in studies conducted with different cell models, intracellular oxidative imbalance occurs as a result of gliadin exposure and oxidant radicals are formed as a result of lipid peroxidation<sup>[21]</sup>. These oxidant radicals have been shown to form -S-S bonds by oxidizing -SH groups found in the side chain of sulfur containing amino acids and consequently increase oxidized metabolites in the cell<sup>[22]</sup>. In this case thiol/disulphide equilibrium, which is balanced under physiological conditions, is weakened and disrupted in favor of disulphide form. As a result of all these reactions, cell morphology, membrane permeability and vital cellular activities such as apoptosis and cell proliferation are disrupted<sup>[1,5]</sup>.

There are certain studies in the literature investigating thiol and disulphide amounts in low molecular thiol compounds that constitute a small portion of total body thiol pool such as cysteine, glutathione (GSH) and oxidized glutathione thiol<sup>[23,24]</sup>. Until 2014, any colorimetric method that measures total thiol and disulphide amount in the body had not yet been developed. But Erel *et al*<sup>[19]</sup> developed a fully-automated method in 2014, by which total thiol, native thiol and disulphide amounts can be measured easily and repetitively with high sensitivity and specificity.

We have not found any study in the literature examining dynamic thiol/disulphide balance in celiac disease. However, there are studies conducted with low molecular weight thiol compounds. Stojiljković *et al*<sup>[25]</sup> have shown that in intestinal tissues of celiac patients in the pediatric age group, GSH level that constitutes a big part of intracellular thiol content decreases and lipid hydroperoxide level which is an oxidant substance that plays a role in cell membrane damage increases. These results have indicated that GDH redox cycle is disrupted in celiac patients. In a

study conducted with asymptomatic celiac patients by Odetti *et al*<sup>[6]</sup> oxidant radicals derived from protein (carboxyl groups) and lipids (thiobarbituric acid-reactive substances) were determined high.

In our study, total and native thiol levels in celiac patients were determined lower compared to the control group; disulphide level, disulphide/total thiol and disulphide/native thiol ratios were found to be higher and eventually dynamic equilibrium was observed to shift to disulphide form. This case may be due to high levels of oxidant radicals in celiac disease. Our hypothesis is supported by the two studies mentioned above, in which the level of oxidant radicals increase in celiac patients. The increase in the level of oxidant radicals in celiac disease may be due to two cases. Firstly, the disease being a chronic inflammatory disease and secondly, being an autoimmune disease. Previously many studies have indicated that oxidant radicals increase in inflammatory/autoimmune diseases and accordingly oxidative stress level increases<sup>[26-28]</sup>.

For instance, Nanda *et al*<sup>[29]</sup> have determined that in autoantibody positive hypothyroidis patients, levels of oxidant radicals are higher compared to those with autoantibody negative and observed a positive correlation between autoantibodies and oxidant radicals. Determining a positive correlation of C-reactive protein and celiac autoantibodies with disulphide/native thiol level and determining the disulphide form significantly high in autoantibody positive celiac patients strongly supports our thesis.

Another reason for determining low thiol reserve in celiac patients compared to the control group may be due to lack of thiol-containing food intake based upon deteriorated intestinal mucosa. However determining disulphide form and albumin level higher in celiac group compared to the control group indicates that this abnormal



thiol/disulphide equilibrium in celiac patients is not due to lack of oral intake but rather to oxidative stress.

In our study, disulphide/native thiol ratio was determined higher in patients who do not comply with gluten diet compared to those who comply with the diet. We think that this case could be associated with inflammation. Because previous studies have indicated that inflammation is higher in patients who do not comply with gluten diet than those who comply with diet<sup>[30,31]</sup>. Another reason may be due to the significant increase in oxidative stress based upon gliadin toxicity in patients who do not comply with gluten diet<sup>[21]</sup>.

The main limitation of our study is its cross-sectional design and that repetitive measurements have not been done in the patient group. The other limitation is that the information whether participants in the control group have taken thiol containing nutrients or not, is only limited to anamnesis.

For the first time in this study, thiol/disulphide balance was shown to shift towards disulphide form in celiac patients, compared to the control group. Also this is the first study to examine the effects of celiac autoantibodies and gluten-free diet on dynamic thiol/disulphide equilibrium. According to all these results, disrupted thiol/disulphide equilibrium in celiac patients was thought to be associated with autoimmunity and inflammation. In order these results to be clarified, further studies are required to examine the association between thiol/disulphide homeostasis parameters and proinflammatory cytokines that play an active role in celiac disease.

## COMMENTS

### Background

Celiac disease (CD), observed in genetically predisposed individuals, is a chronic/autoimmune disease of the small intestine, characterized by symptoms such as mucosal damage induced by gliadin, malabsorption, anemia, diarrhea and growth retardation.

### Research frontiers

Dynamic thiol/disulphide homeostasis began to be measured in an easy and repeatable way in 2014 by a new method developed by Erel and his colleagues with high accuracy and sensitivity.

### Innovations and breakthroughs

The authors to measure native thiol, total thiol, disulphide, disulphide/native thiol, disulphide/total thiol and native thiol/total thiol levels in celiac patients with a new and fully automated method of analysis and determine dynamic thiol/disulphide homeostasis.

### Applications

Disrupted thiol/disulphide equilibrium in celiac patients was thought to be associated with autoimmunity and inflammation. In order these results to be clarified, further studies are required to examine the association between thiol/disulphide homeostasis parameters and proinflammatory cytokines that play an active role in CD.

### Peer-review

In the present paper, entitled "Thiol/disulphide homeostasis in celiac disease", Kaplan *et al* measured thiol/disulphide homeostasis, an indirect evaluation for oxidative stress, in patients with CD.

## REFERENCES

- 1 Di Sabatino A, Corazza GR. Coeliac disease. *Lancet* 2009; **373**: 1480-1493 [PMID: 19394538 DOI: 10.1016/S0140-6736(09)60254-3]
- 2 Lurz E, Scheidegger U, Spalinger J, Schöni M, Schibli S. Clinical presentation of celiac disease and the diagnostic accuracy of serologic markers in children. *Eur J Pediatr* 2009; **168**: 839-845 [PMID: 18923841 DOI: 10.1007/s00431-008-0845-4]
- 3 Cheeseman KH, Slater TF. An introduction to free radical biochemistry. *Br Med Bull* 1993; **49**: 481-493 [PMID: 8221017 DOI: 10.1093/oxfordjournals.bmb.a072625]
- 4 Diosdado B, van Oort E, Wijmenga C. "Coelionomics": towards understanding the molecular pathology of coeliac disease. *Clin Chem Lab Med* 2005; **43**: 685-695 [PMID: 16207126 DOI: 10.1515/CCLM.2005.117]
- 5 Zimmer KP, Fischer I, Mothes T, Weissen-Plenz G, Schmitz M, Wieser H, Büning J, Lerch MM, Ciclitira PC, Weber P, Naim HY. Endocytotic segregation of gliadin peptide 31-49 in enterocytes. *Gut* 2010; **59**: 300-310 [PMID: 19654123 DOI: 10.1136/gut.2008.169656]
- 6 Odetti P, Valentini S, Aragno I, Garibaldi S, Pronzato MA, Rolandi E, Barreca T. Oxidative stress in subjects affected by celiac disease. *Free Radic Res* 1998; **29**: 17-24 [PMID: 9733018 DOI: 10.1080/10715769800300031]
- 7 Finkel T. Redox-dependent signal transduction. *FEBS Lett* 2000; **476**: 52-54 [PMID: 10878249 DOI: 10.1016/S0014-5793(00)01669-0]
- 8 Dean RT, Fu S, Stocker R, Davies MJ. Biochemistry and pathology of radical-mediated protein oxidation. *Biochem J* 1997; **324** (Pt 1): 1-18 [PMID: 9164834 DOI: 10.1042/bj3240001]
- 9 Ates I, Kaplan M, Inan B, Alisik M, Erel O, Yilmaz N, Guler S. How does thiol/disulfide homeostasis change in prediabetic patients? *Diabetes Res Clin Pract* 2015; **110**: 166-171 [PMID: 26421363 DOI: 10.1016/j.diabres.2015.09.011]
- 10 Ates I, Ozkayar N, Inan B, Yilmaz FM, Topcuoglu C, Neselioglu S, Erel O, Dede F, Yilmaz N. Dynamic thiol/disulphide homeostasis in patients with newly diagnosed primary hypertension. *J Am Soc Hypertens* 2016; **10**: 159-166 [PMID: 26725016 DOI: 10.1016/j.jash.2015.12.008]
- 11 Ates I, Kaplan M, Yuksel M, Mese D, Alisik M, Erel Ö, Yilmaz N, Guler S. Determination of thiol/disulphide homeostasis in type 1 diabetes mellitus and the factors associated with thiol oxidation. *Endocrine* 2016; **51**: 47-51 [PMID: 26547218 DOI: 10.1007/s12020-015-0784-6]
- 12 Ates İ, Ozkayar N, Altay M, Yilmaz FM, Topcuoglu C, Alisik M, Erel Ö, Dede F. Is disulphide/thiol ratio related to blood pressure in masked hypertension? *Clin Exp Hypertens* 2016; **38**: 150-154 [PMID: 26418425 DOI: 10.3109/10641963.2015.1060995]
- 13 Jones DP, Liang Y. Measuring the poise of thiol/disulfide couples in vivo. *Free Radic Biol Med* 2009; **47**: 1329-1338 [PMID: 19715755 DOI: 10.1016/j.freeradbiomed.2009.08.021]
- 14 Powis G, Gasdaska JR, Baker A. Redox signaling and the control of cell growth and death. *Adv Pharmacol* 1997; **38**: 329-359 [PMID: 8895815 DOI: 10.1016/S1054-3589(08)60990-4]
- 15 Jones DP, Maellaro E, Jiang S, Slater AF, Orrenius S. Effects of N-acetyl-L-cysteine on T-cell apoptosis are not mediated by increased cellular glutathione. *Immunol Lett* 1995; **45**: 205-209 [PMID: 7558175 DOI: 10.1016/0165-2478(95)00004-O]
- 16 Matteucci E, Giampietro O. Thiol signalling network with an eye to diabetes. *Molecules* 2010; **15**: 8890-8903 [PMID: 21135801 DOI: 10.3390/molecules15128890]
- 17 Kundi H, Ates I, Kiziltunc E, Cetin M, Cicekcioglu H, Neselioglu S, Erel O, Ornek E. A novel oxidative stress marker in acute myocardial infarction; thiol/disulphide homeostasis. *Am J Emerg Med* 2015; **33**: 1567-1571 [PMID: 26143314 DOI: 10.1016/j.ajem.2015.06.016]
- 18 Prabhu A, Sarcar B, Kahali S, Yuan Z, Johnson JJ, Adam KP, Kensicki E, Chinnaiyan P. Cysteine catabolism: a novel metabolic pathway contributing to glioblastoma growth. *Cancer Res* 2014; **74**: 787-796 [PMID: 24351290 DOI: 10.1158/0008-5472.CAN-13-1423]
- 19 Erel O, Neselioglu S. A novel and automated assay for thiol/disulphide homeostasis. *Clin Biochem* 2014; **47**: 326-332 [PMID: 24351290 DOI: 10.1158/0008-5472.CAN-13-1423]

- 25304913 DOI: 10.1016/j.clinbiochem.2014.09.026]
- 20 **Rostom A**, Dubé C, Cranney A, Saloojee N, Sy R, Garritty C, Sampson M, Zhang L, Yazdi F, Mamaladze V, Pan I, McNeil J, Moher D, Mack D, Patel D. Celiac disease. *Evid Rep Technol Assess (Summ)* 2004; **(104)**: 1-6 [PMID: 15346868]
- 21 **Ellis L**, Dolfini E, Bardella MT. Gliadin cytotoxicity and in vitro cell cultures. *Toxicol Lett* 2003; **146**: 1-8 [PMID: 14615062 DOI: 10.1016/j.toxlet.2003.09.004]
- 22 **Cremers CM**, Jakob U. Oxidant sensing by reversible disulfide bond formation. *J Biol Chem* 2013; **288**: 26489-26496 [PMID: 23861395 DOI: 10.1074/jbc.R113.462929]
- 23 **Boda M**, Nemeth I. Decrease in the antioxidant capacity of red blood cells in children with celiac disease. *Acta paediatrica Hungarica* 1992; **32**: 241-255
- 24 **Grunwell JR**, Gillespie SE, Ward JM, Fitzpatrick AM, Brown LA, Gauthier TW, Hebbard KB. Comparison of Glutathione, Cysteine, and Their Redox Potentials in the Plasma of Critically Ill and Healthy Children. *Front Pediatr* 2015; **3**: 46 [PMID: 26075191 DOI: 10.3389/fped.2015.00046]
- 25 **Stojiljković V**, Todorović A, Pejić S, Kasapović J, Saicić ZS, Radlović N, Pajović SB. Antioxidant status and lipid peroxidation in small intestinal mucosa of children with celiac disease. *Clin Biochem* 2009; **42**: 1431-1437 [PMID: 19560448 DOI: 10.1016/j.clinbiochem.2009.06.009]
- 26 **Burek CL**, Rose NR. Autoimmune thyroiditis and ROS. *Autoimmun Rev* 2008; **7**: 530-537 [PMID: 18625441 DOI: 10.1016/j.autrev.2008.04.006]
- 27 **Kannan S**. Free radical theory of autoimmunity. *Theor Biol Med Model* 2006; **3**: 22 [PMID: 16759382 DOI: 10.1186/1742-4682-3-22]
- 28 **Yuksel M**, Ates I, Kaplan M, Alışık M, Erel Ö, Saygılı F, Kayaçetin E. The dynamic thiol/disulphide homeostasis in inflammatory bowel disease and its relation with disease activity and pathogenesis. *Int J Colorectal Dis* 2016; **31**: 1229-1231 [PMID: 26561415 DOI: 10.1007/s00384-015-2439-8]
- 29 **Nanda N**, Bobby Z, Hamide A. Oxidative stress in anti thyroperoxidase antibody positive hypothyroid patient. *Asian J Biochem* 2012; **7**: 54-58 [DOI: 10.3923/ajb.2012.54.58]
- 30 **Rubio-Tapia A**, Murray JA. Celiac disease. *Curr Opin Gastroenterol* 2010; **26**: 116-122 [PMID: 20040864 DOI: 10.1097/MOG.0b013e3283365263]
- 31 **Ferretti G**, Bacchetti T, Masciangelo S, Saturni L. Celiac disease, inflammation and oxidative damage: a nutrigenetic approach. *Nutrients* 2012; **4**: 243-257 [PMID: 22606367 DOI: 10.3390/nu4040243]

**P- Reviewer:** Ierardi E, Tarocchi M   **S- Editor:** Qi Y   **L- Editor:** A  
**E- Editor:** Lu YJ





Published by **Baishideng Publishing Group Inc**  
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA  
Telephone: +1-925-223-8242  
Fax: +1-925-223-8243  
E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
Help Desk: <http://www.f6publishing.com/helpdesk>  
<http://www.wjgnet.com>

