

31858-Review

Quotes Excluded Bibliography Excluded

8% SIMILAR

FAQ

Name of Journal: World Journal of Hematology

ESPS Manuscript NO: 31858

Manuscript Type: Case Report

Management of unstable angina in a patient with Haemophilia A

Andreina Carbone, Tiziana Formisano, Francesco Natale, Maurizio Cappelli Bigazzi, Donato Tartaglione, Enrica Golia, Felice Gragnano, Mario Crisci, Renato Maria Bianchi, Raffaele Calabrò, Maria Giovanna Russo, Paolo Calabrò

Abstract

Hemophilia A is an X-linked recessive disorder characterized by a deficiency of coagulation factor VIII (FVIII) and therefore by a greater risk of bleeding during percutaneous interventional procedures and during the dual antiplatelet

















The management of unstable angina in a patient with Haemophilia A



2

全部

图片

新闻

视频

购物

更多

设置

工具

找到约 96,800 条结果 (用时 0.80 秒)

Percutaneous Coronary Intervention in a Patient with Acute ST ...

www.invasivecardiology.com/article/8763 ▼ 翻译此页

作者: D Quintero - 相关文章

2008年8月1日 - When available, percutaneous coronary intervention (PCI) is the treatment of choice for acute STEMI. 16 However, patients with hemophilia that

Treating symptomatic coronary artery disease in patients with Von ...

www.sciencedirect.com/science/article/pii/S165838761300071X ▼翻译此页

作者: SA Hassan - 2013 - 被引用次数: 2 - 相关文章

2013年10月1日 - Characteristics of chest pain along with changes on the ECG disease risk factors: prevalence and management in adult hemophilia patients.

[PDF] Successful Management of a Patient with Factor XI Deficiency and ...

synapse.koreamed.org/Synapse/Data/PDFData/0054KCJ/kcj-35-860.pdf ▼ 翻译此页

作者: SH Lee - 被引用次数: 3 - 相关文章

when treating a patient with unstable angina and congenital coagulation disorder during and after percutaneous ... Factor XI deficiency is also called hemophilia.

Consensus Review of the Treatment of Cardiovascular Disease in ...

https://www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: VA Ferraris - 2015 - 被引用次数: 3 - 相关文章

2015年2月6日 - Older patients with milder forms of hemophilia (mild or moderate factor Chest pain syndromes may indicate myocardial ischemia in PWH,

Bleeding risk and the management of bleeding complications in ...

www.bloodjournal.org/content/111/10/4871 ▼ 翻译此页



Management of unstable angina in a patient with Haemophilia A





全部

图片 新闻

耳

视频 购物

更多

设置 工具

找到约 658,000 条结果 (用时 0.55 秒)

Google 学术: Management of unstable angina in a patient with Haemophilia A

... for the management of patients with unstable angina ... - Braunwald - 被引用次数: 777

... events in hypercholesterolaemic patients (JELIS): a ... - Yokoyama - 被引用次数: 1633

... of ischaemic heart disease in haemophilia patients; an ... - Schutgens - 被引用次数: 50

Percutaneous Coronary Intervention in a Patient with Acute ST ...

www.invasivecardiology.com/article/8763 ▼ 翻译此页

作者: D Quintero - 相关文章

2008年8月1日 - When available, percutaneous coronary intervention (PCI) is the treatment of choice for acute STEMI. 16 However, patients with hemophilia that ...

Consensus Review of the Treatment of Cardiovascular Disease in ...

https://www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: VA Ferraris - 2015 - 被引用次数: 5 - 相关文章

2015年2月6日 - Older patients with milder forms of hemophilia (mild or moderate reflects the primary clinical manifestation of IHD [ie, stable angina vs an ...

Percutaneous coronary intervention with stent implantation in ... - NCBI



Management of unstable angina in a patient with Haemophilia A





全部

图片 新闻

闻 视频

껮

更多

设置

工具

找到约 663,000 条结果 (用时 0.65 秒)

Google 学术: Management of unstable angina in a patient with Haemophilia A

购物

.... for the management of patients with unstable angina ... - Braunwald - 被引用次数,784

... events in hypercholesterolaemic patients (JELIS): a ... - Yokoyama - 被引用次数: 1648 ... of ischaemic heart disease in haemophilia patients: an ... - Schutgens - 被引用次数: 50

Percutaneous Coronary Intervention in a Patient with Acute ST ... www.invasivecardiology.com/article/8763 ▼ 翻译此页

www.invasivecardiology.com/article/8/63

作者: D Quintero - 相关文章

2008年8月1日 - When available, percutaneous coronary intervention (PCI) is the treatment of choice for acute STEMI.¹⁶ However, patients with hemophilia that ...

Percutaneous coronary intervention with stent implantation in ... - NCBI https://www.ncbi.nlm.nih.gov/pubmed/17610561 - 翻译此页

... with stent implantation in haemophilic A patient with unstable angina. ... regarding safe interventional or surgical treatment of CAD in haemophiliacs.

Consensus Review of the Treatment of Cardiovascular Disease in ...

https://www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) - 翻译此页作者: VA Ferraris - 2015 - 被引用次数: 5 - 相关文章

2015年2月6日 - Older patients with milder forms of hemophilia (mild or moderate reflects the primary clinical manifestation of IHD [ie, stable angina vs an

Treating symptomatic coronary artery disease in patients with Von ... www.sciencedirect.com/science/article/pii/S165838761300071X ▼ 翻译此页