



33162-Review

 Quotes Excluded
 Bibliography Excluded

 7%
 SIMILAR

 Name of Journal: *World Journal of Nephrology*

Manuscript NO: 33162

Manuscript Type: Original Article

Observational Study

Restless legs syndrome is contributing to fatigue and low quality of life levels in hemodialysis patients

Christoforos D Giannaki, Michael Hadjigavriel, Akis Lazarou, Aristos Michael, Loukas Damianou, Efthimios Atmatzidis, Ioannis Stefanidis, Georgios M Hadjigeorgiou, Giorgos K Sakkas, Marios Pantzaris

Abstract

AIM: Evidence reveals that restless legs syndrome (RLS)/Willis-Ekbom Disease negatively affects various aspects of health and quality of life in patients receiving hemodialysis (HD) therapy. The aim of the current study

Match Overview

1	Crossref 43 words Giannaki, C. D., G. M. Hadjigeorgiou, C. Karatzaferi, M. D. Maridaki, Y. Koutedakis, P. Founta, N. Tsianas, I. Stefanidis	2%
2	Internet 27 words crawled on 18-Apr-2014 aasmnet.org	1%
3	Internet 22 words crawled on 18-Aug-2016 www.elsevier.es	1%
4	Crossref 16 words "Abstracts of TheMovement Disorder Society's Eleventh ... ernational Congress of Parkinson's Disease and Movemen	1%
5	Crossref 15 words Cote, I., D. Trojan, M. Kaminska, M. Cardoso, A. Benedetti, D. Weiss, A. Robinson, A. Bar-Or, Y. Lapierre, and R. Kir...	1%
6	Internet 12 words crawled on 17-Oct-2016 journals.lww.com	1%

[全部](#)[新闻](#)[图片](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 2,460,000 条结果 (用时 1.28 秒)

Fatigue in Patients Receiving Maintenance Dialysis: A Review of ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: M Jhamb - 2008 - 被引用次数: 158 - 相关文章

2008年6月24日 - Keywords: **Fatigue**, end stage **renal** disease, **quality of life**, psychometrics, of **dialysis patients**, **leading** to improved treatment of severe **fatigue**. ... in **patients** with higher versus **lower** target hemoglobin **levels**. The ... **Dialysis patients** have high rates of sleep apnea, insomnia, **restless legs syndrome** and ...

Restless Legs Syndrome in Patients on Hemodialysis: Symptom ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: JM Kim - 2008 - 被引用次数: 46 - 相关文章

2008年12月31日 - Keywords: **restless legs syndrome**, **hemodialysis**, blood urea nitrogen ... indicate incorrect **dialysis** treatment and reduce the **quality of life** in **dialysis patients**. ... Predialysis BUN **level** and **symptom** severity as measured by the IRLSSG ... The **lower** prevalence found in our study might be due to our method of ...

Restless Legs Syndrome: Associated Risk Factors in Hemodialysis ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: SS Beladi-Mousavi - 2015 - 被引用次数: 2 - 相关文章

2015年11月29日 - The mean **level** of serum ferritin was **lower** in **patients** without RLS, but it ... These symptoms usually emerge at night or at times of rest, **leading** to ... to severe RLS, the condition causes **exhaustion**, daytime **fatigue**, and ... **Restless legs syndrome** in hemodialysis patients: health-related

[全部](#)[新闻](#)[图片](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 3,480,000 条结果 (用时 0.74 秒)

Fatigue in Patients Receiving Maintenance Dialysis: A Review of ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: M Jhamb - 2008 - 被引用次数: 175 - [相关文章](#)

2008年6月24日 - Keywords: **Fatigue**, end stage **renal** disease, **quality of life**, psychometrics, of **dialysis patients**, **leading** to improved treatment of severe **fatigue**. ... in **patients** with higher versus **lower** target hemoglobin **levels**. The ... **Dialysis patients** have high rates of sleep apnea, insomnia, **restless legs syndrome** and ...

Restless Legs Syndrome: Associated Risk Factors in Hemodialysis ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: SS Beladi-Mousavi - 2015 - 被引用次数: 3 - [相关文章](#)

2015年11月29日 - The mean **level** of serum ferritin was **lower** in **patients** without **RLS**, ... These symptoms usually emerge at night or at times of rest, **leading** ... In most people with moderate to severe **RLS**, the condition causes **exhaustion**, daytime **fatigue**, ... poor compliance with therapy, impaired health-related **quality of life**, ...

Restless leg syndrome: is it a real problem? - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: PE Cotter - 2006 - 被引用次数: 32 - [相关文章](#)

Restless legs syndrome (RLS) is a common condition that is frequently ... iron deficiency, pregnancy and end-stage **renal** failure and, perhaps, neuropathy. significant effect on the **quality of life** of the sufferer, and the chronic insomnia in A recent study noted that 21% of PD **patients** had RLS, with **lower** ferritin **levels** ...

Restless Legs Syndrome in Patients on Hemodialysis: Symptom ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

Restless legs syndrome is contributing to fatigue and low quality of life levels in he

全部 图片 视频 新闻 地图 图书

找到约 4,020,000 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Fatigue in Patients Receiving Maintenance Dialysis: A Review of ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2582327/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2582327/>

24 Jun 2008 ... Keywords: **Fatigue**, end stage **renal** disease, **quality of life**, psychometrics,
of **dialysis patients**, leading to improved treatment of severe **fatigue**. ... in **patients** with higher
versus **lower** target hemoglobin **levels**. The ... **Dialysis patients** have high rates of sleep apnea,
insomnia, **restless legs syndrome** and ...

[Restless Legs Syndrome in Patients on Hemodialysis: Symptom ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2686851/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2686851/>

31 Dec 2008 ... Keywords: **restless legs syndrome**, **hemodialysis**, blood urea nitrogen ...
indicate incorrect **dialysis** treatment and reduce the **quality of life** in **dialysis patients**. ...
Predialysis BUN **level** and **symptom** severity as measured by the IRLSSG ... The **lower**
prevalence found in our study might be due to our method of ...

[Restless leg syndrome in chronic hemodialysis patients in Mashhad ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5423282/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5423282/>

25 Dec 2016 ... Keywords: **Restless leg syndrome**, **Hemodialysis**, Chronic kidney we found
significant correlation of serum hemoglobin **level** and RLS. ... Erythropoietin dosage was
significantly **lower** in RLS group. ... Authors' **contribution** **Restless legs syndrome** in
hemodialysis patients: health-related **quality of life** ...

[Restless leg syndrome: is it a real problem? - NCBI - NIH](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1936366/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1936366/>

Restless legs syndrome (RLS) is a common condition that is frequently ... iron deficiency,
pregnancy and end-stage **renal** failure and, perhaps, neuropathy. significant effect on the
quality of life of the sufferer, and the chronic insomnia in A recent study noted that 21% of
PD **patients** had RLS, with **lower** ferritin **levels** ...

[\[PDF\] Insomnia and Restless Leg Syndrome in Patients Undergoing ...](#)