

Name of Journal: World Journal of Gastrointestinal Endoscopy

ESPS Manuscript NO: 34022

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Oral esomeprazole versus injectable omeprazole for the prevention of
hemorrhage after ESD

Takashi Uchiyama, Takuma Higurashi, Hitoshi Kuriyama, Yoshinobu Kondo,

Yasuo Hata, Atsushi Nakajima

Abstract

AIM

Match Overview

1	Internet 24 words crawled on 12-Oct-2012 jasn.asnjournals.org	1%
2	Internet 18 words crawled on 24-Jun-2016 connection.ebscohost.com	1%
3	Crossref 17 words Watari, Jiro. "Clinical outcomes and risk factors for perforati on in gastric endoscopic submucosal dissection: A prosp...	1%

学术搜索

找到约 46 条结果 (用时0.07秒)

文章

小提示：只搜索中文(简体)结果，可在 [学术搜索设置](#) 指定搜索语言

我的图书馆

[HTML] **Omeprazole** may be superior to famotidine in the management of iatrogenic ulcer **after** endoscopic mucosal resection: a prospective randomized controlled ...

时间不限

BD Ye, JH Cheon, KD Choi, SG Kim... - Alimentary ..., 2006 - Wiley Online Library

2017以来

... Next article in issue: Sequential therapy using high-dose **esomeprazole**-amoxicillin followed by gatifloxacin for ... Effect of daily **oral omeprazole** on 24 hour intragastric acidity. ... A prospective randomized trial of either famotidine or **omeprazole** for the **prevention of bleeding after** ...

2016以来

被引用次数：48 相关文章 所有 5 个版本 引用 保存

2013以来

自定义范围...

[HTML] A case-control study of **esomeprazole** plus rebamipide vs. **omeprazole** plus rebamipide on post-ESD gastric ulcers

M Bunno, K Gouda, K Yamahara... - Japanese clinical ..., 2013 - ncbi.nlm.nih.gov

按相关性排序

... The group E was administered 20 mg **oral esomeprazole** (Nexium; Astra Zeneca Co, Osaka, Japan) and ... continuous NSAID therapy have shown that, by day 5, once-daily **oral esomeprazole** at doses of ... **Esomeprazole**: a review of its use in the management of gastric acid-related ...

按日期排序

被引用次数：7 相关文章 所有 9 个版本 引用 保存

不限语言

[HTML] Meta-analysis: comparison of **oral** vs. intravenous proton pump inhibitors in patients with peptic ulcer **bleeding**

KKF Tsoi, HW Hirai, JJY Sung - Alimentary pharmacology & ..., 2013 - Wiley Online Library

☒ 包括专利... Intravenous Pantoprazole is much more expensive than **orally** administered forms.[33] The ... Intravenous **esomeprazole** for **prevention** of recurrent peptic ulcer **bleeding**: a randomized trial. ... **Oral** versus intravenous proton pump inhibitors in **preventing** re-bleeding for patients with ...☐ 包含引用

被引用次数：25 相关文章 所有 8 个版本 引用 保存

☒ 创建快讯

[HTML] Two-week treatment with proton pump inhibitor is sufficient for healing post endoscopic submucosal dissection ulcers.

M Arai, T Matsumura, K Okimoto... - World journal of ..., 2014 - europepmc.org

... of **omeprazole** 20 mg every 12 h for 1 d, and received **oral esomeprazole** (20 mg/d ... we treated 45 artificial ulcers **after** ESD only by PPI (**esomeprazole** or **omeprazole** ... endoscopic mucosal resection with circumferential mucosal incision assisted by submucosal **injection** of sodium ...

被引用次数：4 相关文章 所有 8 个版本 引用 保存

Oral esomeprazole vs injectable omeprazole for the prevention of hemorrhage

全部 图片 视频 新闻 地图 图书

找到约 142,000 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[A Case-Control Study of Esomeprazole Plus Rebamipide vs ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3742353/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3742353/>

6 Mar 2013 ... Endoscopic submucosal dissection (ESD) is useful for treating gastric tumors. ... by esomeprazole plus rebamipide (group E) and omeprazole plus The group E was administered 20 mg oral esomeprazole (Nexium; Astra ... There were no cases of delayed gastric perforation or bleeding after discharge.

[Effects of esomeprazole on the healing process of post-endoscopic ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5141375/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5141375/>

1 Dec 2016 ... Objectives: Gastric endoscopic submucosal dissection (ESD) is ... First, 20 mg esomeprazole was orally administered to all subjects before ... are effective in preventing bleeding after the procedure and prompt healing of artificial ulcers. ... Esomeprazole, an S-isomer of omeprazole, is a new form of PPI and ...

[Pantoprazole for the Treatment of Peptic Ulcer Bleeding and ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3987766/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3987766/>

17 Sep 2012 ... Endoscopic therapy significantly reduces rebleeding rates, the need for this contrasts to results for omeprazole and esomeprazole, which have resulted in a ... in healthy volunteers after oral or IV pantoprazole, although volume of for the prevention of bleeding after endoscopic submucosal dissection.

[Effective healing of endoscopic submucosal dissection-induced ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4404649/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4404649/>

15 Apr 2015 ... Keywords: Endoscopic submucosal dissection, ESD-induced ulcers, ... Inclusion criteria included the administration of 20 mg of omeprazole orally for 1 or 8 weeks after ... Japan) with the injection of glycerin and hyaluronic acid solutions. Bleeding from the ulcer was observed in 2 patients in the 8-week ...

[Effect of Intravenous Omeprazole on Recurrent Bleeding after ...](http://www.nejm.org/doi/full/10.1056/NEJM200008033430501)

www.nejm.org/doi/full/10.1056/NEJM200008033430501 ▾

3 Aug 2000 ... After endoscopic treatment of bleeding peptic ulcers, bleeding recurs ... Patients with actively bleeding ulcers or ulcers with nonbleeding ... to receive omeprazole



全部

新闻

图片

视频

购物

更多

设置

工具

找到约 11,800 条结果 (用时 0.61 秒)

A Case-Control Study of Esomeprazole Plus Rebamipide vs ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: M Bunno - 2013 - 被引用次数: 8 - 相关文章

2013年3月6日 - Endoscopic submucosal dissection (ESD) is useful for treating gastric tumors. ... by esomeprazole plus rebamipide (group E) and omeprazole plus The group E was administered 20 mg oral esomeprazole (Nexium; Astra ... There were no cases of delayed gastric perforation or bleeding after discharge.

Effects of esomeprazole on the healing process of post-endoscopic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5141375/> - 翻译此页

作者: T Ichida - 2016 - 相关文章

2016年12月1日 - Objectives: Gastric endoscopic submucosal dissection (ESD) is ... First, 20 mg esomeprazole was orally administered to all subjects before ... are effective in preventing bleeding after the procedure and prompt healing of artificial ulcers. ... Esomeprazole, an S-isomer of omeprazole, is a new form of PPI and ...

Pantoprazole for the Treatment of Peptic Ulcer Bleeding and ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: CJ van Rensburg - 2012 - 被引用次数: 10 - 相关文章

2012年9月17日 - Endoscopic therapy significantly reduces rebleeding rates, the need for this contrasts to results for omeprazole and esomeprazole, which have resulted in a ... in healthy volunteers after oral or IV pantoprazole, although volume of for the prevention of bleeding after endoscopic submucosal dissection.

Effect of Intravenous Omeprazole on Recurrent Bleeding after ...

www.nejm.org/doi/full/10.1056/NEJM200008033430501 - 翻译此页

作者: JYW Lau - 2000 - 被引用次数: 757 - 相关文章