

找到约 5 条结果 (用时 0.62 秒)

Comparison of five tumor regression grading systems for gastric ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5351213/> - 翻译此页

作者: Y Zhu - 2017 - 被引用次数: 5 - 相关文章

2017年3月14日 - Keywords: Gastric cancer, Neoadjuvant chemotherapy, Tumor ... evaluation method predicting patient prognosis is urgently needed in clinical practice. ... gastrectomy in National Cancer Center/Cancer Hospital, Chinese ... while the median number of the examined lymph nodes per case was 24 (3–58).

Significance of histopathological tumor regression after neoadjuvant ...

<https://www.ncbi.nlm.nih.gov/pubmed/21490451> - 翻译此页

作者: K Becker - 2011 - 被引用次数: 142 - 相关文章

(1)Institute of Pathology, Klinikum Rechts der Isar, Technische Universität München, ... BACKGROUND: Histopathological tumor regression may have an important ... resection specimens of GC after neoadjuvant cisplatin-based chemotherapy, using an ... RESULTS: Of the patients in this study, 102 (21.2%) had complete or ...

Significance of Histopathological Tumor Regression After ...

https://www.researchgate.net/.../51048877_Significance_of_Histopathologic... - 翻译此页

Histopathological tumor regression may have an important prognostic impact in ... Tumor Regression After Neoadjuvant Chemotherapy in Gastric Adenocarcinomas A ... Tumor regression (P = 0.009) and postoperative Lymph node status (P A retrospective study of 192 cases from National Cancer Center in China. Article.

Comparison of five tumor regression grading systems for gastric ...

https://www.researchgate.net/.../315060382_Comparison_of_five_tumor_re... - 翻译此页

2018年8月1日 - Methods Totally, 192 patients of gastric adenocarcinoma (including ... chemotherapy: a retrospective study of. 192 ... evaluation to predict outcome is urgent in clinical practice. ... Center/Cancer Hospital, Chinese Academy of Medical Sciences and dian number of the examined lymph nodes per case was.

Curative treatment of gastric cancer - Critical Reviews in Oncology ...

[www.croh-online.com/article/S1040-8428\(02\)00160-9/pdf](http://www.croh-online.com/article/S1040-8428(02)00160-9/pdf) - 翻译此页

作者: AD Roth - 2003 - 被引用次数: 63 - 相关文章

Lymph node dissection extension: conceptual considerations 69. * Tel. Adjuvant systemic chemotherapy trial data analysis Specific problems of neoadjuvant treatment in gastric carcinoma Japanese National Cancer Hospital, mortality rates of change the prognosis of

Name of Journal: *World Journal of Gastrointestinal Oncology*

Manuscript NO: 42426

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Is the evaluation of regression change in lymph nodes necessary for predicting the outcome of gastric adenocarcinoma after neoadjuvant chemotherapy? A retrospective study of 192 cases from the Chinese National Cancer Center

Zhu YL *et al.*

Yue-Lu Zhu, Yong-Kun Sun, Xue-Min Xue, Jiang-Ying Yue, Lin Yang, Li-Yan

Match Overview

1 match

1

Crossref 38 words

Fernández-Aceñero, M. J., M. Granja, J. Sastre, B. García-Paredes, and L. Estrada. "Prognostic significance of tumor ...

1%

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 65,700 条结果 (用时 0.49 秒)

Tumor response evaluation after neoadjuvant chemotherapy in locally ...<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5750190/> - 翻译此页作者 : P Achilli - 2017 - [相关文章](#)

Keywords: Gastric cancer, perioperative chemotherapy, tumor regression ... due to the early lymph node invasion, which is a distinguishing feature of gastric and predicting prognosis after neoadjuvant chemotherapy in patients with locally ...

缺少字词 : unnecessary

Histological evaluation for chemotherapeutic responses of metastatic ...<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4690179/> - 翻译此页作者 : O Kinoshita - 2015 - 被引用次数 : 6 - [相关文章](#)

Keywords: Preoperative chemotherapy, Gastric cancer, Metastatic lymph node, ... An experienced radiologist (Ichijo Y) who was blind to the patients' outcome ... Representative slides of tumor regression grades in metastatic lymph nodes from the regression after neoadjuvant chemotherapy in gastric adenocarcinomas: a ...

缺少字词 : unnecessary

The effects of neoadjuvant chemotherapy on resectability of locally ...<https://www.sciencedirect.com/science/article/pii/S1743919114008334> - 翻译此页作者 : S Hashemzadeh - 2014 - 被引用次数 : 11 - [相关文章](#)

However, the overall prognosis of gastric adenocarcinoma is poor and advanced trial to evaluate the outcome of preoperative chemotherapy on resectability of ... Presence of peri-gastric lymph node involvement in the preoperative CT scans ... neoadjuvant chemotherapy and underwent surgery 4–6 weeks after their last ...

Neoadjuvant chemotherapy for locally advanced gastric cancer: the ...tgc.amegroups.com/article/view/5474/6343 - 翻译此页

The 5-year relative survival for gastric cancer (all stages) is 20% to 25% with a median ... The survival outcome of locally advanced tumors is decreased by high ... role of postoperative chemo-radiotherapy in patients with extended D2 lymphnode regression after neoadjuvant chemotherapy in gastric adenocarcinomas: a ...

Histopathological factors predicting response to neoadjuvant therapy ...https://www.researchgate.net/.../317959599_Histopathological_factors_predi... - 翻译此页

It is important to define factors that can predict response to NAT. ... This study has enrolled 80 patients receiving chemotherapy for locally advanced gastric carcinoma. ... to therapy, 10% had grade

找到约 56,400 条结果 (用时 0.49 秒)

Tumor response evaluation after neoadjuvant chemotherapy in locally ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5750190/> - 翻译此页

作者: P Achilli - 2017 - 相关文章

Keywords: Gastric cancer, perioperative chemotherapy, tumor regression ... due to the early lymph node invasion, which is a distinguishing feature of gastric and predicting prognosis after neoadjuvant chemotherapy in patients with locally ...

缺少字词: unnecessary

Histological evaluation for chemotherapeutic responses of metastatic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4690179/> - 翻译此页

作者: O Kinoshita - 2015 - 被引用次数: 6 - 相关文章

Keywords: Preoperative chemotherapy, Gastric cancer, Metastatic lymph node, ... An experienced radiologist (Ichijo Y) who was blind to the patients' outcome ... Representative slides of tumor regression grades in metastatic lymph nodes from the regression after neoadjuvant chemotherapy in gastric adenocarcinomas: a ...

缺少字词: unnecessary

The effects of neoadjuvant chemotherapy on resectability of locally ...

<https://www.sciencedirect.com/science/article/pii/S1743919114008334> - 翻译此页

作者: S Hashemzadeh - 2014 - 被引用次数: 11 - 相关文章

However, the overall prognosis of gastric adenocarcinoma is poor and advanced trial to evaluate the outcome of preoperative chemotherapy on resectability of ... Presence of peri-gastric lymph node involvement in the preoperative CT scans ... neoadjuvant chemotherapy and underwent surgery 4–6 weeks after their last ...

(PDF) Approaches and genetic determinants in predicting response to ...

https://www.researchgate.net/.../309517155_Approaches_and_genetic_deter... - 翻译此页

2018年8月1日 - neoadjuvant chemotherapy in locally advanced gastric cancer predicting histopathological response after receiving ... specific anticancer therapy is survival and patient outcome. ... residual tumor cells(complete tumor regression) could ... lymph nodes following neoadjuvant radiochemotherapy.

Histopathological factors predicting response to neoadjuvant therapy ...

https://www.researchgate.net/.../317959599_Histopathological_factors_predi... - 翻译此页

It is important to define factors that can predict response to NAT. ... This study has enrolled 80 patients receiving chemotherapy for locally advanced gastric carcinoma. ... to therapy, 10% had grade