

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 1,020,000 条结果 (用时 0.44 秒)

Google 学术 : Insulin resistance is associated to subclinical vascular disease in humans

... the insulin resistance syndrome: the Insulin Resistance ... - Festa - 被引用次数 : 3179

... resistance in patients with clinical and subclinical ... - Maratou - 被引用次数 : 334

Subclinical hypothyroidism is associated with insulin ... - Dessein - 被引用次数 : 129

Relationship between insulin resistance and subclinical ... - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5045581/> - 翻译此页

作者 : H Fakhrazadeh - 2016 - 被引用次数 : 9 - 相关文章

2016年10月1日 - More people in diabetic group than non diabetic group had CACS ≥ 10 than

Insulin resistance-associated cardiovascular disease: potential ...

Subclinical Vascular Dysfunction Associated with Metabolic Syndrome ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4702465/> - 翻译此页

作者 : J Shen - 2015 - 被引用次数 : 8 - 相关文章

We have shown greater subclinical vascular dysfunction in AAs compared with whites, even

Prevalence of Cardiovascular Disease Risk Factors by Race ... hyperglycemia, and insulin resistance), resulting in a proinflammatory and prothrombotic state (1, 2, 23). Nitric oxide activity in the human coronary circulation.

Association between insulin resistance and the development of ...

<https://cardiab.biomedcentral.com/articles/10.1186/s12933-018-0762-4> ▼ 翻译此页

作者 : V Ormazabal - 2018 - 被引用次数 : 5

2018年8月31日 - Insulin resistance and cardiovascular disease. Elevated levels of LDL, smoking, elevated blood pressure and type 1 and type 2 diabetes, are well known risk factors for CVD,

Match Overview



There are no matching sources for this report.

Name of Journal: *World Journal of Diabetes*

Manuscript NO: 45714

Manuscript Type: REVIEW

Insulin resistance is associated to subclinical vascular disease in humans

María M Adeva-Andany, Eva Ameneiros-Rodríguez, Carlos Fernández-Fernández,
Alberto Domínguez-Montero, Raquel Funcasta-Calderón

Abstract

Insulin resistance is associated with subclinical vascular disease that is not justified by conventional cardiovascular risk factors, such as smoking or hypercholesterolemia. Vascular injury associated to insulin resistance involves functional and structural damage to the arterial wall that includes impaired

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 1,020,000 条结果 (用时 0.44 秒)

Google 学术 : Insulin resistance is associated to subclinical vascular disease in humans

... the insulin resistance syndrome: the Insulin Resistance ... - Festa - 被引用次数 : 3179

... resistance in patients with clinical and subclinical ... - Maratou - 被引用次数 : 334

Subclinical hypothyroidism is associated with insulin ... - Dessein - 被引用次数 : 129

Relationship between insulin resistance and subclinical ... - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5045581/> - 翻译此页

作者 : H Fakhrzadeh - 2016 - 被引用次数 : 9 - 相关文章

2016年10月1日 - More people in diabetic group than non diabetic group had CACS ≥ 10 than

Insulin resistance-associated cardiovascular disease: potential ...

Subclinical Vascular Dysfunction Associated with Metabolic Syndrome ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4702465/> - 翻译此页

作者 : J Shen - 2015 - 被引用次数 : 8 - 相关文章

We have shown greater subclinical vascular dysfunction in AAs compared with whites, even

Prevalence of Cardiovascular Disease Risk Factors by Race ... hyperglycemia, and insulin resistance), resulting in a proinflammatory and prothrombotic state (1, 2, 23). Nitric oxide activity in the human coronary circulation.

Association between insulin resistance and the development of ...

<https://cardiab.biomedcentral.com/articles/10.1186/s12933-018-0762-4> ▼ 翻译此页

作者 : V Ormazabal - 2018 - 被引用次数 : 5

2018年8月31日 - Insulin resistance and cardiovascular disease. Elevated levels of LDL, smoking, elevated blood pressure and type 1 and type 2 diabetes, are well known risk factors for CVD.