



All

Images

Videos

翻译成中文

关闭取词

26,200 Results

Any time ▾

## Gastric Neuroendocrine Tumors | Request PDF

[https://www.researchgate.net/publication/8237856\\_Gastric\\_Neuroendocrine\\_Tumors](https://www.researchgate.net/publication/8237856_Gastric_Neuroendocrine_Tumors)

Gastric carcinoids are rare **neuroendocrine tumors**, usually classified as **type I**, if associated with atrophic body gastritis; **type II**, if associated with Zollinger-Ellison syndrome and multiple ...

## Endoscopic diagnosis and management of type I ...

[https://www.researchgate.net/publication/275361304\\_Endoscopic\\_diagnosis\\_and\\_management...](https://www.researchgate.net/publication/275361304_Endoscopic_diagnosis_and_management...)

**Type 1 gastric neuroendocrine tumors** (gNETs) are usually small lesions, restricted to mucosal and sub-mucosal layers of corpus and fundus, with low aggressive behavior, for the majority of cases.

## Review article: the investigation and management of ...

[onlinelibrary.wiley.com/doi/10.1111/apt.12698/full](https://onlinelibrary.wiley.com/doi/10.1111/apt.12698/full)

**Review article: the investigation and management of gastric neuroendocrine tumours ...** David A. Kooby, Charles A. Staley, Shishir K. Maithel, A 15-year experience with **gastric neuroendocrine tumors**: Does **type** make a **difference** ... Sergei Aksenov, Pierre F. Saldinger, Giant **type III** well-differentiated **neuroendocrine tumor** of the **stomach**: A case ...

Published in: **Alimentary Pharmacology & Therapeutics** · 2014

Authors: **Ron Basuroy · Aymn Haji · J K Ramage · A Quaglia · Raj Srirajaskanthan**

Affiliation: **King S College**

About: **Zollinger-Ellison syndrome · Endoscopic submucosal dissection · Proton-pump inhibitor**

## Neuroendocrine Tumors - ScienceDirect

<https://www.sciencedirect.com/science/article/pii/B9780123750976100356>

**Type III gastric NETs** are sporadic **tumors**, and usually well-differentiated. ... NSE is a general **neuroendocrine** marker which cannot **differentiate between** different subtypes of **neuroendocrine tumors**. ... Fasting serum gastrin levels are important to **differentiate type I** and **type II neuroendocrine tumors** of the **stomach** from **type III**.33., 65.

## Gastric cancer: Classification, histology and application ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3418539>

Mar 15, 2012 · Several classifications were proposed in order to address this issue. The scheme endorsed by the International **Gastric Cancer** Association separates **gastric** cancers into **type I**, **type II** and **type III**, to represent the **tumors** at distal esophagus, at cardia and at the **stomach** distal to cardia, respectively. This classification, however, has not ...

Cited by: **227**

Author: **Bing Hu, Nassim El Hajj, Scott Sittler, Na...**

Publish Year: **2012**

**Name of Journal:** *World Journal of Clinical Cases*

**Manuscript NO:** 48438

**Manuscript Type:** MINIREVIEW

**Multifocal G1-G2 gastric neuroendocrine tumors: Differentiating between Type I, II and III, a clinicopathologic review**

Algashaamy K *et al.* Multifocal G1-G2 gastric neuroendocrine tumors

Khaled Algashaamy, Monica Garcia-Buitrago

## Abstract

Gastric neuroendocrine tumors (gNETs) are a rare entity that is increasing in incidence.

## Match Overview

1	<b>Internet</b> 86 words crawled on 25-Jun-2015 <a href="http://www.cap.org">www.cap.org</a>	4%
2	<b>Crossref</b> 37 words Joo Young Kim, Seung-Mo Hong, Jae Y. Ro. "Recent updates on grading and classification of neuroendocrine tumors"	2%
3	<b>Internet</b> 32 words crawled on 03-Jul-2017 <a href="http://gipathol.pathology.or.kr">gipathol.pathology.or.kr</a>	2%
4	<b>Internet</b> 27 words crawled on 21-Jul-2016 <a href="http://docslide.us">docslide.us</a>	1%
5	<b>Crossref</b> 11 words Dermot O'Toole, Gianfranco Delle Fave, Robert T. Jensen ... "Gastric and duodenal neuroendocrine tumours", <i>Best Pract</i>	1%





All

Images

Videos

翻译成中文

关闭取词

27,500 Results

Any time ▾

## Gastric Neuroendocrine Tumours | Request PDF

[https://www.researchgate.net/publication/232280006\\_Gastric\\_Neuroendocrine\\_Tumours](https://www.researchgate.net/publication/232280006_Gastric_Neuroendocrine_Tumours)

Important criteria in **gastric** NEN for the risk evaluation of endoscopic treatment alone are the size of the lesion, depth of invasion and the **tumor** biology, e.g. **neuroendocrine tumor** (NET) G1/G2 ...

## Gastric Neuroendocrine Tumors | Gianfranco Delle Fave ...

[https://www.researchgate.net/publication/8237856\\_Gastric\\_Neuroendocrine\\_Tumors](https://www.researchgate.net/publication/8237856_Gastric_Neuroendocrine_Tumors)

**Gastric** carcinoids are rare **neuroendocrine tumors**, usually classified as **type I**, if associated with atrophic body gastritis; **type II**, if associated with Zollinger-Ellison syndrome and multiple ...

## Gastrointestinal Carcinoid Tumors - National Cancer Institute

<https://www.cancer.gov/types/gi-carcinoid-tumors/hp/gi-carcinoid-treatment-pdq> ▾

Carcinoid **tumors** are rare, slow-growing **tumors** that originate in cells of the diffuse **neuroendocrine** system. They occur most frequently in tissues derived from the embryonic gut. Foregut **tumors**, which account for up to 25% of cases, arise in the lung, thymus, **stomach**, or proximal duodenum. Midgut ...

## Prognostic relevance of a novel TNM classification system ...

<https://onlinelibrary.wiley.com/doi/10.1002/cncr.23549>

**Neuroendocrine tumors** (NETs) of the gastroenteropancreatic system (GEP) are rare neoplasms with an estimated incidence of 1 to 2 per 100,000. 1-3 During the last 30 years, 1 the incidence of GEP NET has increased. 2 Both because of their long clinical course and because of their heterogeneity, GEP NETs provide a substantial challenge to all disciplines involved in the management of these patients.

Cited by: 398

Author: Ulrich-Frank Pape, Ulrich-Frank Pape, He...

Publish Year: 2008

## Neuroendocrine Tumors - ScienceDirect

<https://www.sciencedirect.com/science/article/pii/B9780123750976100356>

**Type III gastric** NETs are sporadic **tumors**, and usually well-differentiated. ... NSE is a general **neuroendocrine** marker which cannot **differentiate between** different subtypes of **neuroendocrine tumors**. ... Fasting serum gastrin levels are important to **differentiate type I** and **type II neuroendocrine tumors** of the **stomach** from **type III**.33., 65.

## Neuroendocrine Carcinomas of the Gastroenteropancreatic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4665594>