

Name of Journal: *World Journal of Hepatology*

Manuscript NO: 56366

Manuscript Type: ORIGINAL ARTICLE

Prospective Study

Subclinical proximal tubulopathy in hepatitis B: the roles of nucleot(s)ide analogue treatment and the hepatitis B virus

Subclinical proximal tubulopathy in hepatitis B

4 Anaïs Brayette, Marie Essig, Paul Carrier, Marilyne Debette-Gratien, Anaïs Labrunie, Sophie Alain, Marianne Maynard, Nathalie Ganne-Carrié, Eric Nguyen-Khac, Pauline Pinet, Victor De Ledinghen, Christophe Renou, Philippe Mathurin, Claire Vanlemmens, Vincent Di Martino, Anne Gervais, Juliette Foucher, Fouchard-Hubert Isabelle, Julien Vergniol, Isabelle Hourmand-Ollivier, Daniel Cohen, Xavier Duval, Thierry Poynard, Marc Bardou, Armand Abergel, Manh-Thong Dao, Thierry Thévenot, Jean-Baptiste Hiriart, Valérie Canva, Guillaume Lassailly, Christine Aurières, Nathalie Boyer,

Match Overview

1	Internet 174 words crawled on 19-May-2020 aasidpubs.onlinelibrary.wiley.com	3%
2	Crossref 90 words "Posters (Abstracts 301–2389)", <i>Hepatology</i> , 2018	2%
3	Crossref 89 words "Abstract Supplement 2019 ACR/ARP Annual Meeting", <i>Arthritis & Rheumatology</i> , 2019	2%
4	Internet 78 words crawled on 08-Feb-2019 hal-univ-bourgogne.archives-ouvertes.fr	1%
5	Crossref 60 words "Posters (Abstracts 289–2348)", <i>Hepatology</i> , 2019	1%
6	Internet 31 words crawled on 20-Jun-2017 www.wjgnet.com	1%

激活 Windows

转到“设置”以激活 Windows

Text-Only Report

Subclinical proximal tubulopathy in Hepatitis B: respective role of treatr



Sign in



ALL

IMAGES

VIDEOS

29,300 Results

Any time ▾

Nucleotide Analogue-Related Proximal Renal Tubular ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5107844>

1. Introduction. Over 350 million people around the world have been diagnosed with chronic **hepatitis B** (CHB) [].Progression of **hepatitis B virus** (HBV) related chronic liver diseases depends on host and viral factors [].Risks of cirrhosis and hepatocellular carcinoma development increase significantly along with increase in baseline HBV viral load [2, 3].

Cited by: 1

Author: Abhasnee Sobhonslidsuk, Jirachaya Wan...

Publish Year: 2016

Hepatitis B: The virus and disease - Liang - 2009 ...

<https://aasldpubs.onlinelibrary.wiley.com/doi/full/10.1002/hep.22881>

Apr 27, 2009 · **Hepatitis B virus** (HBV) infects more than 300 million people worldwide and is a common cause of liver disease and liver cancer. HBV, a member of the Hepadnaviridae family, is a small DNA **virus** with unusual features similar to retroviruses. HBV replicates through an RNA intermediate and can integrate into the host genome.

Search Tools

Turn off Hover Translation (关闭取词)

Subclinical proximal tubulopathy in hepatitis B: the roles of nucleot(



Sign in



ALL

IMAGES

VIDEOS



Add the Give with Bing extension >

168 Results

Any time ▾

(PDF) Impact of Nucleos(t)ide Analogue Combination Therapy ...

<https://www.researchgate.net/publication/275050160...>

However, only a few studies have assessed changes in eGFR in patients who have chronic **hepatitis B** (CHB) and are receiving nucleos(t)ide **analogue** (NA) combination therapy.

Update on prevention, diagnosis, and treatment of chronic ...

<https://aasldpubs.onlinelibrary.wiley.com/doi/10.1002/hep.29800>

Diagnosis

Cause

Epidemiology

Prevention

Resources



The presence of HBsAg establishes the diagnosis of hepatitis B. Chronic versus acute infection is defined by the presence of HBsAg for at least 6 months. The prevalence of HBsAg varies greatly across countries, with high prevalence of HBsAgpositive persons defined as 8%, intermediate as 2% to 7%, and low as <2%.^{21, 22} In developed countries, the prevalence is higher among those who immigrated from high or intermediateprevalence countries and in those with highrisk behaviors.^{22, 23} Some persons may test p...

[See more on aasldpubs.onlinelibrary.wiley.com](https://aasldpubs.onlinelibrary.wiley.com)

Cited by: 890

Author: Norah A. Terrault, Anna S.F. Lok, Brian J. M...

Publish Year: 2018

Search Tools

[Turn on Hover Translation \(开启取词\)](#)

激活 Windows

转到"设置"以激活 Windows

[Feedback](#)

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

355 Results

Any time ▾

[\(PDF\) Impact of Nucleos\(t\)ide Analogue Combination Therapy ...](#)

<https://www.researchgate.net/publication/275050160...>

However, only a few studies have assessed changes in eGFR in patients who have chronic **hepatitis B** (CHB) and are receiving nucleos(t)**ide analogue** (NA) combination therapy.

[Tenofovir-induced Fanconi syndrome in chronic hepatitis B ...](#)

<https://www.researchgate.net/publication/268989628...>

Tubular renal toxicity is a side-effect of long-term therapy with nucleos(t)**ide analogue(s)** (NA) in chronic **hepatitis B** (CHB). There are no established surrogate markers in plasma of early NA ...

[Tenofovir alafenamide versus tenofovir disoproxil fumarate ...](#)

<https://www.sciencedirect.com/science/article/pii/S2468125316300243>

Nov 01, 2016 · Introduction. Nearly 250 million people worldwide are thought to be chronically infected with **hepatitis B virus** (HBV). 1 Chronic **hepatitis B** infection causes progressive fibrosis of the liver, which can lead to cirrhosis, liver decompensation, and hepatocellular carcinoma.2, 3, 4 More than 780 000 people die each year as a result of complications of chronic **hepatitis B**. 5 Clinical guidelines ...

Cited by: 164

Author: Maria Buti, Edward Gane, Wai Kay Seto, ...

Publish Year: 2016

[HIV Glasgow 2018, 28–31 October 2018, Glasgow, UK - 2018 ...](#)

<https://onlinelibrary.wiley.com/doi/full/10.1002/jia2.25187>

No renal discontinuations and no cases of **proximal** renal **tubulopathy** were reported. Conclusions: After 96 weeks, **B/F/TAF** achieved virologic suppression in 84.1% of **treatment**-naïve adults with no **treatment**-emergent resistance, and 100% had HIV-1 RNA <50 c/mL in the per-protocol analysis. **B/F/TAF** was safe and well tolerated with fewer ...

[Posters \(Abstracts 301–2389\) - 2018 - Hepatology - Wiley ...](#)

<https://aasldpubs.onlinelibrary.wiley.com/doi/10.1002/hep.30257>

Background: **Hepatitis B virus** (HBV) infects approximate 240 million people worldwide and 93 million in China . HBV has become one of the leading causes for acute-on-chronic liver failure (ACLF), mainly