

## PEER-REVIEW REPORT

**Name of journal:** World Journal of Hepatology

**Manuscript NO:** 61113

**Title:** Two-stage hepatectomy with radioembolization for bilateral colorectal liver metastases: A case report

**Reviewer's code:** 03120371

**Position:** Peer Reviewer

**Academic degree:** PhD

**Professional title:** Chief Doctor, Professor

**Reviewer's Country/Territory:** China

**Author's Country/Territory:** Italy

**Manuscript submission date:** 2020-11-24

**Reviewer chosen by:** Jin-Lei Wang

**Reviewer accepted review:** 2020-12-29 03:48

**Reviewer performed review:** 2021-01-04 09:20

**Review time:** 6 Days and 5 Hours

<b>Scientific quality</b>	<input type="checkbox"/> Grade A: Excellent <input type="checkbox"/> Grade B: Very good <input checked="" type="checkbox"/> Grade C: Good <input type="checkbox"/> Grade D: Fair <input type="checkbox"/> Grade E: Do not publish
<b>Language quality</b>	<input type="checkbox"/> Grade A: Priority publishing <input checked="" type="checkbox"/> Grade B: Minor language polishing <input type="checkbox"/> Grade C: A great deal of language polishing <input type="checkbox"/> Grade D: Rejection
<b>Conclusion</b>	<input type="checkbox"/> Accept (High priority) <input type="checkbox"/> Accept (General priority) <input checked="" type="checkbox"/> Minor revision <input type="checkbox"/> Major revision <input type="checkbox"/> Rejection
<b>Re-review</b>	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No
<b>Peer-reviewer statements</b>	Peer-Review: <input checked="" type="checkbox"/> Anonymous <input type="checkbox"/> Onymous Conflicts-of-Interest: <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No



**Baishideng  
Publishing  
Group**

7041 Koll Center Parkway, Suite  
160, Pleasanton, CA 94566, USA  
**Telephone:** +1-925-399-1568  
**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
<https://www.wjgnet.com>

#### **SPECIFIC COMMENTS TO AUTHORS**

This is the first report on the efficacy and safety of two-step hepatectomy combined with radio-embolization in the treatment of metastatic colorectal liver metastases. The results showed that the two-step hepatectomy combined with radio-embolization was safe and reliable, which not only guaranteed the regeneration of reserved liver volume, but also could control the progression of tumor and had the value of application. But further clinical studies are needed.