

Name of Journal: *World Journal of Diabetes*

Manuscript NO: 61668

Manuscript Type: REVIEW

Anti- and non-tumor necrosis factor-alpha-targeted therapies effects on insulin resistance in rheumatoid arthritis, psoriatic arthritis and ankylosing spondylitis

targeted-therapies on IR in autoimmune-mediated arthritis

Chrong-Reen Wang, Hung-Wen Tsai

Match Overview

| | | |
|---|--|----|
| 1 | Crossref 60 words Hadi Khodabandehloo, Sattar Gorgani-Firuzjaee, Ghodratollah Panahi, Reza Meshkani. "Molecular and cellular m... | 1% |
| 2 | Internet 45 words crawled on 27-Jul-2020 www.mdpi.com | 1% |
| 3 | Crossref 42 words Chrong-Reen Wang, Ming-Fei Liu. "Recombinant Soluble TNF- α Receptor Fusion Protein Therapy Reduces Insul... | 1% |
| 4 | Internet 42 words crawled on 20-Mar-2019 pt.scribd.com | 1% |
| 5 | Internet 40 words crawled on 26-Aug-2019 academic.oup.com | 1% |
| 6 | Crossref 38 words Hong Ruan, Harvey F Lodish. "Insulin resistance in adip... se tissue: direct and indirect effects of tumor necrosis fact | 1% |

Anti- and non-tumor necrosis factor-alpha-targeted therapies effect

Sign in

ALL IMAGES VIDEOS

12,100 Results Any time

Anti-tumor necrosis factor-α therapy in uveitis ...

https://www.sciencedirect.com/science/article/pii/S0039625715001113

Nov 01, 2015 · Since the first reported use in 2001 of an anti-tumor necrosis factor-alpha (TNF-α) agent, infliximab, for the treatment of uveitis, several new anti-TNF-α agents have emerged for the treatment of refractory noninfectious uveitides, although their use remains off-label in the US.

Cited by: 77 Author: Miguel Cordero-Coma, Lucia Sobrin, Lucia ... Publish Year: 2015

Genome-wide association study of genetic predictors of ...

https://onlinelibrary.wiley.com/doi/10.1002/art.30130

Anti-tumor necrosis factor (anti-TNF) agents are successful therapies in rheumatoid arthritis (RA); however, inadequate response occurs in 30-40% of patients treated. Knowledge of the genetic factors that influence response may facilitate personalized therapy.

Search Tools

Turn off Hover Translation (关闭取词)

PEOPLE ALSO ASK

Are biologics good for arthritis?

Is methotrexate a biologic drug?



12,300 Results

Any time ▾

Non-TNF-Targeted Biologic vs a Second Anti-TNF Drug to ...

<https://www.researchgate.net/publication/309550824...>

Primary lack of response and secondary loss of response (LOR) are major obstacles to the use of **anti-tumor necrosis factor (TNF)-based therapies** in patients with **rheumatoid arthritis** or ...

Abatacept vs Anti-TNF: Lower Diabetes Risk in RA ...

<https://www.medpagetoday.com/rheumatology/arthritis/85970> ▾

Apr 15, 2020 · Patients with **rheumatoid arthritis (RA)** treated with abatacept (Orencia) had a lower risk of developing diabetes mellitus compared with those receiving certain tumor **necrosis factor (TNF ...**

Author: Nancy Walsh

Incidence and complications of interstitial lung disease ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC26555431>

Nov 11, 2015 · Introduction. Interstitial lung disease (ILD) is a common extra-articular condition in **rheumatoid arthritis (RA)**, but few studies have systematically investigated its incidence and risk factors in patients receiving **anti-tumor necrosis factor-alpha (anti-TNF α)** agents or alternate mechanisms of action (MOAs) (e.g., T-cell, B-cell, and interleukin-6 inhibitors).

Cited by: 92

Author: Jeffrey R. Curtis, Khaled Sarsour, Pavel Na...

Search Tools

Turn on Hover Translation (开启取词)

激活 Windows

转到“设置”以激活 Windows。

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

12,100 Results

Any time ▾

Open links in new tab

Anti-tumor necrosis factor- α therapy in uveitis ...

<https://www.sciencedirect.com/science/article/pii/S0039625715001113>

Nov 01, 2015 · The first reported use of **anti**-TNF- α agents comes from the 1980s in animal models of sepsis^{11, 141}; however, subsequent clinical trials in sepsis patients yielded...

Cited by: 77**Author:** Miguel Cordero-Coma, Lucia Sobrin, Lu...**Publish Year:** 2015

(PDF) Effect of anti-tumor necrosis factor- α therapies on ...

<https://www.researchgate.net/publication/5451244...>

Conclusion: Tumor **necrosis** factor α inhibitor treatment appears to be associated with an increase in body weight and BMI, and treatment with **anti**-IL-12/23 and **anti**-IL-17 biologics do not.

PEOPLE ALSO ASK

Are biologics good for arthritis? ▾

Is methotrexate a biologic drug? ▾

What is an example of anti inflammatory medication? ▾

[Feedback](#)

Liver diseases associated with anti-tumor necrosis factor ...

<https://academic.oup.com/ibdjournal/article-abstract/17/1/479/4631203> ▾

Overall, most published reports indicate that **anti**-TNF agents are safe in patients infected with hepatitis C, 19, 26, – 44 as recently reviewed. ¹ However, the safety of **anti**-TNF agents in patient...

Incidence and complications of interstitial lung disease ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/pmid/26555431>

Nov 11, 2015 · Introduction. Interstitial lung disease (ILD) is a common extra-articular condition **in** **rheumatoid arthritis** (RA), but few studies have systematically investigated its incidence and risk...

Cited by: 92**Author:** Jeffrey R. Curtis, Khaled Sarsour, Pavel ...**Publish Year:** 2015