

Microsoft Bing

国内版国际版

Paraesophageal hernia and iron deficiency anemia - mechanisms, di

Chat with Bing

Sign in

Add the Give with Bing extension

ALLIMAGESVIDEOS

60,400 ResultsAny timeOpen links in new tab

Iron-Deficiency Anemia Is a Common Presenting Issue with ...

<https://link.springer.com/article/10.1007/s11605-013-2184-7>

Mar 21, 2013 · A significant percentage of patients with paraesophageal hernia (PEH) will have a co-existing diagnosis of iron-deficiency anemia which will resolve following surgical repair. Between 2000 and 2010, 270 patients underwent operative repair of PEH. Of this group, 123 patients (45.6 %) reported a preexisting diagnosis of iron-deficiency anemia.

Cited by: 2Author: Philip W. Carrott, Sheraz Markar, Jean H...Publish Year: 2012

Effect on iron deficiency anemia of laparoscopic repair of ...

<https://academic.oup.com/dote/article-abstract/18/5/329/2419666>

IntroductionMethodsResultsDiscussion

Some patients with iron deficiency anemia have an associated large paraesophageal hernia and no other

Iron-Deficiency Anemia

Medical condition

Iron deficiency anemia is a common type of anemia - a condition in which blood lacks adequate healthy red blood cells.

MayoClinic

Symptoms

The symptoms of moderate to severe iron deficiency anemia include:

- general fatigue
- weakness
- pale skin

国内版

国际版

Paraesophageal hernia and iron deficiency anemia-mechanisms, dia



ALL

IMAGES

VIDEOS

73,700 Results

Any time ▾

Open links in new tab



### [Iron-Deficiency Anemia Is a Common Presenting Issue with ...](#)

<https://link.springer.com/article/10.1007/s11605-013-2184-7> ▾

Mar 21, 2013 · A significant percentage of patients with paraesophageal hernia (PEH) will have a co-existing diagnosis of iron-deficiency anemia which will resolve following surgical repair. Between 2000 and 2010, 270 patients underwent operative repair of PEH. Of this group, 123 patients (45.6 %) reported a preexisting **diagnosis** of iron-deficiency anemia.

Cited by: 2

Author: Philip W. Carrott, Sheraz Markar, Jean ...

Publish Year: 2012

### [Large hiatal hernia in patients with iron deficiency ...](#)

<https://www.researchgate.net/publication/5265901...>

Iron deficiency anemia in association with a large **paraesophageal hernia** can be treated by laparoscopic repair with acceptable morbidity and minimal mortality. The complications of a large ...

#### PEOPLE ALSO ASK

Does hiatal hernia cause iron deficiency?



Is a hernia repair necessary?



What is laparoscopic Nissen fundoplication?



Feedback

### [\(PDF\) Hiatal Hernia: A Rare Cause of Iron-Deficiency ...](#)

<https://www.researchgate.net/publication/312464644...>

Background: A significant percentage of patients with **paraesophageal hernia** (PEH) will have a co-existing **diagnosis** of **iron-deficiency** anemia which will resolve following surgical repair.

[Iron-Deficiency Anemia Is a Common Presenting Issue with ...](#)



## Iron-Deficiency Anemia Is a Common Presenting Issue with ...

<https://link.springer.com/article/10.1007/s11605-013-2184-7> ▾

Mar 21, 2013 · A significant percentage of patients with paraesophageal hernia (PEH) will have a co-existing diagnosis of iron-deficiency anemia which will resolve following surgical repair. Between 2000 and 2010, 270 patients underwent operative repair of PEH. Of this group, 123 patients (45.6 %) reported a preexisting diagnosis of iron-deficiency anemia.

Cited by: 2

Author: Philip W. Carrott, Sheraz Markar, Jean H...

Publish Year: 2012

## Effect on iron deficiency anemia of laparoscopic repair of ...

<https://academic.oup.com/dote/article-abstract/18/5/329/2419666> ▾

### Introduction

### Methods

### Results

### Discussion

Some patients with iron deficiency anemia have an associated large paraesophageal hernia and no other identifiable source of bleeding.<sup>1–5</sup> Cameron's lesions are areas of linear gastric erosion that have been identified at the neck of the sac of large paraesophageal hernias in less than half of these patients.<sup>5</sup> Several studies have shown that the incidence of paraesophageal hernia in those undergoing investigation for iron deficiency anemia may be in the region of 9%.<sup>3,6</sup> The frequency of anemia in pati...

See more on [academic.oup.com](https://academic.oup.com)

Cited by: 24

Author: J. D. Hayden, G. G. Jamieson

Publish Year: 2005

Published: Oct 01, 2005

### PEOPLE ALSO ASK

Can hernia cause iron deficiency? ▾

What causes iron deficiency anaemia? ▾

Can hiatal hernia cause anaemia? ▾

Feedback

## (PDF) Iron-Deficiency Anemia Is a Common Presenting Issue ...

<https://www.researchgate.net/publication/226067097>