

World Journal of *Gastroenterology*

World J Gastroenterol 2021 May 21; 27(19): 2251-2433



EDITORIAL

- 2251 Celiac Disease in Asia beyond the Middle East and Indian subcontinent: Epidemiological burden and diagnostic barriers

Poddighe D, Abdukhakimova D

FRONTIER

- 2257 Biomarkers in autoimmune pancreatitis and immunoglobulin G4-related disease

Hara A, Watanabe T, Minaga K, Yoshikawa T, Kamata K, Kudo M

REVIEW

- 2270 Receptor for advanced glycation end-products axis and coronavirus disease 2019 in inflammatory bowel diseases: A dangerous liaison?

Rojas A, Schneider I, Lindner C, González I, Morales MA

- 2281 Individualized treatment options for patients with non-cirrhotic and cirrhotic liver disease

Hartl L, Elias J, Prager G, Reiberger T, Unger LW

MINIREVIEWS

- 2299 Clinical characteristics and outcomes of patients with hepatic angiomyolipoma: A literature review

Calame P, Tyrode G, Weil Verhoeven D, Félix S, Klompenhouwer AJ, Di Martino V, Delabrousse E, Thévenot T

- 2312 Risk of hepatitis B virus reactivation in patients with autoimmune diseases undergoing non-tumor necrosis factor-targeted biologics

Akiyama S, Cotter TG, Sakuraba A

- 2325 Burden of venous thromboembolism in patients with pancreatic cancer

Frere C

- 2341 Pathophysiological mechanisms underlying gastrointestinal symptoms in patients with COVID-19

Jin B, Singh R, Ha SE, Zogg H, Park PJ, Ro S

ORIGINAL ARTICLE

Retrospective Cohort Study

- 2353 Risk factors and prognostic value of acute severe lower gastrointestinal bleeding in Crohn's disease

Yoon J, Kim DS, Kim YJ, Lee JW, Hong SW, Hwang HW, Hwang SW, Park SH, Yang DH, Ye BD, Byeon JS, Myung SJ, Yang SK

Retrospective Study

- 2366** Changes in the nutritional status of nine vitamins in patients with esophageal cancer during chemotherapy
Liang LQ, Meng LL, Cai BN, Cui ZP, Ma N, Du LH, Yu W, Qu BL, Feng SQ, Liu F

Observational Study

- 2376** Effects of sepsis and its treatment measures on intestinal flora structure in critical care patients
Yang XJ, Liu D, Ren HY, Zhang XY, Zhang J, Yang XJ
- 2394** Gut microbiota dysbiosis in Chinese children with type 1 diabetes mellitus: An observational study
Liu X, Cheng YW, Shao L, Sun SH, Wu J, Song QH, Zou HS, Ling ZX

META-ANALYSIS

- 2415** Selection of first-line systemic therapies for advanced hepatocellular carcinoma: A network meta-analysis of randomized controlled trials
Han Y, Zhi WH, Xu F, Zhang CB, Huang XQ, Luo JF

ABOUT COVER

Editorial Board Member of *World Journal of Gastroenterology*, Pietro Fusaroli, MD, Associate Professor, Department of Medical and Surgical Sciences, University of Bologna/Hospital of Imola, Via Montericco 4, Imola (BO) 40026, Italy. pietro.fusaroli@unibo.it

AIMS AND SCOPE

The primary aim of *World Journal of Gastroenterology* (WJG, *World J Gastroenterol*) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. WJG mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The WJG is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central, and Scopus. The 2020 edition of Journal Citation Report® cites the 2019 impact factor (IF) for WJG as 3.665; IF without journal self cites: 3.534; 5-year IF: 4.048; Ranking: 35 among 88 journals in gastroenterology and hepatology; and Quartile category: Q2. The WJG's CiteScore for 2019 is 7.1 and Scopus CiteScore rank 2019: Gastroenterology is 17/137.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Yan-Jie Ma*; Production Department Director: *Xiang Li*; Editorial Office Director: *Ze-Mao Gong*.

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Andrzej S Tarnawski, Subrata Ghosh

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

PUBLICATION DATE

May 21, 2021

COPYRIGHT

© 2021 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Risk of hepatitis B virus reactivation in patients with autoimmune diseases undergoing non-tumor necrosis factor-targeted biologics

Shintaro Akiyama, Thomas G Cotter, Atsushi Sakuraba

ORCID number: Shintaro Akiyama 0000-0003-0727-7638; Thomas G Cotter 0000-0002-0376-0107; Atsushi Sakuraba 0000-0003-2519-6129.

Author contributions: Akiyama S performed the literature search, figures, tables, and data collection; all authors conducted study design, data interpretation, and drafting of manuscripts.

Conflict-of-interest statement: The authors have nothing to disclose.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Specialty type: Gastroenterology and hepatology

Country/Territory of origin: United

Shintaro Akiyama, Thomas G Cotter, Atsushi Sakuraba, Section of Gastroenterology, Hepatology and Nutrition, Department of Medicine, University of Chicago, Chicago, IL 60637, United States

Corresponding author: Shintaro Akiyama, MD, MSc, PhD, Academic Fellow, Doctor, Section of Gastroenterology, Hepatology and Nutrition, Department of Medicine, University of Chicago, 5841 S. Maryland Ave. MC 4076, Chicago, IL 60637, United States.
akiyama_s@hotmail.co.jp

Abstract

Hepatitis B virus reactivation (HBVr) can occur in patients treated with immunosuppressive medications. Risk stratification for HBVr based on hepatitis B virus (HBV) serology and viral load is an important strategy to determine appropriate HBV monitoring and antiviral prophylaxis use. Recent advances in the understanding of pathophysiology of autoimmune diseases have led the development of cytokine-targeted therapies. Tumor necrosis factor (TNF)- α inhibitors have been widely used for patients with inflammatory bowel disease, psoriasis, and rheumatic diseases. Further, the clinical benefits of interleukin (IL)-12/23, IL-17, or Janus kinases inhibitors have been demonstrated in these patients. It is well known that TNF- α inhibitor use can lead to HBVr, however, the risk of HBVr in patients undergoing non-TNF-targeted biologics have not been fully understood. In this review, we discuss the risk of HBVr in patients treated with non-TNF-targeted biologics, and immunological mechanisms of these medications causing HBVr.

Key Words: Hepatitis B virus; Autoimmune diseases; Biological therapy; Interleukin-23; Interleukin-17; Janus kinases

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Although the risk of hepatitis B virus reactivation (HBVr) in patients undergoing non-tumor necrosis factor (TNF)-targeted biologics have not been fully understood, some previous studies showed that the risk of HBVr in patients with non-TNF-targeted biologics might be higher than that in patients with TNF- α inhibitors. While patients with chronic hepatitis B virus (HBV) should receive antiviral

States

Peer-review report's scientific quality classification

Grade A (Excellent): A, A, A

Grade B (Very good): 0

Grade C (Good): C

Grade D (Fair): 0

Grade E (Poor): 0

Received: January 24, 2021**Peer-review started:** January 24, 2021**First decision:** February 23, 2021**Revised:** February 27, 2021**Accepted:** April 22, 2021**Article in press:** April 22, 2021**Published online:** May 21, 2021**P-Reviewer:** Georgiev T,
Karagiannakis D, Mohsenzadegan M**S-Editor:** Fan JR**L-Editor:** A**P-Editor:** Wang LL

prophylaxis when they start non-TNF-targeted biologics, antiviral prophylaxis may be a favorable strategy rather than the pre-emptive strategy in patients with resolved HBV. Large-scale studies are needed to ascertain the differential risk of HBVr between patients with TNF- α inhibitors and non-TNF-targeted biologics.

Citation: Akiyama S, Cotter TG, Sakuraba A. Risk of hepatitis B virus reactivation in patients with autoimmune diseases undergoing non-tumor necrosis factor-targeted biologics. *World J Gastroenterol* 2021; 27(19): 2312-2324

URL: <https://www.wjgnet.com/1007-9327/full/v27/i19/2312.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v27.i19.2312>

INTRODUCTION

Hepatitis B virus reactivation (HBVr) can occur in patients treated with immunosuppressive therapy and chemotherapy. In the current era of biologics, physicians need to understand the risk of HBVr in patients with autoimmune diseases undergoing anti-cytokine therapies.

The following three components are important for the development of HBVr: (1) The host immune response; (2) The covalently closed circular DNA of the viral genome of HBV (cccDNA); and (3) The use of immunosuppressive drugs[1]. HBV infection induces a series of innate[2] and adaptive[3] immune responses[1]. The host immune responses against hepatitis B virus (HBV) infection recruit adaptive cytotoxic T (Tc) cells to induce both cytolytic-dependent and -independent antiviral effects. In the cytolytic-independent effect, interferons (IFN) play an important role to suppress the HBV replication. To produce neutralizing antibodies to clear circulating HBV, B cells are also recruited to limit the viral spread of HBV (Figure 1)[4]. However, even when clinical resolution of HBV infection is achieved, it does not mean complete elimination of HBV-DNA because cccDNA can persist in the nucleus of hepatocytes and it can be a source of HBVr when immunosuppressive medications are used.

Tumor necrosis factor (TNF)- α is a key cytokine not only in the pathogenesis of autoimmune diseases but also in the host immune reactions against HBV infection. TNF- α is synthesized by macrophages and T cells and induce the production of a variety of inflammatory cytokines, suppressing viral replication[5]. TNF- α is also necessary for the proliferation of HBV-specific Tc cells that are essential for suppression of HBV replication[6]. Hence, TNF- α inhibitors (*e.g.*, infliximab, adalimumab, and etanercept) can inhibit the anti-HBV immune response, leading to HBV replication[7]. Indeed, the pooled prevalence of HBVr in patients with autoimmune diseases undergoing TNF- α inhibitors was reported to be 4.2% (95%CI: 1.4%-8.2%)[8].

The signaling pathways involving interleukin (IL)-12/23, IL-17, and Janus kinases (JAKs) have been highlighted as novel specific therapeutic targets for autoimmune diseases. A recent multicenter observational study for patients with psoriasis showed that HBVr was significantly more common among patients receiving anti-TNF- α therapies than IL-17 inhibitors[9]. However, there is still limited data in understanding the risk of HBVr in patients who are treated with biologics which inhibit such specific inflammatory pathways. In the present article, we aimed to review previous literatures which assessed the risk of HBVr in patients treated with non-TNF-targeted biologics and discuss how each medication can influence the development of HBVr.

DEFINITIONS OF HBV INFECTION AND REACTIVATION

The professional societies in the United States [American Association for the Study of Liver Diseases (AASLD); American Gastroenterological Association (AGA)], Europe [European Association for the Study of the Liver (EASL)] and Asia [Asian Pacific Association for the Study of the Liver (APASL)] have published guidelines to assist providers with HBVr management[10-13]. In this review article, we divide patients into 2 risk groups which is consistent with the professional society guidelines[10-13] when assessing practical management of HBVr.




Figure 1 The possible immunological mechanism to explain how non-tumor necrosis factor-targeted biologics can induce the development of hepatitis B reactivation. cccDNA: Covalently closed circular DNA; HBV: Hepatitis B virus; IFN: Interferon; IL: Interleukin; T_H17 cells: IL-17 producing T helper cells; T_H1 cells: T helper 1 cells; T_c cells: Cytotoxic T cells.

Chronic HBV

Chronic HBV (CHB) [*i.e.*, Hepatitis B surface antigen (HBsAg)-positive and antibody to hepatitis B core antigen (anti-HBc)-positive] which includes patients with chronic active [serum HBV DNA ≥ 2000 IU/mL and normal or elevated serum alanine transaminase (ALT)] or inactive (serum HBV DNA < 2000 IU/mL and normal ALT) HBV infection.

Resolved HBV

Resolved HBV (*i.e.*, HBsAg-negative and anti-HBc-positive). Of note, there is insufficient evidence to support the use of anti-HBs titers as a decision aid when making a recommendation regarding prophylaxis[11].

There are subtle differences in the definition of HBVr among the professional society guidelines, however, the general concept is the same[10-13]. In patients with CHB, HBVr is defined by a rise in HBV DNA above baseline. In patients with resolved HBV, HBVr is defined by either the appearance of HBV DNA in the blood or conversion to the HBsAg+ state (*i.e.*, seroreversion). The heterogeneity observed in HBVr definition is also reflected in the existing studies on HBVr. The majority of studies included the following parameters: (1) An acute rise in HBV-DNA levels compared with baseline; (2) Elevated levels of serum aminotransferases; and (3) Seroreversion[1]. In this review article, we followed the criteria of HBVr described in each article.

THE MANAGEMENT OF PATIENTS WITH HBV INFECTION UNDERGOING IMMUNOSUPPRESSION

Risk stratification

Patients with CHB have an increased risk of HBVr when undergoing immunosuppressive therapy compared to patients with resolved HBV. For example, among patients treated with TNF- α inhibitors there is an estimated 5-fold increased risk of

HBVr in patients with CHB compared to patients with resolved HBV (15.4% *vs* 3.0% risk of HBVr)[8]. Further, another study showed that the pooled rate of HBVr without antiviral prophylaxis was 15.6% (95%CI: 2.3-35.7) in patients with CHB who were treated with TNF- α inhibitors. In patients with resolved HBV, the pooled rates of HBVr without antiviral prophylaxis in patients who were treated with TNF- α inhibitors and non-TNF-targeted biologics were 1.4% (95%CI: 0.5%-2.6%) and 6.1% (95%CI: 0.0%-16.6%), respectively[14]. Each of the 4 professional societal guidelines recommend testing HBV serology on all candidates for immunosuppressive therapy or chemotherapy to enable appropriate risk stratification (*i.e.* CHB *vs* resolved HBV)[10-13].

Next, the degree of expected iatrogenic immunosuppression should be assessed. Hematopoietic stem cell transplant (HSCT) recipients and B cell-depleting therapies (*e.g.*, rituximab) are high-potency regimens and confer the highest risk of HBVr[10-13]. The AGA guidelines ascertain that anthracyclines (*e.g.*, doxorubicin) and moderate- to high-dose corticosteroids (CS) (*i.e.*, ≥ 10 mg of daily prednisone or equivalent for ≥ 4 wk) confer higher risk than other immunosuppressants[11].

Therapeutic prophylaxis

CHB: In general, the professional societal guidelines recommend HBV prophylaxis, typically entecavir or tenofovir, for all candidates for immunosuppression who have CHB, apart from patients treated with traditional immunosuppressive agents (*e.g.*, thiopurines, methotrexate), intra-articular CSs, or oral CSs ≤ 1 wk[10-13]. The AGA risk stratify this cohort of patients into moderate (1%-10%) and high risk ($> 10\%$) groups for HBVr[11]. Antiviral prophylaxis should be started before and continued after cessation of immunosuppression, generally 12 to 18 mo if high-potency therapies are used and 6 to 12 mo for other therapies[10-13].

Resolved HBV: Guidelines largely agree that resolved HBV patients on high-potency immunosuppression (HSCT recipients and B cell-depleting therapies) should receive HBVr prophylaxis, with the AGA placing this group of patients in the high risk HBVr group ($> 10\%$)[10-13].

For resolved HBV patients not on a high-potency regimen, the guidelines are more dissimilar. AGA recommend prophylaxis for resolved HBV patients at moderate risk (1%-10%) of HBVr, which include patients treated with TNF- α inhibitors, other cytokine or integrin inhibitors, tyrosine kinase inhibitors, moderate- or high-dose CSs for ≥ 4 wk and anthracycline derivatives. In contrast, AASLD, EASL and APASL recommend a pre-emptive therapy for this patient cohort, not prophylaxis, whereby serial lab monitoring (HBV DNA, HBsAg) is performed at 1- to 3-mo intervals on therapy and up to 12 mo after cessation of immunosuppression with on-demand antiviral therapy if needed[10,12,13]. Given that HBsAg seroreversion can lead to fatal acute hepatitis, antiviral therapy should be started immediately, independently of ALT level[11]. Of note, both EASL and APASL recommend treating resolved HBV patients similarly to HBsAg-positive patients if baseline serum HBV-DNA is positive[12,13].

AGA classify resolved HBV patients who are treated with traditional immunosuppressive agents (*e.g.*, thiopurines, methotrexate), low-dose CSs ≥ 4 wk, intra-articular CSs, or any dose of oral CSs for ≤ 1 wk, as low-risk ($< 1\%$) for HBVr and do not recommend prophylaxis, similar to the other society guidelines[10-13].

The risk of HBVr in patients who are treated with non-TNF-targeted biologics

Given the paucity of data on the HBVr risk among patients treated with non-TNF-targeted biologics, we reviewed the existing literature on the risk of HBVr in patients with autoimmune diseases who received non-TNF-targeted biologics and summarized the findings in Tables 1-3. A majority of articles focused on patients with CHB or resolved infection. According to the AGA guideline, CHB and resolved HBV patients treated with non-TNF-targeted therapies are categorized into the moderate-risk HBVr group and therapeutic prophylaxis is recommended. However, AASLD, EASL and APASL recommend serial monitoring among resolved HBV (if HBV DNA is negative) with pre-emptive prophylaxis if HBVr is observed. Therefore, it is important to determine the precise HBVr with non-TNF-targeted biologics to ascertain if a strategy of monitoring/pre-emptive may be too lax, and perhaps a uniform strategy of prophylaxis may be more optimal as recommended by the AGA. Given that patients with resolved infection should be treated similarly to those with CHB patients if their serum HBV-DNA tests are positive at baseline[12,13], we present their baseline HBV-DNA in Tables 1-3.

Table 1 The risk of hepatitis B virus reactivation in patients treated with interleukin-12/23 or interleukin-23 inhibitors

Ref.	Number of HBV patients	HBV status	Disease	Drugs	Prophylaxis	Follow-up	HBV reactivation
Ting <i>et al</i> [27], 2018	54	(1) 10 CHB; and (2) 44 resolved HBV. HBV-DNA at baseline (-)	Psoriasis	Ustekinumab	Yes: 2 patients with CHB	24 mo	(1) 2 patients with CHB without prophylaxis. (no hepatitis); and (2) 1 patient with resolved HBV (mild hepatitis)
Solay <i>et al</i> [5], 2018	29	29 resolved HBV. HBV-DNA at baseline (-)	Psoriasis/HS/AS/RA/CD	Ustekinumab (n = 7)	NA	22 wk	1 patient with psoriasis without prophylaxis (no data regarding hepatitis)
Sanz-Bueno <i>et al</i> [68], 2015	20	20 resolved HBV. HBV-DNA at baseline (-) but viral load was assessed in 7 of 20 patients	Psoriasis	Ustekinumab (n = 6)	No	40 mo	0
Chiu <i>et al</i> [28], 2013	14	(1) 11 CHB; and (2) 3 resolved HBV. HBV-DNA at baseline was not available	Psoriasis	Ustekinumab	Yes: 4 patients with CHB	10 mo	(1) 2 patients with CHB without prophylaxis (No hepatitis); and (2) 0
Navarro <i>et al</i> [69], 2013	5	5 CHB	Psoriasis	Ustekinumab (n = 1)	Yes	25 mo	0
Hayashi <i>et al</i> [70], 2014	5	5 resolved HBV. HBV-DNA at baseline was not available	Psoriasis	Ustekinumab	No	52 wk	0
Koskinas <i>et al</i> [41], 2013	1	Resolved HBV. HBV-DNA at baseline was not available	Psoriasis	Ustekinumab	No	16 mo	1 with hepatitis (ALT 65 IU/mL)
Steglich <i>et al</i> [71], 2014	1	Resolved HBV. HBV-DNA at baseline (-)	Psoriasis	Ustekinumab	Yes	36 mo	0
Duncan <i>et al</i> [43], 2019	1	Resolved HBV. HBV-DNA at baseline was not available	Palmoplantar Psoriasis	Guselkumab	No	12 mo	0

If a study included both of patients with chronic hepatitis B virus (HBV) and those with resolved HBV, we labeled the former with (1) and the latter with (2) in the column with HBV status. HBV: Hepatitis B virus; AS: Ankylosing spondylitis; CD: Crohn's disease; CHB: Chronic HBV; HS: Hidradenitis suppurativa; RA: Rheumatoid arthritis; NA: Not available; ALT: Alanine transaminase.

IL-12/23 INHIBITORS

Mechanism of HBV reactivation

The cytokine IL-12 contributes to the differentiation of naïve T cells to T helper 1 (T_H1) cells and IL-23 maintains and expand IL-17 producing T helper (T_H17) cells (Figure 1) [15]. These two cytokines play a central role to regulate T cell-mediated immune responses, which are dysregulated in various autoimmune diseases including psoriasis and Crohn's disease (CD)[15,16]. The clinical benefit of IL-12 and IL-23 inhibition has been demonstrated in psoriasis, CD, and ulcerative colitis by ustekinumab[17-19], which is an antibody against p40, the common subunit of IL-12 and IL-23. IL-12 plays an important role in achieving sustained control of HBV replication. IL-12 can promote cell-mediated immunity by facilitating the production of IFN-γ production by T_H1 cells, resulting in the inhibition of HBV replication[20,21] and the induction of antiviral effects of HBV-specific Tc cells[22,23]. Indeed, patients with CHB who were treated with recombinant human IL-12 exhibited a high proportion of HBV clearance in a dose-dependent manner[24] and the addition of IL-12 to lamivudine enhanced T cell reactivity to HBV and IFN-γ production[25]. Furthermore, patients with CHB responding to IFN-α treatment were shown to have higher IL-12 and IFN-γ expression levels during the treatment[26]. These findings suggest that ustekinumab might theoretically increase the risk of HBVr.

Table 2 The risk of hepatitis B virus reactivation in patients with interleukin-17 inhibitors

Ref.	Number of HBV patients	HBV status	Disease	Drugs	Prophylaxis	Follow-up	HBV reactivation
Chiu <i>et al</i> [52], 2018	49	(1) 25 CHB; and (2) 24 resolved HBV. HBV-DNA at baseline (-) in 11 patients with resolved HBV	Psoriasis	Secukinumab	Yes: 3 patients with CHB	3 mo	(1) 6 patients with CHB without prophylaxis. (no hepatitis); and (2) 1 patient with resolved HBV with positive viral load at baseline (no hepatitis)
Moneva-Leniz <i>et al</i> [72], 2020	4	(1) 2 CHB; and (2) 2 resolved HBV. HBV-DNA at baseline (-)	Psoriasis/palmoplantar psoriasis	Secukinumab	Yes: 1 patient with CHB and 1 patient with resolved HBV	20 mo	(1) 0; and (2) 0
Feaster <i>et al</i> [73], 2018	1	A carrier of congenital HBV infection ¹	Psoriasis and PsA	Secukinumab	No	24 mo	0
Bevans <i>et al</i> [74], 2018	1	Seropositive hepatitis ¹	Palmoplantar psoriasis and AS	Secukinumab	No	14 mo	0
Yanagihara <i>et al</i> [75], 2017	1	CHB	Psoriasis vulgaris	Secukinumab	Yes	9 mo	0
Peccerillo <i>et al</i> [76], 2018	1	Resolved HBV. HBV-DNA at baseline (-)	Psoriasis	Secukinumab	Yes	14 mo	0
Koike <i>et al</i> [53], 2019	1	CHB	Psoriasis and PsA	Ixekizumab	Yes	18 mo	0
Lora <i>et al</i> [54], 2019	1	Resolved HBV. HBV-DNA at baseline (-)	Psoriasis	Ixekizumab	Yes	12 mo	0

¹A diagnosis in an article was used due to lack of data regarding hepatitis B virus (HBV) serology and viral load. If a study included both of patients with chronic HBV and those with resolved HBV, we labeled the former with (1) and the latter with (2) in the column with HBV status. HBV: Hepatitis B virus; AS: Ankylosing spondylitis; CHB: Chronic HBV; PsA: Psoriatic arthritis.

Table 3 The risk of hepatitis B virus reactivation in patients with Janus kinase inhibitors

Ref.	Number of HBV patients	Serology for HBV infectious	Disease	Drugs	Prophylaxis	Follow-up	HBV reactivation
Chen <i>et al</i> [66], 2018	81	(1) 6 CHB; and (2) 75 resolved HBV. HBV-DNA at baseline (-) but viral load was assessed in 53 patients with resolved HBV	RA	Tofacitinib	Yes: 2 patients with CHB	3-6 mo	(1) 2 patients with CHB without prophylaxis (1 patient developed hepatitis); and (2) 0
Serling-Boyd <i>et al</i> [67], 2021	8	8 resolved HBV. HBV-DNA was assessed in 6 patients, but viral loads were not available	7 RA, 1 PsA	Tofacitinib	Yes: 2 patients	3.1 yr	0
Harigai <i>et al</i> [62], 2020	215	215 resolved HBV. HBV-DNA (-) at baseline in 30 patients with resolved HBV who had detectable post-baseline HBV-DNA	RA	Baricitinib	NA	2.7 yr	8 patients with resolved HBV had HBV-DNA \geq 29 IU/mL (4 patients met the criteria of HBVr in this study, no hepatitis)

If a study included both of patients with chronic hepatitis B virus (HBV) and those with resolved HBV, we labeled the former with (1) and the latter with (2) in the column with HBV status. HBV: Hepatitis B virus; CHB: Chronic HBV; PsA: Psoriatic arthritis; RA: Rheumatoid arthritis; NA: Not available; HBVr: Hepatitis B virus reactivation.

Clinical studies in patients with autoimmune diseases

Several studies have assessed the risk of HBVr in patients treated with ustekinumab (Table 1). A retrospective study showed that no HBVr occurred among 2 patients with CHB taking antiviral prophylaxis after starting ustekinumab, whereas, 2 patients (25%) developed HBVr without hepatitis among 8 patients with CHB without antiviral prophylaxis[27]. Another study demonstrated a 29% rate of HBVr after ustekinumab initiation without antiviral prophylaxis in patients with CHB[28]. Given that patients

with CHB have a high risk of HBVr without antiviral prophylaxis, these patients require antiviral prophylaxis and appropriate monitoring for HBV-DNA and serology tests after initiating ustekinumab treatment.

A retrospective study on 44 patients with resolved HBV who initiated ustekinumab without antiviral prophylaxis found that 1 patient (2.3%) developed HBVr complicated with mild hepatitis[27]. This patient discontinued concurrent methotrexate when reactivation occurred and HBV-DNA became undetectable without antiviral therapy in 6 mo. In another study involving 7 patients with resolved HBV on ustekinumab, 1 patient (14.3%) developed HBVr. This patient was not on antiviral prophylaxis and started entecavir for treatment of HBVr[5]. These data suggest that there is a certain risk of HBVr in patients with resolved HBV even without detectable HBV-DNA at baseline after starting ustekinumab, suggesting that these patients might need antiviral prophylaxis as is the AGA guidelines preferred option. While the guidelines of AASLD, EASL, APASL recommend pre-emptive therapy if HBV DNA is negative, further studies are warranted in order to understand if these patients require antiviral prophylaxis.

IL-23 INHIBITORS

Mechanism of HBV reactivation

IL-23-specific antagonists, such as tildrakizumab[29,30], risankizumab[31,32], guselkumab[33,34], and brazikumab[35], have been shown to be effective for psoriasis and CD. These medications bind to the p19 subunit on IL-23 and inhibit its interaction with the IL-23 receptors[36]. The potential mechanism of HBVr in patients treated with IL-23 inhibitors is still unclear. Previous studies found that T_H17 cells, which are expanded by IL-23 (Figure 1), increase with the severity of liver damage in patients with CHB[37-39]. An observational, clinical-controlled study also demonstrated that the expression levels of IL-23 and IL-17 were associated with increased possibilities of hepatitis B e-antigen (HBeAg) clearance and HBsAg decline in patients with HBeAg-positive CHB during pegylated IFN therapy[40]. This study also found that high serum IL-23 Levels can predict the response to IFN therapy in patients with HBeAg-positive CHB[40]. Given that T_H17 cells promote the differentiation and function of B cells[41,42], IL-23 might activate the humoral immune response against circulating HBV and play a role to facilitate HBV clearance by IFN therapy (Figure 1). Although this hypothesis suggest that IL-23 inhibitors may abrogate the HBV clearance, it still remains to be elucidated whether these medications contribute to the development of HBVr.

Clinical studies in patients with autoimmune diseases

The data regarding the safety of IL-23 inhibitors in patients with HBV infection is limited. A case report showed that a patient with resolved HBV infection did not develop HBVr 1 year after starting guselkumab (Table 1)[43]. There have been no reported studies focusing on the risk of HBVr in patients treated with other IL-23 inhibitors.

IL-17 INHIBITORS

Mechanism of HBV reactivation

IL-17 is a major effector cytokine of T_H17 cells and mediate host defense mechanisms[44]. Inhibition of IL-17 with secukinumab, ixekizumab, and brodalumab have demonstrated clinical benefits in patients with psoriasis[45-47], psoriatic arthritis[48], and ankylosing spondylitis[49]. T_H17/IL-17 axis is involved in the process of fibrogenesis and increases the expression of proinflammatory cytokines, promoting the recruitment of inflammatory cells in patients with CHB[50]. A previous study showed that Th17 cells were significantly increased in patients with CHB, as well as the expression level of IL-17[51]. They also demonstrated that the suppression of viral replication induced by IFN- α resulted in a decrease in T_H17 cells and IL-17 expression, suggesting that T_H17 cells might play an important role during IFN- α treatment to eliminate HBV[51]. As we described above, T_H17 cells also facilitate B cells[41,42] and would enhance the humoral response to clear circulating HBV (Figure 1). These findings implicate that T_H17/IL-17 axis might be associated with HBV clearance and its inhibition may increase the risk of HBVr.

Clinical studies in patients with autoimmune diseases

A prospective multicenter study on 22 patients with CHB with no antiviral prophylaxis after starting secukinumab showed that 6 patients (27.3%) developed HBVr (Table 2)[52]. Three patients with HBVr started antiviral treatments and their viral loads decreased rapidly within 3 mo. The remaining three patients with HBVr were followed without antiviral drugs and their viral loads remained low without acute hepatitis. Notably, none of the 3 patients with CHB who received antiviral prophylaxis developed HBVr[52]. Hence, this study reinforced the importance of antiviral prophylaxis in patients with CHB starting treatment with IL-17 inhibitors. This study also included 24 patients with resolved HBV who did not receive antiviral prophylaxis and identified one patient (4.2%) with a positive viral load at baseline who developed HBVr without acute hepatitis[52]. This study re-affirmed the EASL and APASL guidelines which recommend antiviral prophylaxis in patients with resolved HBV if their baseline viral loads are positive.

A case report on a patient with CHB treated with ixekizumab and entecavir simultaneously did not develop HBVr after 18 mo of treatment[53]. Another report showed that a patient with resolved HBV did not experience HBVr during follow-up (Table 2)[54]. Given that data regarding the risk of HBVr in patients treated with ixekizumab or brodalumab are still limited, further studies with larger sample sizes are warranted.

JAK INHIBITORS

Mechanism of HBV reactivation

JAKs bind to type I and II cytokine receptors and transmit extracellular cytokine signals to activate various signal transducers and activators of transcription, which drive the proinflammatory machinery of the cellular immune response[55]. The clinical benefit of JAK inhibitors has been demonstrated in patients with rheumatoid arthritis[56,57], psoriatic arthritis[58,59], and ulcerative colitis[60,61]. Important signaling pathways in host-defense include innate antiviral responses *via* IFN- α/β mediated by JAK1-tyrosine kinase 2 complexes, and IFN- γ mediated by JAK1-JAK2 complexes[55]. Hence, JAK inhibitors might counteract the suppressive effects of IFN on viral replication[62,63]. Further, dendritic cells and effective T cell lineages including T_H cells and Tc cells play important roles to defense against HBV-infection (Figure 1)[64]. A previous study demonstrated that a JAK inhibitor can block the differentiation and function of dendritic cells, leading to impaired T cell activation (Figure 1)[65]. Thus, it was suggested that JAK inhibitors might negatively interact with the defense mechanism against HBV infection. Further studies investigating how JAK inhibitors influence the development of HBVr are warranted.

Clinical studies in patients with autoimmune diseases

A retrospective cohort study including 6 patients with CHB showed that 2 out of 4 patients (50%) without antiviral prophylaxis developed HBVr after starting tofacitinib. One patient had an elevated ALT level and started entecavir, resulting in declines in HBV-DNA and ALT levels. Another patient started entecavir and did not develop acute hepatitis. Both patients continued tofacitinib after the development of HBVr. Meanwhile, 2 patients with CHB who received antiviral prophylaxis did not develop HBVr after initiating tofacitinib. Further, in this study, none of 75 patients with resolved HBV received antiviral prophylaxis and no HBVr was observed in this group[66]. Another study also demonstrated that patients with resolved HBV did not develop HBVr after starting tofacitinib (Table 3)[67].

A study assessing data which was integrated from four phase 3 trials of baricitinib in patients showed that, among 215 patients with resolved HBV, 8 patients (3.7%) had a single quantifiable result of HBV-DNA viral load (HBV-DNA level ≥ 29 IU/mL) after initiating baricitinib. Among these 8 patients, 4 patients met the definition of HBVr (HBV-DNA ≥ 100 IU/mL), but no patients developed hepatitis. HBV-DNA at baseline was assessed in 6 patients and all examined patients did not have detectable HBV-DNA level. Antiviral therapy was not used in 5 of 8 patients[62].

All these findings suggest that patients with CHB should receive antiviral prophylaxis when they start JAK inhibitors. As for patients with resolved HBV infection, given that HBVr was occasionally reported even if their HBV-DNA levels were not detected at baseline, an appropriate consultation with hepatologists is necessary. There has been limited data regarding the risk of HBVr in patients with autoimmune diseases who are treated with other JAK inhibitors (*e.g.*, upadacitinib,

filgotinib, peficitinib).

CONCLUSION

In summary, considering antiviral prophylaxis with an appropriate risk stratification is necessary when we start non-TNF-targeted biologics for patients with autoimmune diseases. The frequencies of HBVr without antiviral prophylaxis in patients with CHB on IL-12/23, IL-17, and JAK inhibitors are up to 29%, 27%, and 50%, respectively. A meta-analysis demonstrated that the pooled rate of HBVr without antiviral prophylaxis was 15.6% (95%CI: 2.3-35.7) in patients with CHB who were treated with TNF- α inhibitors, suggesting that non-TNF-targeted biologics, particularly JAK inhibitors, may have a higher risk of HBVr compared with TNF- α inhibitors. Given that no patients who received antiviral prophylaxis developed HBVr, HBVr is preventable with antiviral therapy in patients with CHB on non-TNF-targeted biologics. As all of professional societies recommended in their guidelines, patients with CHB should receive antiviral prophylaxis when they start non-TNF-targeted biologics. In patients with resolved HBV, the rates of HBVr without antiviral prophylaxis in patients on IL-12/23, IL-17, and JAK inhibitors are up to 2.3%, 4.2%, and 0%, respectively. The meta-analysis showed that the pooled rates of HBVr without antiviral prophylaxis in patients who were treated with TNF- α inhibitors and non-TNF-targeted biologics were 1.4% (95%CI: 0.5%-2.6%) and 6.1% (95%CI: 0.0%-16.6%), respectively[14]. These data supported that the risk of HBVr in patients treated with non-TNF-targeted biologics might be higher than that in patients with TNF- α inhibitors even if their HBV status is resolved HBV. According to the AGA guideline, patients with resolved HBV who are treated with non-TNF-targeted biologics are categorized into the moderate risk group and antiviral prophylaxis are recommended for this patient cohort[11]. However, as stated previously, AASLD, EASL and APASL recommend the pre-emptive therapeutic strategy for this cohort, although APASL and EASL do include the caveat of potentially using HBV DNA assessment to aid decision-making[10,12,13]. Given the higher risk of HBVr with non-TNF-targeted biologics compared with TNF- α inhibitors, antiviral prophylaxis may be a favorable strategy rather than the pre-emptive strategy to prevent HBVr in patients with resolved HBV. Large-scale studies are needed to ascertain the differential risk of HBVr between patients with TNF- α inhibitors and non-TNF-targeted biologics and to stratify the risk of HBVr by the type of non-TNF-targeted biologics. While HBsAg seroreversion can lead to fatal acute hepatitis, a consultation with hepatologists or infectious disease specialists is recommended.

REFERENCES

- 1 Gentile G, Antonelli G. HBV Reactivation in Patients Undergoing Hematopoietic Stem Cell Transplantation: A Narrative Review. *Viruses* 2019; **11** [PMID: 31717647 DOI: 10.3390/v11111049]
- 2 Maini MK, Gehring AJ. The role of innate immunity in the immunopathology and treatment of HBV infection. *J Hepatol* 2016; **64**: S60-S70 [PMID: 27084038 DOI: 10.1016/j.jhep.2016.01.028]
- 3 Bertolotti A, Ferrari C. Adaptive immunity in HBV infection. *J Hepatol* 2016; **64**: S71-S83 [PMID: 27084039 DOI: 10.1016/j.jhep.2016.01.026]
- 4 Guidotti LG, Chisari FV. Immunobiology and pathogenesis of viral hepatitis. *Annu Rev Pathol* 2006; **1**: 23-61 [PMID: 18039107 DOI: 10.1146/annurev.pathol.1.110304.100230]
- 5 Solay AH, Acar A, Eser F, Kuşcu F, Tütüncü EE, Kul G, Şentürk GÇ, Gürbüz Y. Reactivation rates in patients using biological agents, with resolved HBV infection or isolated anti-HBc IgG positivity. *Turk J Gastroenterol* 2018; **29**: 561-565 [PMID: 30260778 DOI: 10.5152/tjg.2018.18032]
- 6 Kasahara S, Ando K, Saito K, Sekikawa K, Ito H, Ishikawa T, Ohnishi H, Seishima M, Kakumu S, Moriwaki H. Lack of tumor necrosis factor alpha induces impaired proliferation of hepatitis B virus-specific cytotoxic T lymphocytes. *J Virol* 2003; **77**: 2469-2476 [PMID: 12551985 DOI: 10.1128/jvi.77.4.2469-2476.2003]
- 7 Manzano-Alonso ML, Castellano-Tortajada G. Reactivation of hepatitis B virus infection after cytotoxic chemotherapy or immunosuppressive therapy. *World J Gastroenterol* 2011; **17**: 1531-1537 [PMID: 21472116 DOI: 10.3748/wjg.v17.i12.1531]
- 8 Cantini F, Boccia S, Goletti D, Iannone F, Leoncini E, Panic N, Prignano F, Gaeta GB. HBV Reactivation in Patients Treated with Antitumor Necrosis Factor-Alpha (TNF- α) Agents for Rheumatic and Dermatologic Conditions: A Systematic Review and Meta-Analysis. *Int J Rheumatol* 2014; **2014**: 926836 [PMID: 25114684 DOI: 10.1155/2014/926836]
- 9 Chiu HY, Chiu YM, Chang Liao NF, Chi CC, Tsai TF, Hsieh CY, Hsieh TY, Lai KL, Chiu TM, Wu NL, Hui RC, Lee CN, Wang TS, Chen PH, Yang CC, Huang YH. Predictors of hepatitis B and C

- virus reactivation in patients with psoriasis treated with biological agent: A nine-year multicenter cohort study. *J Am Acad Dermatol* 2019 [PMID: [31821860](#) DOI: [10.1016/j.jaad.2019.12.001](#)]
- 10 **Terrault NA**, Lok ASF, McMahon BJ, Chang KM, Hwang JP, Jonas MM, Brown RS Jr, Bzowej NH, Wong JB. Update on prevention, diagnosis, and treatment of chronic hepatitis B: AASLD 2018 hepatitis B guidance. *Hepatology* 2018; **67**: 1560-1599 [PMID: [29405329](#) DOI: [10.1002/hep.29800](#)]
 - 11 **Reddy KR**, Beavers KL, Hammond SP, Lim JK, Falck-Ytter YT; American Gastroenterological Association Institute. American Gastroenterological Association Institute guideline on the prevention and treatment of hepatitis B virus reactivation during immunosuppressive drug therapy. *Gastroenterology* 2015; **148**: 215-9; quiz e16 [PMID: [25447850](#) DOI: [10.1053/j.gastro.2014.10.039](#)]
 - 12 **European Association for the Study of the Liver**. EASL 2017 Clinical Practice Guidelines on the management of hepatitis B virus infection. *J Hepatol* 2017; **67**: 370-398 [PMID: [28427875](#) DOI: [10.1016/j.jhep.2017.03.021](#)]
 - 13 **Sarin SK**, Kumar M, Lau GK, Abbas Z, Chan HL, Chen CJ, Chen DS, Chen HL, Chen PJ, Chien RN, Dokmeci AK, Gane E, Hou JL, Jafri W, Jia J, Kim JH, Lai CL, Lee HC, Lim SG, Liu CJ, Locarnini S, Al Mahtab M, Mohamed R, Omata M, Park J, Piratvisuth T, Sharma BC, Sollano J, Wang FS, Wei L, Yuen MF, Zheng SS, Kao JH. Asian-Pacific clinical practice guidelines on the management of hepatitis B: a 2015 update. *Hepatol Int* 2016; **10**: 1-98 [PMID: [26563120](#) DOI: [10.1007/s12072-015-9675-4](#)]
 - 14 **Lin TC**, Yoshida K, Tedeschi SK, de Abreu MM, Hashemi N, Solomon DH. Risk of Hepatitis B Virus Reactivation in Patients With Inflammatory Arthritis Receiving Disease-Modifying Antirheumatic Drugs: A Systematic Review and Meta-Analysis. *Arthritis Care Res (Hoboken)* 2018; **70**: 724-731 [PMID: [28834412](#) DOI: [10.1002/acr.23346](#)]
 - 15 **Yen D**, Cheung J, Scheerens H, Poulet F, McClanahan T, McKenzie B, Kleinschek MA, Owyang A, Mattson J, Blumenschein W, Murphy E, Sathe M, Cua DJ, Kastelein RA, Rennick D. IL-23 is essential for T cell-mediated colitis and promotes inflammation via IL-17 and IL-6. *J Clin Invest* 2006; **116**: 1310-1316 [PMID: [16670770](#) DOI: [10.1172/JCI21404](#)]
 - 16 **Lee E**, Trepicchio WL, Oestreicher JL, Pittman D, Wang F, Chamian F, Dhodapkar M, Krueger JG. Increased expression of interleukin 23 p19 and p40 in lesional skin of patients with psoriasis vulgaris. *J Exp Med* 2004; **199**: 125-130 [PMID: [14707118](#) DOI: [10.1084/jem.20030451](#)]
 - 17 **Papp KA**, Langley RG, Lebwohl M, Krueger GG, Szapary P, Yelding N, Guzzo C, Hsu MC, Wang Y, Li S, Dooley LT, Reich K; PHOENIX 2 study investigators. Efficacy and safety of ustekinumab, a human interleukin-12/23 monoclonal antibody, in patients with psoriasis: 52-week results from a randomised, double-blind, placebo-controlled trial (PHOENIX 2). *Lancet* 2008; **371**: 1675-1684 [PMID: [18486740](#) DOI: [10.1016/S0140-6736\(08\)60726-6](#)]
 - 18 **Sandborn WJ**, Gasink C, Gao LL, Blank MA, Johans J, Guzzo C, Sands BE, Hanauer SB, Targan S, Rutgeerts P, Ghosh S, de Villiers WJ, Panaccione R, Greenberg G, Schreiber S, Lichtiger S, Feagan BG; CERTIFI Study Group. Ustekinumab induction and maintenance therapy in refractory Crohn's disease. *N Engl J Med* 2012; **367**: 1519-1528 [PMID: [23075178](#) DOI: [10.1056/NEJMoa1203572](#)]
 - 19 **Sands BE**, Sandborn WJ, Panaccione R, O'Brien CD, Zhang H, Johans J, Adedokun OJ, Li K, Peyrin-Biroulet L, Van Assche G, Danese S, Targan S, Abreu MT, Hisamatsu T, Szapary P, Marano C; UNIFI Study Group. Ustekinumab as Induction and Maintenance Therapy for Ulcerative Colitis. *N Engl J Med* 2019; **381**: 1201-1214 [PMID: [31553833](#) DOI: [10.1056/NEJMoa1900750](#)]
 - 20 **Zeuzem S**, Carreño V. Interleukin-12 in the treatment of chronic hepatitis B and C. *Antiviral Res* 2001; **52**: 181-188 [PMID: [11672828](#) DOI: [10.1016/S0166-3542\(01\)00183-8](#)]
 - 21 **Cavanaugh VJ**, Guidotti LG, Chisari FV. Interleukin-12 inhibits hepatitis B virus replication in transgenic mice. *J Virol* 1997; **71**: 3236-3243 [PMID: [9060687](#) DOI: [10.1128/JVI.71.4.3236-3243.1997](#)]
 - 22 **Guidotti LG**. The role of cytotoxic T cells and cytokines in the control of hepatitis B virus infection. *Vaccine* 2002; **20** Suppl 4: A80-A82 [PMID: [12477433](#) DOI: [10.1016/S0264-410X\(02\)00392-4](#)]
 - 23 **McClary H**, Koch R, Chisari FV, Guidotti LG. Relative sensitivity of hepatitis B virus and other hepatotropic viruses to the antiviral effects of cytokines. *J Virol* 2000; **74**: 2255-2264 [PMID: [10666256](#) DOI: [10.1128/jvi.74.5.2255-2264.2000](#)]
 - 24 **Carreño V**, Zeuzem S, Hopf U, Marcellin P, Cooksley WG, Fevery J, Diago M, Reddy R, Peters M, Rittweger K, Rakhit A, Pardo M. A phase I/II study of recombinant human interleukin-12 in patients with chronic hepatitis B. *J Hepatol* 2000; **32**: 317-324 [PMID: [10707873](#) DOI: [10.1016/S0168-8278\(00\)80078-1](#)]
 - 25 **Rigopoulou EI**, Suri D, Chokshi S, Mullerova I, Rice S, Tedder RS, Williams R, Naoumov NV. Lamivudine plus interleukin-12 combination therapy in chronic hepatitis B: antiviral and immunological activity. *Hepatology* 2005; **42**: 1028-1036 [PMID: [16250037](#) DOI: [10.1002/hep.20888](#)]
 - 26 **Rossol S**, Marinos G, Carucci P, Singer MV, Williams R, Naoumov NV. Interleukin-12 induction of Th1 cytokines is important for viral clearance in chronic hepatitis B. *J Clin Invest* 1997; **99**: 3025-3033 [PMID: [9185527](#) DOI: [10.1172/JCI119498](#)]
 - 27 **Ting SW**, Chen YC, Huang YH. Risk of Hepatitis B Reactivation in Patients with Psoriasis on Ustekinumab. *Clin Drug Investig* 2018; **38**: 873-880 [PMID: [29968197](#) DOI: [10.1007/s40261-018-0671-z](#)]
 - 28 **Chiu HY**, Chen CH, Wu MS, Cheng YP, Tsai TF. The safety profile of ustekinumab in the treatment of patients with psoriasis and concurrent hepatitis B or C. *Br J Dermatol* 2013; **169**: 1295-1303

- [PMID: [23746170](#) DOI: [10.1111/bjd.12461](#)]
- 29 **Reich K**, Papp KA, Blauvelt A, Tying SK, Sinclair R, Thaçi D, Nogales K, Mehta A, Cichanowitz N, Li Q, Liu K, La Rosa C, Green S, Kimball AB. Tildrakizumab vs placebo or etanercept for chronic plaque psoriasis (reSURFACE 1 and reSURFACE 2): results from two randomised controlled, phase 3 trials. *Lancet* 2017; **390**: 276-288 [PMID: [28596043](#) DOI: [10.1016/S0140-6736\(17\)31279-5](#)]
 - 30 **Papp K**, Thaçi D, Reich K, Riedl E, Langley RG, Krueger JG, Gottlieb AB, Nakagawa H, Bowman EP, Mehta A, Li Q, Zhou Y, Shames R. Tildrakizumab (MK-3222), an anti-interleukin-23p19 monoclonal antibody, improves psoriasis in a phase IIb randomized placebo-controlled trial. *Br J Dermatol* 2015; **173**: 930-939 [PMID: [26042589](#) DOI: [10.1111/bjd.13932](#)]
 - 31 **Gordon KB**, Strober B, Lebwohl M, Augustin M, Blauvelt A, Poulin Y, Papp KA, Sofen H, Puig L, Foley P, Ohtsuki M, Flack M, Geng Z, Gu Y, Valdes JM, Thompson EH, Bachelez H. Efficacy and safety of risankizumab in moderate-to-severe plaque psoriasis (UltIMMA-1 and UltIMMA-2): results from two double-blind, randomised, placebo-controlled and ustekinumab-controlled phase 3 trials. *Lancet* 2018; **392**: 650-661 [PMID: [30097359](#) DOI: [10.1016/S0140-6736\(18\)31713-6](#)]
 - 32 **Feagan BG**, Sandborn WJ, D'Haens G, Panés J, Kaser A, Ferrante M, Louis E, Franchimont D, Dewit O, Seidler U, Kim KJ, Neurath MF, Schreiber S, Scholl P, Pamulapati C, Lalovic B, Visvanathan S, Padula SJ, Herichova I, Soaita A, Hall DB, Böcher WO. Induction therapy with the selective interleukin-23 inhibitor risankizumab in patients with moderate-to-severe Crohn's disease: a randomised, double-blind, placebo-controlled phase 2 study. *Lancet* 2017; **389**: 1699-1709 [PMID: [28411872](#) DOI: [10.1016/S0140-6736\(17\)30570-6](#)]
 - 33 **Gordon KB**, Duffin KC, Bissonnette R, Prinz JC, Wasfi Y, Li S, Shen YK, Szapary P, Randazzo B, Reich K. A Phase 2 Trial of Guselkumab vs Adalimumab for Plaque Psoriasis. *N Engl J Med* 2015; **373**: 136-144 [PMID: [26154787](#) DOI: [10.1056/NEJMoa1501646](#)]
 - 34 **Reich K**, Armstrong AW, Foley P, Song M, Wasfi Y, Randazzo B, Li S, Shen YK, Gordon KB. Efficacy and safety of guselkumab, an anti-interleukin-23 monoclonal antibody, compared with adalimumab for the treatment of patients with moderate to severe psoriasis with randomized withdrawal and retreatment: Results from the phase III, double-blind, placebo- and active comparator-controlled VOYAGE 2 trial. *J Am Acad Dermatol* 2017; **76**: 418-431 [PMID: [28057361](#) DOI: [10.1016/j.jaad.2016.11.042](#)]
 - 35 **Sands BE**, Chen J, Feagan BG, Penney M, Rees WA, Danese S, Higgins PDR, Newbold P, Faggioni R, Patra K, Li J, Klekotka P, Morehouse C, Pulkstenis E, Drappa J, van der Merwe R, Gasser RA Jr. Efficacy and Safety of MEDI2070, an Antibody Against Interleukin 23, in Patients With Moderate to Severe Crohn's Disease: A Phase 2a Study. *Gastroenterology* 2017; **153**: 77-86. e6 [PMID: [28390867](#) DOI: [10.1053/j.gastro.2017.03.049](#)]
 - 36 **Frieder J**, Kivelevitch D, Haugh I, Watson I, Menter A. Anti-IL-23 and Anti-IL-17 Biologic Agents for the Treatment of Immune-Mediated Inflammatory Conditions. *Clin Pharmacol Ther* 2018; **103**: 88-101 [PMID: [28960267](#) DOI: [10.1002/cpt.893](#)]
 - 37 **Zhang JY**, Zhang Z, Lin F, Zou ZS, Xu RN, Jin L, Fu JL, Shi F, Shi M, Wang HF, Wang FS. Interleukin-17-producing CD4(+) T cells increase with severity of liver damage in patients with chronic hepatitis B. *Hepatology* 2010; **51**: 81-91 [PMID: [19842207](#) DOI: [10.1002/hep.23273](#)]
 - 38 **Xia L**, Tian D, Huang W, Zhu H, Wang J, Zhang Y, Hu H, Nie Y, Fan D, Wu K. Upregulation of IL-23 expression in patients with chronic hepatitis B is mediated by the HBx/ERK/NF-κB pathway. *J Immunol* 2012; **188**: 753-764 [PMID: [22174449](#) DOI: [10.4049/jimmunol.1101652](#)]
 - 39 **Ge J**, Wang K, Meng QH, Qi ZX, Meng FL, Fan YC. Implication of Th17 and Th1 cells in patients with chronic active hepatitis B. *J Clin Immunol* 2010; **30**: 60-67 [PMID: [19756987](#) DOI: [10.1007/s10875-009-9328-2](#)]
 - 40 **Yu C**, Gong X, Yang Q, Lian J, Xu K, Ruan B, Li L. The serum IL-23 Level predicts the response to pegylated interferon therapy in patients with chronic hepatitis B. *Liver Int* 2015; **35**: 1549-1556 [PMID: [25312687](#) DOI: [10.1111/liv.12701](#)]
 - 41 **Koskinas J**, Tampaki M, Doumba PP, Rallis E. Hepatitis B virus reactivation during therapy with ustekinumab for psoriasis in a hepatitis B surface-antigen-negative anti-HBs-positive patient. *Br J Dermatol* 2013; **168**: 679-680 [PMID: [23121260](#) DOI: [10.1111/bjd.12120](#)]
 - 42 **Mitsdoerffer M**, Lee Y, Jäger A, Kim HJ, Korn T, Kolls JK, Cantor H, Bettelli E, Kuchroo VK. Proinflammatory T helper type 17 cells are effective B-cell helpers. *Proc Natl Acad Sci USA* 2010; **107**: 14292-14297 [PMID: [20660725](#) DOI: [10.1073/pnas.1009234107](#)]
 - 43 **Duncan JR**, Orlowski TJ, Elewski BE. Safety of guselkumab in hepatitis B virus infection. *Dermatol Online J* 2019; **25** [PMID: [31735016](#)]
 - 44 **Curtis MM**, Way SS. Interleukin-17 in host defence against bacterial, mycobacterial and fungal pathogens. *Immunology* 2009; **126**: 177-185 [PMID: [19125888](#) DOI: [10.1111/j.1365-2567.2008.03017.x](#)]
 - 45 **Langley RG**, Elewski BE, Lebwohl M, Reich K, Griffiths CE, Papp K, Puig L, Nakagawa H, Spelman L, Sigurgeirsson B, Rivas E, Tsai TF, Wasel N, Tying S, Salko T, Hampele I, Notter M, Karpov A, Helou S, Papavassilis C; ERASURE Study Group; FIXTURE Study Group. Secukinumab in plaque psoriasis--results of two phase 3 trials. *N Engl J Med* 2014; **371**: 326-338 [PMID: [25007392](#) DOI: [10.1056/NEJMoa1314258](#)]
 - 46 **Gordon KB**, Blauvelt A, Papp KA, Langley RG, Luger T, Ohtsuki M, Reich K, Amato D, Ball SG, Braun DK, Cameron GS, Erickson J, Konrad RJ, Muram TM, Nickoloff BJ, Osuntokun OO, Secrest RJ, Zhao F, Mallbris L, Leonardi CL; UNCOVER-1 Study Group; UNCOVER-2 Study Group; UNCOVER-3 Study Group. Phase 3 Trials of Ixekizumab in Moderate-to-Severe Plaque Psoriasis. *N*

- Engl J Med* 2016; **375**: 345-356 [PMID: [27299809](#) DOI: [10.1056/NEJMoa1512711](#)]
- 47 **Lebwohl M**, Strober B, Menter A, Gordon K, Weglowska J, Puig L, Papp K, Spelman L, Toth D, Kerdel F, Armstrong AW, Stingl G, Kimball AB, Bachelez H, Wu JJ, Crowley J, Langley RG, Blicharski T, Paul C, Lacour JP, Tying S, Kircik L, Chimenti S, Callis Duffin K, Bagel J, Koo J, Aras G, Li J, Song W, Milmont CE, Shi Y, Erond N, Klekotka P, Kotzin B, Nirula A. Phase 3 Studies Comparing Brodalumab with Ustekinumab in Psoriasis. *N Engl J Med* 2015; **373**: 1318-1328 [PMID: [26422722](#) DOI: [10.1056/NEJMoa1503824](#)]
 - 48 **Nash P**, Kirkham B, Okada M, Rahman P, Combe B, Burmester GR, Adams DH, Kerr L, Lee C, Shuler CL, Genovese M; SPIRIT-P2 Study Group. Ixekizumab for the treatment of patients with active psoriatic arthritis and an inadequate response to tumour necrosis factor inhibitors: results from the 24-week randomised, double-blind, placebo-controlled period of the SPIRIT-P2 phase 3 trial. *Lancet* 2017; **389**: 2317-2327 [PMID: [28551073](#) DOI: [10.1016/S0140-6736\(17\)31429-0](#)]
 - 49 **Baeten D**, Sieper J, Braun J, Baraliakos X, Dougados M, Emery P, Deodhar A, Porter B, Martin R, Andersson M, Mpfu S, Richards HB; MEASURE 1 Study Group; MEASURE 2 Study Group. Secukinumab, an Interleukin-17A Inhibitor, in Ankylosing Spondylitis. *N Engl J Med* 2015; **373**: 2534-2548 [PMID: [26699169](#) DOI: [10.1056/NEJMoa1505066](#)]
 - 50 **Paquissi FC**. Immunity and Fibrogenesis: The Role of Th17/IL-17 Axis in HBV and HCV-induced Chronic Hepatitis and Progression to Cirrhosis. *Front Immunol* 2017; **8**: 1195 [PMID: [29033929](#) DOI: [10.3389/fimmu.2017.01195](#)]
 - 51 **Feng H**, Yin J, Han YP, Zhou XY, Chen S, Yang L, Yan JR, Zhang GX. Sustained Changes of Treg and Th17 Cells During Interferon- α Therapy in Patients with Chronic Hepatitis B. *Viral Immunol* 2015; **28**: 412-417 [PMID: [26266573](#) DOI: [10.1089/vim.2015.0024](#)]
 - 52 **Chiu HY**, Hui RC, Huang YH, Huang RY, Chen KL, Tsai YC, Lai PJ, Wang TS, Tsai TF. Safety Profile of Secukinumab in Treatment of Patients with Psoriasis and Concurrent Hepatitis B or C: A Multicentric Prospective Cohort Study. *Acta Derm Venereol* 2018; **98**: 829-834 [PMID: [29972221](#) DOI: [10.2340/00015555-2989](#)]
 - 53 **Koike Y**, Fujiki Y, Higuchi M, Fukuchi R, Kuwatsuka S, Murota H. An interleukin-17 inhibitor successfully treated a complicated psoriasis and psoriatic arthritis patient with hepatitis B virus infection and end-stage kidney disease on hemodialysis. *JAAD Case Rep* 2019; **5**: 150-152 [PMID: [30733983](#) DOI: [10.1016/j.jdc.2018.11.016](#)]
 - 54 **Lora V**, Graceffa D, De Felice C, Morrone A, Bonifati C. Treatment of severe psoriasis with ixekizumab in a liver transplant recipient with concomitant hepatitis B virus infection. *Dermatol Ther* 2019; **32**: e12909 [PMID: [30964590](#) DOI: [10.1111/dth.12909](#)]
 - 55 **Winthrop KL**. The emerging safety profile of JAK inhibitors in rheumatic disease. *Nat Rev Rheumatol* 2017; **13**: 234-243 [PMID: [28250461](#) DOI: [10.1038/nrrheum.2017.23](#)]
 - 56 **Burmester GR**, Blanco R, Charles-Schoeman C, Wollenhaupt J, Zerbini C, Benda B, Gruben D, Wallenstein G, Krishnaswami S, Zwillich SH, Konec T, Soma K, Bradley J, Mebus C; ORAL Step investigators. Tofacitinib (CP-690,550) in combination with methotrexate in patients with active rheumatoid arthritis with an inadequate response to tumour necrosis factor inhibitors: a randomised phase 3 trial. *Lancet* 2013; **381**: 451-460 [PMID: [23294500](#) DOI: [10.1016/S0140-6736\(12\)61424-X](#)]
 - 57 **Kremer J**, Li ZG, Hall S, Fleischmann R, Genovese M, Martin-Mola E, Isaacs JD, Gruben D, Wallenstein G, Krishnaswami S, Zwillich SH, Konec T, Riese R, Bradley J. Tofacitinib in combination with nonbiologic disease-modifying antirheumatic drugs in patients with active rheumatoid arthritis: a randomized trial. *Ann Intern Med* 2013; **159**: 253-261 [PMID: [24026258](#) DOI: [10.7326/0003-4819-159-4-201308200-00006](#)]
 - 58 **Gladman D**, Rigby W, Azevedo VF, Behrens F, Blanco R, Kaszuba A, Kudlacz E, Wang C, Menon S, Hendrikx T, Kanik KS. Tofacitinib for Psoriatic Arthritis in Patients with an Inadequate Response to TNF Inhibitors. *N Engl J Med* 2017; **377**: 1525-1536 [PMID: [29045207](#) DOI: [10.1056/NEJMoa1615977](#)]
 - 59 **Mease P**, Hall S, FitzGerald O, van der Heijde D, Merola JF, Avila-Zapata F, Cieślak D, Graham D, Wang C, Menon S, Hendrikx T, Kanik KS. Tofacitinib or Adalimumab vs Placebo for Psoriatic Arthritis. *N Engl J Med* 2017; **377**: 1537-1550 [PMID: [29045212](#) DOI: [10.1056/NEJMoa1615975](#)]
 - 60 **Sandborn WJ**, Su C, Sands BE, D'Haens GR, Vermeire S, Schreiber S, Danese S, Feagan BG, Reinisch W, Niezychowski W, Friedman G, Lawendy N, Yu D, Woodworth D, Mukherjee A, Zhang H, Healey P, Panés J; OCTAVE Induction 1, OCTAVE Induction 2, and OCTAVE Sustain Investigators. Tofacitinib as Induction and Maintenance Therapy for Ulcerative Colitis. *N Engl J Med* 2017; **376**: 1723-1736 [PMID: [28467869](#) DOI: [10.1056/NEJMoa1606910](#)]
 - 61 **Sandborn WJ**, Ghosh S, Panes J, Vranic I, Su C, Rousell S, Niezychowski W; Study A3921063 Investigators. Tofacitinib, an oral Janus kinase inhibitor, in active ulcerative colitis. *N Engl J Med* 2012; **367**: 616-624 [PMID: [22894574](#) DOI: [10.1056/NEJMoa1112168](#)]
 - 62 **Harigai M**, Winthrop K, Takeuchi T, Hsieh TY, Chen YM, Smolen JS, Burmester G, Walls C, Wu WS, Dickson C, Liao R, Genovese MC. Evaluation of hepatitis B virus in clinical trials of baricitinib in rheumatoid arthritis. *RMD Open* 2020; **6** [PMID: [32098857](#) DOI: [10.1136/rmdopen-2019-001095](#)]
 - 63 **Kirito K**, Sakamoto M, Enomoto N. Elevation of the Hepatitis B Virus DNA during the Treatment of Polycythemia Vera with the JAK Kinase Inhibitor Ruxolitinib. *Intern Med* 2016; **55**: 1341-1344 [PMID: [27181544](#) DOI: [10.2169/internalmedicine.55.5529](#)]
 - 64 **Rehermann B**, Nascimbeni M. Immunology of hepatitis B virus and hepatitis C virus infection. *Nat Rev Immunol* 2005; **5**: 215-229 [PMID: [15738952](#) DOI: [10.1038/nri1573](#)]
 - 65 **Heine A**, Held SA, Daecke SN, Wallner S, Yajnanarayana SP, Kurts C, Wolf D, Brossart P. The

- JAK-inhibitor ruxolitinib impairs dendritic cell function *in vitro* and *in vivo*. *Blood* 2013; **122**: 1192-1202 [PMID: 23770777 DOI: 10.1182/blood-2013-03-484642]
- 66 **Chen YM**, Huang WN, Wu YD, Lin CT, Chen YH, Chen DY, Hsieh TY. Reactivation of hepatitis B virus infection in patients with rheumatoid arthritis receiving tofacitinib: a real-world study. *Ann Rheum Dis* 2018; **77**: 780-782 [PMID: 28663308 DOI: 10.1136/annrheumdis-2017-211322]
- 67 **Serling-Boyd N**, Mohareb AM, Kim AY, Hyle EP, Wallace ZS. The use of tocilizumab and tofacitinib in patients with resolved hepatitis B infection: a case series. *Ann Rheum Dis* 2021; **80**: 274-276 [PMID: 32732241 DOI: 10.1136/annrheumdis-2020-218289]
- 68 **Sanz-Bueno J**, Vanaclocha F, García-Doval I, Torrado R, Carretero G, Daudén E, Patricia Ruiz-Genao D, Alsina-Gibert MM, Pérez-Zafrilla B, Pérez-Rial G, Rivera R; members of the BIOBADADERM group. Risk of Reactivation of Hepatitis B Virus Infection in Psoriasis Patients Treated With Biologics: A Retrospective Analysis of 20 Cases From the BIOBADADERM Database. *Actas Dermosifiliogr* 2015; **106**: 477-482 [PMID: 25776200 DOI: 10.1016/j.ad.2015.01.010]
- 69 **Navarro R**, Vilarrasa E, Herranz P, Puig L, Bordas X, Carrascosa JM, Taberner R, Ferrán M, García-Bustinduy M, Romero-Maté A, Pedragosa R, García-Diez A, Daudén E. Safety and effectiveness of ustekinumab and antitumour necrosis factor therapy in patients with psoriasis and chronic viral hepatitis B or C: a retrospective, multicentre study in a clinical setting. *Br J Dermatol* 2013; **168**: 609-616 [PMID: 22985451 DOI: 10.1111/bjd.12045]
- 70 **Hayashi M**, Umezawa Y, Fukuchi O, Ito T, Saeki H, Nakagawa H. Efficacy and safety of ustekinumab treatment in elderly patients with psoriasis. *J Dermatol* 2014; **41**: 974-980 [PMID: 25346301 DOI: 10.1111/1346-8138.12653]
- 71 **Steglich RB**, Meneghello LP, Carvalho AV, Cheinquer H, Muller FM, Reginatto FP. The use of ustekinumab in a patient with severe psoriasis and positive HBV serology. *An Bras Dermatol* 2014; **89**: 652-654 [PMID: 25054756 DOI: 10.1590/abd1806-4841.20143013]
- 72 **Moneva-Leniz LM**, Sahuquillo-Torralba A, Vila-Payeras A, Mateu-Puchades A. Risk of Hepatitis B Virus Reactivation in Patients on Secukinumab for Psoriasis: A Series of 4 Cases. *Actas Dermosifiliogr* 2020; **111**: 613-614 [PMID: 32589963 DOI: 10.1016/j.ad.2019.02.022]
- 73 **Feaster B**, Cline A, Feldman SR. Secukinumab for psoriasis in a patient with hepatitis B. *Dermatol Online J* 2018; **24** [PMID: 30677838]
- 74 **Bevans SL**, Mayo TT, Elewski BE. Safety of secukinumab in hepatitis B virus. *J Eur Acad Dermatol Venereol* 2018; **32**: e120-e121 [PMID: 28960490 DOI: 10.1111/jdv.14608]
- 75 **Yanagihara S**, Sugita K, Yoshida Y, Tsuruta D, Yamamoto O. Psoriasis vulgaris in a hepatitis B virus carrier successfully treated with secukinumab and entecavir combination therapy. *Eur J Dermatol* 2017; **27**: 185-186 [PMID: 27965188 DOI: 10.1684/ejd.2016.2939]
- 76 **Peccerillo F**, Odorici G, Pellacani G, Conti A. Secukinumab: A positive outcome in a patient with severe psoriasis and HBV-HCV co-infection. *Dermatol Ther* 2018; **31**: e12601 [PMID: 29633448 DOI: 10.1111/dth.12601]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

