

国内版 国际版

The risk of hepatitis B virus reactivation in patients with autoimmuni



ALL IMAGES VIDEOS

11,400 Results Any time ▾

Hepatitis B virus reactivation in hepatitis B virus ...

<https://europepmc.org/article/MED/23898372> ▾

Jul 01, 2013 - Core tip: Patients with occult or resolved hepatitis B undergoing chemotherapy or immunosuppression are potentially at risk of hepatitis B virus (HBV) reactivation which can be disastrous since it can lead to acute liver failure and death. In this report, we describe the characteristic and outcome in one of the larger series of patients (n = 14) with occult or resolved HBV who ...

Cited by: 28 Author: Kalliopi Zachou, Alexandros Sarantopoulo...
Publish Year: 2013

Risk of hepatitis B virus reactivation in rheumatoid ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC25848463>

Mar 27, 2015 - Core tip: Hepatitis B virus (HBV) infection represents a major issue in patients with rheumatoid arthritis (RA) undergoing biological disease-modifying anti-rheumatic drugs (bDMARDs). While several observational studies and trials deal with the risk of HBV reactivation under anti-TNF agents, there is limited experience with newer drugs as tocilizumab (TCZ) and abatacept (ABA).

Cited by: 63 Author: Francesca De Nard, Monica Todoerti, Vitt...
Publish Year: 2015

Hepatitis B virus persistence and reactivation | The BMJ

<https://www.bmj.com/content/370/bmj.m2200> ▾

Sep 01, 2020 - Hepatitis B virus (HBV) infection causes chronic hepatitis and has long term complications. Individuals ever infected with HBV are at risk of viral reactivation under certain circumstances. This review summarizes studies on HBV persistence and reactivation with a focus on the definitions and mechanisms. Emphasis is placed on the interplay between HBV replication and ...

Author: Yu Shi, Min Zheng Publish Year: 2020

Prevention of Hepatitis B reactivation in the setting of ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC27291888>

INTRODUCTION. Hepatitis B is a disease prevalent globally, with approximately 2 billion people exposed to hepatitis B and of these, 240 million are chronically infected (chronic hepatitis B, CHB) []. This review will highlight the impact of concomitant hepatitis B virus (HBV) infection amongst patients with malignant, inflammatory and autoimmune conditions who undergo immunosuppression.

Cited by: 67 Author: Venessa Pattullo
Publish Year: 2016

Moderate Risk of Hepatitis B Virus Reactivation in HBsAg ...

<https://www.nature.com/articles/s41598-020-59406-4>

Feb 12, 2020 · Hepatitis B reactivation in patients with previous hepatitis B virus exposure undergoing rituximab-containing chemotherapy for lymphoma: a prospective study. *J Clin Oncol* 32, 3736–3743, <https://doi.org/10.1200/JCO.2019.42.19.3736>

Cited by: 3 Author: Meng Hsuan Kuo, Chih-Wei Tseng, Chi-H...

PDF Risk of hepatitis B virus reactivation in rheumatoid ...

<https://air.unimi.it/retrieve/handle/2434/662680/1291355/WJH-7-344.pdf>

Virus, hepatitis C and hepatitis B virus (HBV). The prevalence of HBV infection in patients affected by rheumatic diseases is similar to the one of the general population. In the last decade, increasing attention has been paid to the possibility of HBV reactivation in RA patients undergoing ...

Risk of hepatitis B virus reactivation in rheumatoid ...

europepmc.org/articles/PMC4381162

Mar 01, 2015 · Core tip: Hepatitis B virus (HBV) infection represents a major issue in patients with rheumatoid arthritis (RA) undergoing biological disease-modifying anti-rheumatic drugs (bDMARDs). While several observational studies and trials deal with the risk of HBV reactivation under anti-TNF agents, there is limited experience with newer drugs as tocilizumab (TCZ) and abatacept (ABA).

Cited by: 63 Author: Francesca De Nard, Monica Todoerti, Vit...

Hepatitis B virus

Dna Virus



Hepatitis B virus, is a partially double-stranded DNA virus, a species of the genus Orthohepadnavirus and a member of the Hepadnaviridae family of viruses. This virus causes the disease hepatitis B.

Wikipedia

Scientific name: Hepatitis B virus

Family: Hepadnaviridae

Genus: Orthohepadnavirus

Domain: Virus


Class: Reoviricetes

Biological rank: Species

Data from: Wikipedia Freebase

Suggest an edit

激活 Windows
转到“设置”以激活 Windows。



Name of Journal: *World Journal of Gastroenterology*
Manuscript NO: 63080
Manuscript Type: MINIREVIEWS

Risk of hepatitis B virus reactivation in patients with autoimmune diseases undergoing non-tumor necrosis factor-targeted biologics

HBV reactivation and non-TNF-targeted biologics

Shintaro Akiyama, Thomas G. Cotter, Atsushi Sakuraba

Match Overview

Rank	Source	Words	Percentage
1	Crossref: Anthony Myint, Myron J. Tong, Simon W. Beaven. "Reactivation of Hepatitis B Virus: A Review of Clinical Guidelines"	122	3%
2	Crossref: Chiu, H.-K., C.-H. Chen, M.-S. Wu, Y.-P. Cheng, and T.-F. Tsai. "The safety profile of ustekinumab in the treatment..."	50	1%
3	Crossref: Tae Hoon Lee, Thomas D. Szstano. "Hepatology Clearance for Immunomodulation in Patients With Hepatitis B"	38	1%
4	Internet: crawled on 13-Jan-2020 www.nature.com	33	1%
5	Crossref: Chengbo Yu, Xianli Gong, Qian Yang, Jiangshan Lian, Kaijin Xu, Bin Ruan, Lanjuan Li. "The serum IL-23 level..."	29	1%
6	Internet: crawled on 14-Jul-2020 deepblue.lib.umich.edu	17	<1%

国内版 国际版

Risk of hepatitis B virus reactivation in patients with autoimmune di:



ALL IMAGES VIDEOS

12,300 Results Any time ▾

Moderate Risk of Hepatitis B Virus Reactivation in HBsAg ...

<https://www.nature.com/articles/s41598-020-59406-4>

Feb 12, 2020 · Hepatitis B reactivation in patients with previous hepatitis B virus exposure undergoing rituximab-containing chemotherapy for lymphoma: a prospective study. J Clin Oncol 32 , 3736–3743, [https ...](https://doi.org/10.1200/JCO.2019.18.10000)

Cited by: 3 Author: Meng Hsuan Kuo, Chih-Wei Tseng, Chi-H...

Publish Year: 2020

Risk of hepatitis B virus reactivation in rheumatoid ...

europepmc.org/articles/PMC4381162

Mar 01, 2015 · Core tip: Hepatitis B virus (HBV) infection represents a major issue in patients with rheumatoid arthritis (RA) undergoing biological disease-modifying anti-rheumatic drugs (bDMARDs). While several observational studies and trials deal with the risk of HBV reactivation under anti-TNF agents, there is limited experience with newer drugs as tocilizumab (TCZ) and abatacept (ABA).

Cited by: 63 Author: Francesca De Nard, Monica Todoerti, Vitt...

Publish Year: 2015

[PDF] Risk of hepatitis B virus reactivation in rheumatoid ...

<https://air.unimi.it/retrieve/handle/2434/662680/1291355/WJH-7-344.pdf>

Virus, hepatitis C and hepatitis B virus (HBV). The prevalence of HBV infection in patients affected by rheumatic diseases is similar to the one of the general population. In the last decade, increasing attention has been paid to the possibility of HBV reactivation in RA patients undergoing ...

Infection risk in patients undergoing treatment for ...

<https://pubmed.ncbi.nlm.nih.gov/31852268>

Introduction: Despite the therapeutic effectiveness of biologics targeting immune cells or cytokines in patients with inflammatory arthritis, which reflects their pathogenic roles, an increased infection risk is observed in those undergoing biological treatment. However, there are limited data regarding the comparison of infection risks in inflammatory arthritis patients treated with non ...

Cited by: 7 Author: Ying-Ming Chiu, Der-Yuan Chen

Publish Year: 2020

Hepatitis B Reactivation During Immunosuppressive Therapy ...

europepmc.org/articles/PMC4887960