

Management of afferent loop obstruction: Reoperation or ...

<https://pubmed.ncbi.nlm.nih.gov/26425267>

Surgery includes: (1) adhesiolysis and reconstruction for benign causes; and (2) by-pass or excision and reconstruction for malignant causes. However, endoscopic **enteral stenting**, transhepatic percutaneous...

Cited by: 23 Author: Konstantinos Blouhos, Konstantinos Andr...
Publish Year: 2015

Afferent Loop Syndrome Article - StatPearls

<https://www.statpearls.com/articlelibrary/viewarticle/17273>

Nov 24, 2020 - Patients with acute **afferent loop syndrome** due to malignant causes that are deemed non-operative should undergo palliative treatment targeted at relieving the **acute obstruction**. Excision...

Author: Rebecca K. Grotewiel, Renford Cindass Publish Year: 2019

PEOPLE ALSO ASK

- What is the treatment for afferent loop syndrome? ▾
- What is the function of the afferent loop? ▾
- What is the distal jejunal segment? ▾

Feedback

[PDF Management of Obstructed Afferent-Loop due to ...](#)

Related searches

- [afferent loop syndrome treatment](#)
- [what is afferent loop syndrome](#)
- [afferent loop syndrome radiology](#)
- [afferent loop syndrome lcd 10](#)
- [afferent loop syndrome ct](#)

Search Tools

Turn off Hover Translation (关闭取词)

Microsoft Bing

国内版 国际版

Chat with Bing

Sign in

Add the Bing with extension

Clinical management for malignant afferent loop obstruction

ALL IMAGES VIDEOS

23,900 Results Any time

Management of afferent loop obstruction: Reoperation or ...
<https://pubmed.ncbi.nlm.nih.gov/26425267>
Surgery includes: (1) adhesiolysis and reconstruction for benign causes; and (2) by-pass or excision and reconstruction for malignant causes. However, endoscopic enteral stenting, transhepatic percutaneous enteral stenting and direct percutaneous tube enterostomy have the principal role in management of malignant and radiation-induced obstruction.
Cited by: 25 Author: Konstantinos Blouhos, Konstantinos And...
Publish Year: 2015

Management of afferent loop obstruction: Reoperation or ...
<https://europepmc.org/articles/PMC4582236>
Sep 01, 2015 Core tip: Management strategy of afferent loop obstruction (ALO) depends on: (1) the benign or malignant nature of the obstruction. ALO caused by a benign lesion needs definitive repair of the primary cause by surgery. ALO caused by a malignant lesion needs palliative treatment (percutaneous and endoscopic interventions, by-pass surgery) or excision; and (2) the site of obstruction.
Cited by: 25 Author: Konstantinos Blouhos, Konstantinos And...
Publish Year: 2015

Related searches

- afferent loop obstruction
- afferent limb obstruction
- afferent loop syndrome treatment
- afferent loop syndrome
- afferent loop syndrome radiology
- afferent loop syndrome icd 10
- afferent loop syndrome ct

Search Tools

Turn off Hover Translation (关闭翻译)

激活 Windows
转到“设置”以激活 Windows。

Name of Journal: *World Journal of Gastrointestinal Oncology*

Manuscript NO: 65339

Manuscript Type: MINIREVIEWS

Clinical management for malignant afferent loop obstruction

Clinical management for malignant afferent loop obstruction

Arata Sakai, Hideyuki Shiomi, Atsuhiko Masuda, Takashi Kobayashi, Yasutaka Yamada,
Yuzo Kodama

Match Overview

Rank	Source	Words	Percentage
1	Internet www.ncbi.nlm.nih.gov	115 words	5%
2	Crossref Arata Sakai, Hideyuki Shiomi, Atsuhiko Masuda, Takashi Kobayashi, Yasutaka Yamada, Yuzo Kodama. "Effectiveness of endoscopic self-expandable"	64 words	3%
3	Internet crawled on 08-Jan-2021 pesquisa.bvsalud.org	61 words	3%
4	Internet crawled on 01-Feb-2021 www.elsevier.com	56 words	3%
5	Crossref Akshika Kida, Hidenori Kido, Toshiaki Matsuo, Atsuyoshi M Izukami et al. "Usefulness of endoscopic metal stent p..."	51 words	2%
6	Internet crawled on 05-Dec-2020 synapse.koreamed.org	45 words	2%
	Crossref 转到"设置"以激活 Windows...	34 words	

国内版 国际版

Clinical management for malignant afferent loop obstruction



ALL IMAGES VIDEOS

120,000 Results Any time ▾

Management for patients with **afferent loop** syndrome can largely subdivide according to underlying etiology. For most benign causes, surgery usually provides definitive treatment. In the setting of **malignancy**, initial neoadjuvant **management** may be followed by surgical treatment with curative/palliative intent.

Author: Rebecca K. Grotewiel, Renford Cindass

Publish Year: 2019

[Afferent Loop Syndrome Article - StatPearls](#)

www.statpearls.com/articlelibrary/viewarticle/17273/

Was this helpful?

PEOPLE ALSO ASK

Is there a cure for afferent loop syndrome? ▾

What are the symptoms of afferent loop syndrome? ▾

What causes an afferent loop in the limb? ▾

When does afferent loop syndrome occur after anastomosis? ▾

Feedback

[Management of afferent loop obstruction: Reoperation or ...](#)

<https://pubmed.ncbi.nlm.nih.gov/26425267>

Surgery includes: (1) adhesiolysis and reconstruction for benign causes; and (2) by-pass or excision and reconstruction for malignant causes. However, endoscopic **enteral stenting**, transhepatic percutaneous...

Cited by: 27

Author: Konstantinos Blouhos, Konstantinos Andre...

Publish Year: 2015

[Management of Obstructed Afferent-Loop due to Recurrence ...](#)