

**Risk of bile duct carcinogenesis after excision of ...**<https://www.sciencedirect.com/science/article/pii/S003960609970036X>

Nov 01, 1999 · Background: A reflux of pancreatic juice into the biliary tract caused by pancreaticobiliary maljunction (PBM) has been considered important in the development of biliary tract carcinogenesis i...

Cited by: 194

Author: Susumu Kobayashi, Takehide Asano, Ma...

Publish Year: 1999

**[PDF] Bile Duct Cyst as Precursor to Biliary Tract Cancer**<https://link.springer.com/content/pdf/10.1245/s10434-006-9294-3.pdf>

Bile duct cysts (BDC) are rare and of unknown origin,1 with a predominating occurrence in females of the Asian population. BDC may represent a pre-cursor for biliary tract cancer development, especially in...

Cited by: 81

Author: Kjetil Søreide, Jon Arne Søreide, Jon Arn...

Publish Year: 2007

## PEOPLE ALSO ASK

What is the risk of biliary tract malignancy? ▾

Can obesity cause cholangiocarcinoma? ▾

What is bile duct sclerosis? ▾

Is there a biliary change in cholangiocarcinoma? ▾

[Feedback](#)**Clinical presentation, risk factors and staging systems of ...**<https://www.sciencedirect.com/science/article/pii/S1521691815000177>

Apr 01, 2015 · In patients with bile-duct cysts the incidence of CCA in the first decade of their life and the risk of malignancy decreases after cyst excision , , , . Other confirmed risk factors are virus infection...

**The risk of carcinogenesis in congenital choledochal cyst ...**<https://www.sciencedirect.com/science/article/pii/S1665268119309858>

国内版

国际版

Risk factors for preoperative carcinogenesis of bile duct cysts in adu



ALL

IMAGES

VIDEOS

31,900 Results

Any time ▾

### [The risk of carcinogenesis in congenital choledochal cyst ...](#)

<https://www.sciencedirect.com/science/article/pii/S1665268119309858>

Nov 01, 2014 · Bile stasis due to stenosis of the intrahepatic **bile duct** and **repeated cholangitis** with multiple stones may be a high-risk factor. 32 Another reasonable explanation is that the epithelium of the remnant bile duct wall may have already progressed to a precancerous stage by the time of the operation and that genetic changes may have occurred or continued during the postoperative period. 33

Cited by: 25

Author: Xiao-Dong He, Lei Wang, Wei Liu, Qi Liu,...

Publish Year: 2014

### [\[PDF\] Bile Duct Cyst as Precursor to Biliary Tract Cancer](#)

<https://link.springer.com/content/pdf/10.1245/s10434-006-9294-3.pdf>

**Bile duct cysts** (BDC) are rare and of unknown origin,<sup>1</sup> with a predominating occurrence in females of the Asian population. BDC may represent a pre-cursor for biliary tract cancer development, especially in **adults**.<sup>2–4</sup> While more than 60% of patients with **bile duct cysts** are diagnosed during the first decade

Cited by: 81

Author: Kjetil Søreide, Jon Arne Søreide, Jon Arn...

Publish Year: 2007

**3** Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 66955

Manuscript Type: ORIGINAL ARTICLE

*Retrospective Study*

Risk factors for preoperative carcinogenesis of bile duct cysts in adults

Xin Wu, Bing-Lu Li, Chao-Ji Zheng, Xiao-Dong He

## Match Overview

1	Internet 48 words crawled on 11-Oct-2020 <a href="http://www.hindawi.com">www.hindawi.com</a>	1%
2	Internet 30 words crawled on 19-Mar-2021 <a href="http://www.cureus.com">www.cureus.com</a>	1%
3	Internet 20 words crawled on 07-Oct-2020 <a href="http://fpublishing.bibcore.windows.net">fpublishing.bibcore.windows.net</a>	1%
4	Internet 17 words crawled on 05-Jul-2020 <a href="http://www.mysciencework.com">www.mysciencework.com</a>	<1%
5	Crossref 15 words Xin Wu, Binglu Li, Chaoji Zheng, Wei Liu, Tao Hong, Xiaodong He. "Risk Factors for Lateral Lymph Node Metastase ...	<1%
6	Internet 15 words crawled on 02-May-2016	<1%

国内版 国际版

Risk factors for preoperative carcinogenesis of bile duct cysts in adu



ALL

IMAGES

VIDEOS

61,800 Results

Any time ▾

People with these abnormalities are at higher risk of **cholangiocarcinoma**. Choledochal cysts are bile-filled sacs that are connected to the bile duct. The cells lining the sac often have areas of pre-cancerous growth, which can increase the risk for developing cholangiocarcinoma. It is a rare congenital abnormality.

[Risk Factors - Cholangiocarcinoma Foundation](https://cholangiocarcinoma.org/risk-factors/)  
[cholangiocarcinoma.org/risk-factors/](https://cholangiocarcinoma.org/risk-factors/)

Was this helpful?

#### PEOPLE ALSO ASK

What is Congenital biliary cyst? ▾

Can obesity cause cholangiocarcinoma? ▾

Is there an increase in biliary tract cancer? ▾

What is bile duct sclerosis? ▾

Feedback

#### Bile Duct Cyst as Precursor to Biliary Tract Cancer

<https://pubmed.ncbi.nlm.nih.gov/17187167>

Background: Bile duct cysts (BDC) are rare, of uncertain origin, and occur most often in young females of Asian descent. Increasingly, BDCs are reported in the Western population, often with coexistent biliary tract cancer. Methods: The PubMed and Medline literature databases were searched for pertinent publications regarding the clinical association and molecular biological development of ...

Cited by: 80

Author: Kjetil Søreide, Jon Arne Søreide, Jon Arn...

Publish Year: 2007

#### The risk of carcinogenesis in congenital choledochal cyst ...

<https://www.sciencedirect.com/science/article/pii/S1665268119309858>

May 01, 2014 - Bile atresia due to stenosis of the intrahepatic bile duct and associated cholangitis with