

检测到您输入了英文，试试切换到国际版？搜英文结果更丰富更准确 >

5,410,000 条结果 时间不限 ▾

COVID-19 and non-alcoholic fatty liver disease: Two intersecting pandemics **NAFLD** can play a role in the outcome of COVID-19 illness due to frequent association with comorbidities. Initial evidences suggest that increased liver fibrosis in NAFLD might affect COVID-19 outcome.

Author: Piero Portincasa, Marcin Krawczyk, Marcin Krawczyk, Wiktor Smyk, Frank Lammert, Agostino Di Ciaula

Cited by: 27

Publish Year: 2020

COVID-19 and non-alcoholic fatty liver disease: Two ...

 pubmed.ncbi.nlm.nih.gov/32589264/

这是否有帮助？  

Name of Journal: *World Journal of Hepatology*

Manuscript NO: 68617

Manuscript Type: OPINION REVIEW

COVID-19 and non-alcoholic fatty liver disease

Bellanti F *et al.* COVID-19 and NAFLD

Match Overview

1	Internet 22 words crawled on 06-Apr-2021 www.seac.gov	1%
2	Crossref 18 words Quentin M. Anstee, Christopher P. Day, "Epidemiology, Natural History, and Evaluation of Nonalcoholic Fatty Liver	1%
3	Internet 16 words crawled on 26-Jul-2021 www.medrxiv.org	<1%
4	Crossref 15 words Feng Gao, Kenneth I. Zheng, Hua-Dong Yan, Qing-Feng Sun et al. "Association and Interaction Between Serum Int	<1%
5	Internet 15 words crawled on 07-Jan-2016 www.arthritis-research.com	<1%
6	Crossref 13 words M. Nasirul Rahman, Md. Golam Kibria, Nigar Sultana, M. ahfuza Akhter et al. " Seroprevalence of and its associat	<1%
7	Internet 12 words crawled on 22-Jan-2021 biomarkers.biomedcentral.com	<1%
8	Internet 12 words crawled on 27-Oct-2020 www.emboexpress.org	<1%

国内版

国际版

COVID-19 and non-alcoholic fatty liver disease



ALL

IMAGES

VIDEOS

1,710,000 Results

Any time ▾

COVID-19 and non-alcoholic fatty liver disease: Two ...

<https://pubmed.ncbi.nlm.nih.gov/32589264>

Initial evidences suggest that increased liver fibrosis in NAFLD might affect COVID-19 outcome. In addition, long-term monitoring of post-COVID-19 NAFLD patients is advisable, to document further ...

COVID-19 and non-alcoholic fatty liver disease: Two intersecting pandemics

Cited by: 29

Author: Piero Portincasa, Marcin Krawczyk, Marc...

Publish Year: 2020

COVID-19 and Nonalcoholic Fatty Liver Disease - Full Text ...

<https://clinicaltrials.gov/ct2/show/NCT04982328> ▾

Jul 29, 2021 · COVID-19 is currently the leading public health problem, associated with a high risk of complications and death in risk groups of patients. Non-alcoholic fatty liver disease (NAFLD) is the most common liver disease with a prevalence of 30% in the Western population and is also recognized as an independent risk factor for the development of severe COVID-19.

[PDF] COVID-19 and Nonalcoholic Fatty Liver Disease

<https://www.aasld.org/sites/default/files/2020-10/COVID19-Flyer-NAFLD.pdf>

you understand coronavirus disease 2019 (COVID-19) infection and prevention in people with nonalcoholic fatty liver disease (NAFLD), including those with the more serious form of fatty liver disease, nonalcoholic steatohepatitis (NASH). What We Know Our understanding of COVID-19 in people with NAFLD is evolving. When making decisions about COVID-19

Nonalcoholic fatty liver disease is associated with COVID ...

<https://pubmed.ncbi.nlm.nih.gov/32868652>

Aim: Coronavirus disease 2019 (COVID-19) is a recently encountered disease that was declared a pandemic by WHO in 2020. Obesity and other components of the metabolic syndrome may aggravate the severity of COVID-19. Nonalcoholic fatty liver disease (NAFLD) represents the hepatic manifestation of metabolic syndrome.

Cited by: 13

Author: Mahmud Mahamid, Mahmud Mahamid, ...

Publish Year: 2020

Coronavirus Disease-2019 (COVID-19) and the Liver

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8111008>