

World Journal of *Gastrointestinal Pathophysiology*

World J Gastrointest Pathophysiol 2022 July 22; 13(4): 114-127



ORIGINAL ARTICLE

Case Control Study

- 114 Common polymorphisms of protein tyrosine phosphate non-receptor type 2 gene are not associated with risk of Crohn's disease in Indian

Chatterjee K, Dutta AK, Goel A, Aaron R, Balakrishnan V, Thomas A, John A, Jaleel R, David D, Kurien RT, Chowdhury S, Simon EG, Joseph A, Premkumar P, Pulimood AB

LETTER TO THE EDITOR

- 124 Hepatomusculoskeletal disorders: Coining a new term might improve the management of the musculoskeletal manifestations of chronic liver disease

Tsagkaris C, Papadakis SP, Moysidis DV, Papazoglou AS, Koutsogianni A, Papadakis M

ABOUT COVER

Associate Editor of *World Journal of Gastrointestinal Pathophysiology*, Nalu Navarro-Alvarez, MD, PhD, Assistant Professor, Department of Gastroenterology, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City 14080, Mexico. nalu.navarroa@incmnsz.mx

AIMS AND SCOPE

The primary aim of the *World Journal of Gastrointestinal Pathophysiology* (WJGP, *World J Gastrointest Pathophysiol*) is to provide scholars and readers from various fields of gastrointestinal pathophysiology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGP mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal pathophysiology and covering a wide range of topics including disorders of the esophagus, stomach and duodenum, small intestines, pancreas, biliary system, and liver.

INDEXING/ABSTRACTING

The WJGP is now abstracted and indexed in PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yi-Xuan Cai, Production Department Director: Xu Guo, Editorial Office Director: Jia-Ping Yan.

NAME OF JOURNAL

World Journal of Gastrointestinal Pathophysiology

ISSN

ISSN 2150-5330 (online)

LAUNCH DATE

April 15, 2010

FREQUENCY

Bimonthly

EDITORS-IN-CHIEF

Somchai Amornyotin, Tsutomu Nishida, Kusum K Kharbanda

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2150-5330/editorialboard.htm>

PUBLICATION DATE

July 22, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/gerinfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/gerinfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/gerinfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Hepatomusculoskeletal disorders: Coining a new term might improve the management of the musculoskeletal manifestations of chronic liver disease

Christos Tsagkaris, Stavros P Papadakos, Dimitrios V Moysidis, Andreas S Papazoglou, Alexandra Koutsogianni, Marios Papadakis

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C, C, C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Caballero-Mateos AM, Spain; Kim DJ, South Korea; Rodrigues AT, Brazil; Wu SZ, China

Received: December 25, 2021

Peer-review started: December 25, 2021

First decision: February 15, 2022

Revised: April 7, 2022

Accepted: June 26, 2022

Article in press: June 26, 2022

Published online: July 22, 2022



Christos Tsagkaris, Public Health and Policy Working Group, Stg European Student Think Tank, Amsterdam, Netherlands

Stavros P Papadakos, Alexandra Koutsogianni, Laiko General Hospital of Athens, National and Kapodistrian University of Athens, Athens 18233, Greece

Dimitrios V Moysidis, Hippokration University Hospital, Aristotle University of Thessaloniki, Thessaloniki 54642, Greece

Andreas S Papazoglou, Athens Naval Hospital, Athens 18233, Greece

Marios Papadakis, Department of Surgery II, University Hospital Witten-Herdecke, University of Witten-Herdecke, Wuppertal 42283, Germany

Corresponding author: Christos Tsagkaris, MD, Academic Fellow, Public Health and Policy Working Group, Stg European Student Think Tank, Postjeskade 29, 1058 DE, Amsterdam, Netherlands. chriss20x@gmail.com

Abstract

Chronic liver disease can affect many body systems including the musculoskeletal system. The pathogenetic crosstalk between the liver and organs such as the brain and the kidneys has already been described with compound terms merging the organs affected by the pathology, such as the hepatorenal syndrome. Nevertheless, the musculoskeletal manifestations of chronic liver disease have not been coined with such a term to date. Because of this shortage, documenting the musculoskeletal implications of chronic liver disease in both research and clinical practice is challenging. To fill this gap, the authors propose the term hepatomusculoskeletal disorders, a compound term of Greek origin that encompasses all the body structures involved in the aforementioned pathologic crosstalk.

Key Words: Chronic liver disease; Hepatomusculoskeletal disorders; Musculoskeletal system; Hepatology; Pathophysiology; Osteodystrophy

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The authors recommend coining the umbrella term “hepatomusculoskeletal disorders” in response to the need to expand knowledge about chronic liver disorders and capitalize it in the form of practice guidelines.

Citation: Tsagkaris C, Papadakis SP, Moysidis DV, Papazoglou AS, Koutsogianni A, Papadakis M. Hepatomusculoskeletal disorders: Coining a new term might improve the management of the musculoskeletal manifestations of chronic liver disease. *World J Gastrointest Pathophysiol* 2022; 13(4): 124-127

URL: <https://www.wjgnet.com/2150-5330/full/v13/i4/124.htm>

DOI: <https://dx.doi.org/10.4291/wjgp.v13.i4.124>

TO THE EDITOR

Chronic liver disease (CLD) is the 11th leading cause of mortality globally accounting for up to 2% of disability-adjusted life years worldwide[1]. It encompasses ailments of infectious (viral hepatitis) and non-infectious (alcohol abuse, non-alcoholic steatohepatitis, cancer) origin leading to progressive structural and functional depletion of hepatic physiology in the form of liver cirrhosis. CLD is associated with multisystem complications involving the kidneys, the heart, the nervous system and the musculoskeletal system[2]. Research in the field has recently sought hematological and electrocardiographic CLD biomarkers addressing CLD's extrahepatic manifestations as a potential standpoint for the management of the disease and for the identification of novel therapeutic targets[3-5]. Nevertheless, research regarding the musculoskeletal implications of CLD remains limited. Action is needed to expand the existing knowledge and its clinical applications.

The impact of CLD on the musculoskeletal system has been better understood during the last years [6]. The musculoskeletal manifestations can be classified into two categories according to their etiology: (1) On the causative disease which insults the liver; and (2) On the type and the degree of liver disease. In more detail, Hepatitis C is frequently associated with rheumatologic phenomena. Polyarthralgia either in the context of mixed cryoglobulinemia triad of purpura, fatigue and arthralgia or alone as hepatitis C virus (HCV)-induced arthritis is documented frequently[7,8]. Overt arthritis and fibromyalgia are less frequently diagnosed in parallel with HCV infection. Polyarthritides and polyarthralgia are commonly presented as manifestations of hepatitis B virus, hepatitis A virus and hepatitis E virus infections[7] while erosive arthritis is encountered in anti-cyclic citrullinated peptide positive type I autoimmune hepatitis[9]. In regards to the alcoholic liver disease, ethanol exerts direct cytotoxic effects into the muscular system causing alcoholic myopathy while affects bone metabolism causing matrix decomposition and suppression of bone synthesis[10]. Nonalcoholic fatty liver disease is frequently associated with low bone mineral density[11] while in diseases characterized by defective metabolism of metals (*e.g.*, copper in Wilson's disease and iron in haemochromatosis), arthritis, chondrocalcinosis and muscle stiffness and pain are regularly noticed[7,12]. On the other side, the severity of liver disease impacts the musculoskeletal health. Alterations in endogenous steroid metabolism and the use of proton pump inhibitors and diuretics results in fluctuations of mineral metabolism which result in hepatic osteodystrophy[13]. The defective immune responses due to poor complement system and opsonization sufficiency, portosystemic shunt and bacterial intestinal overgrowth render the patients prone to infections like septic arthritis, osteomyelitis, cellulitis and necrotizing fasciitis[14]. Finally, sarcopenia [15], non-traumatic osteonecrosis[16] and a higher rate of periprosthetic complications[17] are manifestations from the musculoskeletal system that compromise severely the quality of a patient's life.

On these grounds, healthcare professionals specializing in the management of musculoskeletal conditions (rheumatologists, orthopedic surgeons, physiatrists, physiotherapists, *etc*) can substantially contribute to CLD management. Prevention-wise, patients with CLD history can benefit from regular screening for osteopenia and osteoporosis and from falls' prevention training[18]. Similarly, physiotherapy to maintain muscle mass, improve patients' functionality and prevent sarcopenia-associated injury and disability can also be provided[19]. Treatment-wise, orthopedists and rheumatologists need to be aware of septic arthritis in CLD patients presenting with joint pain, and for spondylodiscitis and vertebral tuberculosis - in regions where the disease is endemic-in CLD patients presenting with low back pain[20-22]. Performing orthopedic surgery should also entail special considerations in CLD patients. Given their 3.5-fold higher risk for periprosthetic infections, cellulitis and necrotizing fasciitis, conservative management of fractures or osteoarthritis can be prioritized. In case of surgery, the patients and their formal and informal caregivers need to be instructed about the risk of infection and the need to carefully inspect surgical wounds and areas of plaster casting and seek medical attention when appropriate[23].

To contribute towards this end, musculoskeletal healthcare professionals need updated practice guidelines and relevant training. Developing concrete guidelines in turn requires systematic research in the field, with large scale observational studies and clinical trials confirming the existing knowledge and

optimizing the recommended interventions. Currently, it appears that research in the field is heterogeneous, with the majority of studies being observational and having been conducted independently in inconsistent time intervals.

A search for relevant publications on Medline, Scopus and other databases reveals a plethora of terms used to describe CLD musculoskeletal implications. The wording is often alternating (musculoskeletal disorders in patients with CLD, hepatic osteodystrophy) and rather descriptive words addressing particular alterations associated with CLD (sarcopenia, osteosarcopenia, skeletal muscle mass) rather than the phenomenon as a whole[2,24-26]. A term grouping all of the aforementioned together has not been included in the Medical Subject Headings thesaurus and in the International Disease Classification (ICD10) system to date. To the best of the authors' knowledge, no relevant term can be found in hospital records and documentation systems as well. Therefore, the lack of a consistent nomenclature poses significant obstacles to the appraisal of the existing knowledge, let alone its expansion.

The authors recommend coining the umbrella term "hepatomusculoskeletal disorders" in response to the need to expand relevant knowledge and capitalize it in the form of practice guidelines. The term is a compound word of Greek origin. It emphasizes the implications of liver conditions (hepato-) on muscles (musculo-), bones and connective tissue (skeletal). The composition of the term is similar to other relevant clinical terms such as the hepatorenal or the cardiorenal syndrome. In both these examples, the organs whose pathologies affect each other (liver, heart and kidneys respectively) are merged in a single term. Coining the new term in a similar linguistic format to other terms that are established in clinical practice makes it easily comprehensible to physicians and researchers. Therefore, the proposed term can benefit future research, clinical practice and medical education. Certainly, to address the musculoskeletal implications of CLD sufficiently, several steps involving clinicians, researchers, health bodies, healthcare administrators and stakeholders are required. Nonetheless, the new term can hopefully serve as common ground underlining the need to take relevant action.

FOOTNOTES

Author contributions: Tsagkaris C conceptualized this letter; Tsagkaris C, Papadakis SP and Moysidis DV performed the literature search; Tsagkaris C, Papadakis SP and Papazoglou AS wrote the first draft; Moysidis DV, Koutsogianni A and Papadakis M revised the manuscript and wrote the second draft; All authors have further revised, read and approve the final manuscript.

Conflict-of-interest statement: All authors report no relevant conflict of interest for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Switzerland

ORCID number: Christos Tsagkaris 0000-0002-4250-574X; Stavros P Papadakis 0000-0003-1583-1125; Dimitrios V Moysidis 0000-0001-9083-0267; Andreas S Papazoglou 0000-0003-4981-8121; Alexandra Koutsogianni 0000-0002-4215-3078; Marios Papadakis 0000-0002-9020-874X.

S-Editor: Wu YXJ

L-Editor: Filipodia

P-Editor: Wu YXJ

REFERENCES

- 1 Cheemerla S, Balakrishnan M. Global Epidemiology of Chronic Liver Disease. *Clin Liver Dis (Hoboken)* 2021; **17**: 365-370 [PMID: 34136143 DOI: 10.1002/cld.1061]
- 2 Ranjan R, Rampal S, Jaiman A, Tokgözü MA, Koong JK, Ramayah K, Rajaram R. Common musculoskeletal disorders in chronic liver disease patients. *Jt Dis Relat Surg* 2021; **32**: 818-823 [PMID: 34842121 DOI: 10.52312/jdrs.2021.25]
- 3 Surana P, Hercun J, Takyar V, Kleiner DE, Heller T, Koh C. Platelet count as a screening tool for compensated cirrhosis in chronic viral hepatitis. *World J Gastrointest Pathophysiol* 2021; **12**: 40-50 [PMID: 34084591 DOI: 10.4291/wjgp.v12.i3.40]
- 4 Rajapaksha IG, Angus PW, Herath CB. Current therapies and novel approaches for biliary diseases. *World J Gastrointest Pathophysiol* 2019; **10**: 1-10 [PMID: 30622832 DOI: 10.4291/wjgp.v10.i1.1]
- 5 Tsiompanidis E, Siakavellas SI, Tentolouris A, Eleftheriadou I, Chorepsima S, Manolakis A, Oikonomou K, Tentolouris N. Liver cirrhosis-effect on QT interval and cardiac autonomic nervous system activity. *World J Gastrointest Pathophysiol* 2018; **9**: 28-36 [PMID: 29487764 DOI: 10.4291/wjgp.v9.i1.28]

- 6 **Arora A**, Rajesh S, Bansal K, Sureka B, Patidar Y, Thapar S, Mukund A. Cirrhosis-related musculoskeletal disease: radiological review. *Br J Radiol* 2016; **89**: 20150450 [PMID: [27356209](#) DOI: [10.1259/bjr.20150450](#)]
- 7 **Fernandes B**, Dias E, Mascarenhas-Saraiva M, Bernardes M, Costa L, Cardoso H, Macedo G. Rheumatologic manifestations of hepatic diseases. *Ann Gastroenterol* 2019; **32**: 352-360 [PMID: [31263357](#) DOI: [10.20524/aog.2019.0386](#)]
- 8 **Zignego AL**, Gragnani L, Giannini C, Laffi G. The hepatitis C virus infection as a systemic disease. *Intern Emerg Med* 2012; **7** Suppl 3: S201-S208 [PMID: [23073858](#) DOI: [10.1007/s11739-012-0825-6](#)]
- 9 **Fusconi M**, Vannini A, Dall'Aglio AC, Pappas G, Cassani F, Ballardini G, Frisoni M, Grassi A, Bianchi FB, Zauli D. Anti-cyclic citrullinated peptide antibodies in type 1 autoimmune hepatitis. *Aliment Pharmacol Ther* 2005; **22**: 951-955 [PMID: [16268969](#) DOI: [10.1111/j.1365-2036.2005.02686.x](#)]
- 10 **González-Reimers E**, Quintero-Platt G, Rodríguez-Rodríguez E, Martínez-Riera A, Alvisa-Negrín J, Santolaria-Fernández F. Bone changes in alcoholic liver disease. *World J Hepatol* 2015; **7**: 1258-1264 [PMID: [26019741](#) DOI: [10.4254/wjh.v7.i9.1258](#)]
- 11 **Eshraghian A**. Bone metabolism in non-alcoholic fatty liver disease: vitamin D status and bone mineral density. *Minerva Endocrinol* 2017; **42**: 164-172 [PMID: [27973461](#) DOI: [10.23736/S0391-1977.16.02587-6](#)]
- 12 **Martinon F**, Pétrilli V, Mayor A, Tardivel A, Tschopp J. Gout-associated uric acid crystals activate the NALP3 inflammasome. *Nature* 2006; **440**: 237-241 [PMID: [16407889](#) DOI: [10.1038/nature04516](#)]
- 13 **Nakchbandi IA**, van der Merwe SW. Current understanding of osteoporosis associated with liver disease. *Nat Rev Gastroenterol Hepatol* 2009; **6**: 660-670 [PMID: [19881518](#) DOI: [10.1038/nrgastro.2009.166](#)]
- 14 **Park CH**, Joo YE, Choi SK, Rew JS, Kim SJ. Klebsiella pneumoniae septic arthritis in a cirrhotic patient with hepatocellular carcinoma. *J Korean Med Sci* 2004; **19**: 608-610 [PMID: [15308857](#) DOI: [10.3346/jkms.2004.19.4.608](#)]
- 15 **Kalafateli M**, Mantzoukis K, Choi Yau Y, Mohammad AO, Arora S, Rodrigues S, de Vos M, Papadimitriou K, Thorburn D, O'Beirne J, Patch D, Pinzani M, Morgan MY, Agarwal B, Yu D, Burroughs AK, Tsochatzis EA. Malnutrition and sarcopenia predict post-liver transplantation outcomes independently of the Model for End-stage Liver Disease score. *J Cachexia Sarcopenia Muscle* 2017; **8**: 113-121 [PMID: [27239424](#) DOI: [10.1002/jcsm.12095](#)]
- 16 **Deleuran T**, Overgaard S, Vilstrup H, Jepsen P. Cirrhosis is a risk factor for total hip arthroplasty for avascular necrosis. *Acta Orthop* 2016; **87**: 231-234 [PMID: [26900635](#) DOI: [10.3109/17453674.2016.1151122](#)]
- 17 **Salomon B**, Krause PC, Dasa V, Shi L, Jones D, Chapple AG. The Impact of Hepatitis C and Liver Disease on Risk of Complications After Total Hip and Knee Arthroplasty: Analysis of Administrative Data From Louisiana and Texas. *Arthroplast Today* 2021; **7**: 200-207 [PMID: [33553550](#) DOI: [10.1016/j.artd.2020.12.016](#)]
- 18 **Lara-Medrano R**, Alcázar-Quiñones C, Galarza-Delgado DÁ, Baena-Trejo L. Impact of a fall prevention program in the Internal Medicine wards of a tertiary care university hospital. *Med Univ* 2014; **16**: 156-160 Available from: <https://www.elsevier.es/enrevista-medicina-universitaria-304-articulo-impact-fall-prevention-program-in-X1665579614675994>
- 19 **Naseer M**, Turse EP, Syed A, Dailey FE, Zatreh M, Tahan V. Interventions to improve sarcopenia in cirrhosis: A systematic review. *World J Clin Cases* 2019; **7**: 156-170 [PMID: [30705893](#) DOI: [10.12998/wjcc.v7.i2.156](#)]
- 20 **Hung TH**, Hsieh MH, Lay CJ, Tsai CC. Increased occurrence of native septic arthritis in adult cirrhotic patients: a population-based three-year follow-up study in Taiwan. *Prz Gastroenterol* 2014; **9**: 342-347 [PMID: [25653729](#) DOI: [10.5114/pg.2014.47896](#)]
- 21 **Abdelrahman H**, Shousha M, Bahrami R, Boehm H. Haematogenous Spondylodiscitis in Patients With Liver Cirrhosis: Case Series of 36 Patients. *Spine (Phila Pa 1976)* 2020; **45**: E425-E429 [PMID: [31770341](#) DOI: [10.1097/BRS.0000000000003326](#)]
- 22 **Sharma D**, Kc S, Jaisi B. Prevalence of Tuberculosis in Patients with Liver Cirrhosis. *J Nepal Health Res Counc* 2018; **15**: 264-267 [PMID: [29353900](#) DOI: [10.3126/jnhrc.v15i3.18852](#)]
- 23 **Jiang SL**, Schairer WW, Bozic KJ. Increased rates of periprosthetic joint infection in patients with cirrhosis undergoing total joint arthroplasty. *Clin Orthop Relat Res* 2014; **472**: 2483-2491 [PMID: [24711129](#) DOI: [10.1007/s11999-014-3593-y](#)]
- 24 **Barbu EC**, Chițu-Tișu CE, Lazăr M, Olariu C, Bojincă M, Ionescu RA, Ion DA, Bădărău IA. Hepatic Osteodystrophy: A Global (Re)View of the Problem. *Acta Clin Croat* 2017; **56**: 512-525 [PMID: [29479918](#) DOI: [10.20471/acc.2017.56.03.19](#)]
- 25 **Saeki C**, Tsubota A. Influencing Factors and Molecular Pathogenesis of Sarcopenia and Osteosarcopenia in Chronic Liver Disease. *Life (Basel)* 2021; **11** [PMID: [34575048](#) DOI: [10.3390/Life11090899](#)]
- 26 **Tokuchi Y**, Suda G, Kimura M, Maehara O, Kitagataya T, Kubo A, Yoshida S, Fu Q, Yang Z, Hosoda S, Ohara M, Yamada R, Suzuki K, Kawagishi N, Nakai M, Sho T, Natsuizaka M, Morikawa K, Ogawa K, Ohnishi S, Sakamoto N. Possible correlation between increased serum free carnitine levels and increased skeletal muscle mass following HCV eradication by direct acting antivirals. *Sci Rep* 2021; **11**: 16616 [PMID: [34400736](#) DOI: [10.1038/s41598-021-96203-z](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

