

World Journal of *Clinical Cases*

World J Clin Cases 2023 May 26; 11(15): 3369-3663



REVIEW

- 3369** Superior mesenteric artery syndrome: Diagnosis and management
Oka A, Awoniyi M, Hasegawa N, Yoshida Y, Tobita H, Ishimura N, Ishihara S

MINIREVIEWS

- 3385** Astrocytes in the central nervous system and their functions in health and disease: A review
Gradisnik L, Velnar T
- 3395** Progress in diagnosis and treatment of acute injury to the anterior talofibular ligament
Chen RP, Wang QH, Li MY, Su XF, Wang DY, Liu XH, Li ZL
- 3408** Synchronous manifestation of colorectal cancer and intraductal papillary mucinous neoplasms
Mirchev MB, Boeva I, Peshevska-Sekulovska M, Stoitsov V, Peruhova M
- 3418** Clinical infections in neurosurgical oncology: An overview
Velnar T, Kocivnik N, Bosnjak R
- 3434** Effectiveness and safety of subthreshold vibration over suprathreshold vibration in treatment of muscle fatigue in elderly people
Mohamed AA, Khaled E, Hesham A, Khalf A

ORIGINAL ARTICLE

Clinical and Translational Research

- 3444** Establishment of a prognostic model related to tregs and natural killer cells infiltration in bladder cancer
Yang YJ, Xu XQ, Zhang YC, Hu PC, Yang WX

Retrospective Study

- 3457** New native tissue repair for pelvic organ prolapse: Medium-term outcomes of laparoscopic vaginal stump-round ligament fixation
Kakinuma T, Kaneko A, Kakinuma K, Imai K, Takeshima N, Ohwada M
- 3464** Demographic characteristics of patients who underwent anterior cruciate ligament reconstruction at a tertiary care hospital in India
Malv SK, Mahmood A, Vatsya P, Garika SS, Mittal R, Nagar M
- 3471** Usefulness of transcatheter arterial embolization for eighty-three patients with secondary postpartum hemorrhage: Focusing on difference in angiographic findings
Kim BM, Jeon GS, Choi MJ, Hong NS
- 3481** Chronic otitis media and middle ear variants: Is there relation?
Gökharman FD, Şenbil DC, Aydın S, Karavaş E, Özdemir Ö, Yalçın AG, Koşar PN

Observational Study

- 3491** Observation of the effect of angiojet to treat acute lower extremity arterial embolization
Meng XH, Xie XP, Liu YC, Huang CP, Wang LJ, Liu HY, Fang X, Zhang GH
- 3502** Outbreak of methanol-induced optic neuropathy in early COVID-19 era; effectiveness of erythropoietin and methylprednisolone therapy
Tabatabaei SA, Amini M, Haydar AA, Soleimani M, Cheraqpour K, Shahriari M, Hassanian-Moghaddam H, Zamani N, Akbari MR

META-ANALYSIS

- 3511** Impact of heart failure on outcomes in patients with sepsis: A systematic review and meta-analysis
Zhu MY, Tang XK, Gao Y, Xu JJ, Gong YQ

CASE REPORT

- 3522** New clinical application of digital intraoral scanning technology in occlusal reconstruction: A case report
Hou C, Zhu HZ, Xue B, Song HJ, Yang YB, Wang XX, Sun HQ
- 3533** Rare adult neuronal ceroid lipofuscinosis associated with *CLN6* gene mutations: A case report
Wang XQ, Chen CB, Zhao WJ, Fu GB, Zhai Y
- 3542** Enzyme replacement therapy in two patients with classic Fabry disease from the same family tree: Two case reports
Harigane Y, Morimoto I, Suzuki O, Temmoku J, Sakamoto T, Nakamura K, Machii K, Miyata M
- 3552** Immune-mediated necrotizing myopathy: Report of two cases
Chen BH, Zhu XM, Xie L, Hu HQ
- 3560** Retroperitoneal cavernous hemangioma misdiagnosed as lymphatic cyst: A case report and review of the literature
Hou XF, Zhao ZX, Liu LX, Zhang H
- 3571** Malignant melanoma resection and reconstruction with the first manifestation of lumbar metastasis: A case report
Guo ZX, Zhao XL, Zhao ZY, Zhu QY, Wang ZY, Xu M
- 3578** Promising way to address massive intragastric clotting in patients with acute upper gastrointestinal bleeding: A case report
Liu SX, Shi B, Liu YF, Shan JY, Sun B
- 3583** Pyogenic spondylitis caused by *Escherichia coli*: A case report and literature review
Zou LC, Qian J, Bian ZY, Wang XP, Xie T
- 3592** Primary ovarian choriocarcinoma occurring in a postmenopausal woman: A case report
Dai GL, Tang FR, Wang DQ

- 3599** Treatment of severe open bite and mandibular condyle anterior displacement by mini-screws and four second molars extraction: A case report
Huang ZW, Yang R, Gong C, Zhang CX, Wen J, Li H
- 3612** Application of apical negative pressure irrigation in the nonsurgical treatment of radicular cysts: A case report
Chen GP, Zhang YZ, Ling DH
- 3619** Treatment of postherpetic neuralgia by bone marrow aspirate injection: A case report
Honda Pazili T
- 3625** Non-target lung embolization during portal vein embolization due to an unrecognized portosystemic venous fistula: A case report
Alharbi SR, Bin Nasif M, Alwaily HB
- 3631** Acute abdomen caused by spontaneous rupture of degenerative hysteromyoma during pregnancy: A case report
Xu Y, Shen X, Pan XY, Gao S
- 3637** Atypical progress of frozen shoulder after COVID-19 vaccination: A case report
Jo HS, Kim HM, Han JY, Park HK
- 3643** Co-existing squamous cell carcinoma and chronic myelomonocytic leukemia with *ASXL1* and *EZH2* gene mutations: A case report
Deng LJ, Dong Y, Li MM, Sun CG
- 3651** Diagnosis based on electromagnetic navigational bronchoscopy-guided biopsied peripheral lung lesions in a 10-year-old girl: A case report
Meng FZ, Chen QH, Gao M, Zeng L, Lin JR, Zheng JY
- 3658** Relationship between intralobar pulmonary sequestration and type A aortic dissection: A case report
Wang YJ, Chen YY, Lin GH

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Gulali Aktas, MD, Professor, Department of Internal Medicine, Abant Izzet Baysal University Hospital, Bolu 14030, Turkey. draliaktas@yahoo.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*, Production Department Director: *Xiang Li*, Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

May 26, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Clinical and Translational Research

Establishment of a prognostic model related to tregs and natural killer cells infiltration in bladder cancer

Yan-Jie Yang, Xiao-Qing Xu, Yi-Chao Zhang, Peng-Cheng Hu, Wu-Xia Yang

Specialty type: Oncology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B, B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Cassell III AK, Liberia;
Muro M, Spain

Received: December 16, 2022

Peer-review started: December 16, 2022

First decision: February 28, 2023

Revised: March 8, 2023

Accepted: April 12, 2023

Article in press: April 12, 2023

Published online: May 26, 2023



Yan-Jie Yang, Shunde Hospital, Southern Medical University (The First People's Hospital of Shunde), Foshan 528308, Guangdong Province, China

Xiao-Qing Xu, The Graduate School, Tianjin Medical University, Tianjin 300041, China

Yi-Chao Zhang, The Graduate School, Qinghai University, Xi'ning 810000, Qinghai Province, China

Peng-Cheng Hu, Department of Ophthalmology, The Second Affiliated Hospital of Chongqing Medical University, Chongqing 400010, China

Wu-Xia Yang, The Graduate School/Department of Traditional Chinese Medicine, Tianjin Medical University/Tianjin Medical University General Hospital, Tianjin 300041, China

Corresponding author: Wu-Xia Yang, MM, Doctor, The Graduate School/Department of Traditional Chinese Medicine, Tianjin Medical University/Tianjin Medical University General Hospital, Tianjin 300041, China. yangwuxia_09@tmu.edu.cn

Abstract

BACKGROUND

Regulatory T cells (Tregs) and natural killer (NK) cells play an essential role in the development of bladder urothelial carcinoma (BUC).

AIM

To construct a prognosis-related model to judge the prognosis of patients with bladder cancer, meanwhile, predict the sensitivity of patients to chemotherapy and immunotherapy.

METHODS

Bladder cancer information data was obtained from The Cancer Genome Atlas and GSE32894. The CIBERSORT was used to calculate the immune score of each sample. Weighted gene co-expression network analysis was used to find genes that will have the same or similar expression patterns. Subsequently, multivariate cox regression and lasso regression was used to further screen prognosis-related genes. The prprphetic package was used to predict phenotype from gene expression data, drug sensitivity of external cell line and predict clinical data.

RESULTS

The stage and risk scores are independent prognostic factors in patients with

BUC. Mutations in *FGFR3* lead to an increase in Tregs percolation and affect the prognosis of the tumor, and additionally, *EMP1*, *TCHH* and *CNTNAP3B* in the model are mainly positively correlated with the expression of immune checkpoints, while *CMTM8*, *SORT1* and *IQSEC1* are negatively correlated with immune checkpoints and the high-risk group had higher sensitivity to chemotherapy drugs.

CONCLUSION

Prognosis-related models of bladder tumor patients, based on Treg and NK cell percolation in tumor tissue. In addition to judging the prognosis of patients with bladder cancer, it can also predict the sensitivity of patients to chemotherapy and immunotherapy. At the same time, patients were divided into high and low risk groups based on this model, and differences in genetic mutations were found between the high and low risk groups.

Key Words: Natural killer cells; Tregs; Bladder cancer; Weighted gene coexpression network analysis; Bladder cancer treatment; Immunotherapy; Computational molecular biology

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Tregs are thought to be connected to tumor cells evading the immune system, which is linked to cancer patients' poor prognoses. Natural killer (NK) cells control different immune responses and show antitumor cytotoxicity without being previously sensitized. We determined the immunological scores of several cell types in bladder urothelial carcinoma, discovered a gene set that was favorably connected with the Tregs score and negatively correlated with the NK cells score, and built a model that was related to prognosis. The model can predict the sensitivity of patients to chemotherapy and immunotherapy in addition to their prognosis for bladder cancers.

Citation: Yang YJ, Xu XQ, Zhang YC, Hu PC, Yang WX. Establishment of a prognostic model related to tregs and natural killer cells infiltration in bladder cancer. *World J Clin Cases* 2023; 11(15): 3444-3456

URL: <https://www.wjgnet.com/2307-8960/full/v11/i15/3444.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i15.3444>

INTRODUCTION

Bladder urothelial carcinoma (BUC) is a typical malignant growth of the urinary tract that has a high likelihood of return and death[1]. Immunotherapy has lately attracted more research attention than traditional surgery methods and chemotherapy[2]. Understanding the changes in the tumor microenvironment and finding potential immunotherapy strategies have received a lot of attention from scientific specialists. Thus, it is crucial to research how different immune cells infiltrate the tumor's inflammatory environment.

The start and progression of BUC are influenced by three processes: Immune monitoring, immunological balance, and immune escape. T cells (Tregs) are currently believed to be implicated in tumor cell immune evasion; Tregs are frequently found in tumors and are typically identified by the expression of CD4⁺ and Foxp3⁺, which is linked to the bad outcome of cancer patients[3]. The grade and stage of BUC were considerably correlated with the proportion of Foxp3⁺ Tregs[4]. Inhibiting Tregs can improve the treatment of BUC patients. Natural killer (NK) cells are mainly related to killing infected microorganisms and malignant transformed allogeneic and autologous cells. NK cells exhibit antitumor cytotoxicity without prior sensitization and production of cytokines and chemokines that regulate various immune responses[5]. NK cells patrol the body and are recruited to tumor sites to destroy malignant cells[6].

Based on the infiltration of Treg cells and NK cells into the tumor tissue, we built a prognosis-related model of bladder tumor patients for this investigation. It can predict the susceptibility of patients to chemotherapy and immunotherapy in addition to assessing the prognosis of bladder cancer patients.

MATERIALS AND METHODS

Downloading and processing of data

The validated database is derived from GSE32894. The platform of the data set is GPL6947. In the Gene Expression Omnibus (GEO) database, the end-point event is determined according to the disease-

specific survival. GSE32894 included 308 patients with BUC. The data source of the training data set is the transcriptome data of The Cancer Genome Atlas (TCGA's) bladder cancer, including 19 patients' normal samples and 411 patients' cancer samples. The Data Category and Workflow Type of TCGA data are transcriptome profiling and Fragments per Kilobase Million (FPKM). Then, the TranscriptsPerKilobase of exonmodel per Million mapped reads (TPM) method was used to standardize the TCGA data for model verification. The mutation data were obtained from the simple nucleoside variation data of 409 patients with bladder urothelial carcinoma in TCGA database.

Calculation of immune cell score

Each sample's immunological score was determined using "CIBERSORT". 22 immune cells in total were scored, and "perm = 1000, qn = true" was specified as the option in "CIBERSORT".

Weighted gene coexpression network analysis

Finding genes that will have the same or comparable expression patterns may be done using the useful method known as weighted gene coexpression network analysis (WGCNA). These genes could have comparable physiological effects. WGCNA differs from other straightforward clustering methods in this way (such as based on Euclidean distance). It is a clustering technique with biologically relevant applications. In this work, we employed WGCNA to categorize the bladder cancer expression profile data into different groups.

Screening of prognostic related genes

Then, the prognosis-related diagnostic genes in the TCGA database were screened using the univariate cox regression model. Multivariate cox regression and lasso regression were utilized to further screen prognosis-related genes based on the findings of the univariate cox screening of the TCGA database and GEO data. We determined the riskcore for each sample using the multivariate Cox model (riskcore = gene expression * coef).

Drug sensitivity analysis

Prrophetic package is an ancient R package. Its main purpose is to predict phenotype from gene expression data (predict clinical results using CGP cell line data of cancer genome project), predict drug sensitivity of external cell line, and predict clinical data.

Statistical methods

The clinical data were compared using the chi square test. The correlation between different values was determined using Pearson correlation analysis. The following is the study's flowchart (Figure 1).

Waterfall diagram of gene mutation

Maftools package is used to analyze the overall situation of gene mutation between high-risk and low-risk groups. The top 20 genes with the highest frequency of mutations in both high-risk and low-risk groups are displayed in a waterfall chart.

The Cancer Immunome Atlas

The Cancer Immunome Atlas (TCIA) is developed based on TCGA data. The difference is that TCIA only provides immune data analysis of 20 cancer species. The immunophenoscore (IPS) of each sample in the high and low risk categories, as well as the difference between them, were determined using TCIA. The reactivity of CTLA-4 and PD-1 can be accurately predicted by IPS.

RESULTS

Immune cell score and WGCNA

In the TCGA database, we first assigned a number to each sample's immune cells (Figure 2A). The total genes were divided into 9 gene sets based on the immune cell scores of each sample using the WGCNA technique, and the correlation between these 9 gene sets and the immune cell scores of each type was determined (Figure 2B-E). Then we chose a group of genes (module yellow) that had a positive correlation with the score for Regulatory T cells (Treg) ($r = 0.45$) and a negative correlation with the score for NK cell activation ($r = -0.33$). According to earlier studies, Tregs inhibit long-lasting immune reactions to viruses, tumors, and self-antigens; the infiltrate tumor tissue and are linked to a bad prognosis in cancer patients; the quantity of Treg was negatively correlated with recurrence-free survival in BUC and substantially increased in peripheral blood and tumor tissue[7]. The grade and stage of BUC were significantly associated with the proportion of Foxp3+ Tregs[8]. The most significant component of natural immunity is played by NK cells, which can directly destroy target cells like cancerous or virus-infected ones. They serve as the first line of protection against tumors and the first obstacle of human antibodies. From this perspective, the gene collection we chose is also consistent with




Figure 1 The flowchart of our study. TCGA: The Cancer Genome Atlas; NK: Natural killer; WGCNA: Weighted gene coexpression network analysis.

the findings of earlier studies. More significantly, prior research has shown that tumor-infiltrating Treg cells can cause NK cells, B cells, Dendritic cells (DC) and cytotoxic T cells in the Tumor microenvironment (TME) to undergo death[9]. This strengthens the logic of our study concepts even more.

Screening of prognosis related genes and construction of model


By using multivariate Cox regression analysis, the prognostic-related genes in the yellow gene group were chosen (Figure 3A). The alleles with high association were then eliminated using lasso regression (Figure 3B). Lastly, the prognostic association model was built using multivariate Cox regression. 13 genes in total (*CHAT*, *EMP1*, *ADSL*, *TRIM16*, *CMTM8*, *DNASE2B*, *NTN4*, *SORT1*, *IQSEC2*, *TCHH*, *CNTNAP3B*, *ESD*, *NPSR1*) were used to build this model (Figure 4A and B). The riskscore of each sample was then determined using the multivariate Cox model (riskscore = gene expression * coef). Samples that scored above the median for hazards were placed in the high-risk category, while those that scored below the median were placed in the low-risk group. Then it was looked at how long the high-risk group survived compared to the low-risk group. Clearly, the prognosis between the high-risk group and the low-risk group is extremely different (Figure 3C). In comparison to the high-risk group, the prognosis for the low-risk group was noticeably better. The GEO database has verified the model's efficacy in determining prognosis (Figure 4D). In the supporting materials, a survival analysis of individual genes will be presented. Furthermore, we determined the association ($r > 0.4$ and P value > 0.05 were deemed significant) between each gene's expression and immune cell infiltration in the model. The prognosis of patients was then predicted using Nomo map (Figure 2E; stage: "1", "2", "3", "4" represent Stage I, Stage II, Stage III, and Stage IV, respectively; grade: "1" and "2" represent G1 and G2, respectively; gender: "0" and "1" represent females and males, respectively).

Model independence and diagnostic efficiency

Cox regression analysis was conducted to determine whether our model could predict the prognosis of BUC patients independently of other clinical data. Stage and risk score were shown to be strongly correlated with the prognosis of BUC patients by univariate Cox regression analysis. Stage and risk score were identified as independent predictive variables in multivariate Cox regression analysis of BUC patients (Figure 4E and F). Furthermore, the scatter plot demonstrates that patients who scored highly for hazards have much poorer prognoses (Figure 4C and D). The receiver operating characteristic curve (AUC at 1 year: 0.699, AUC at 3 years: 0.745, AUC at 5 years: 0.753) further proves that the model can more accurately predict the prognosis of patients (Figure 4G). We employed chi-square test to evaluate the difference in clinical data between high and low risk groups ("***" = $P < 0.001$, "**" = $P < 0.01$, "*" = $P < 0.05$) in order to further investigate the difference between high and low risk groups. Patients in the high-risk group underwent much more distant metastases and had significantly later clinical stages. (Figure 4H).

Differences in genetic mutations between high and low risk

The variations in genetic mutations between high and low risk were then looked at. According to the findings, the prevalence of *FGFR3* mutations varied considerably between the high-risk and low-risk groups (Figure 5A and B). One of the frequent mutations in bladder cancer is the *FGFR3* mutation, which is also one of the key elements in carcinogenesis. Moreover, *FGFR3*-targeting medications have been utilized to treat BUC patients. It will be fascinating to explore if the *FGFR3* mutation would boost Treg infiltration and have an impact on the prognosis of tumors.





DOI: 10.12998/wjcc.v11.i15.3444 Copyright ©The Author(s) 2023.

Figure 2 Immune infiltration status of TCGA database samples and gene sets related to immune cell infiltration. A: Heatmap of the distribution of immune cell infiltration in each sample; B: The scale independence and the mean connectivity; C: Hierarchical clustering was used to obtain coexpressive gene modules; Different gene modules are assigned to different colors; D: Clustering diagram of each gene module; E: Clustering diagram of each gene module.

The difference of IPS (immunophenoscore) and gene mutation between high-risk and low-risk groups, as well as the difference between the model and each immune checkpoint

The model and each immune checkpoint differ in that *EMP1*, *TCHH*, and *CNTNAP3B* are mostly favorably connected with the expression of immunological checkpoints, whereas *CMTM8*, *SORT1*, and *IQSEC1* are adversely linked with immune checkpoints (Figure 5C). Moreover, the IPS between the





DOI: 10.12998/wjcc.v11.i15.3444 Copyright ©The Author(s) 2023.

Figure 3 Screening of survival related gene sets (based on immune infiltration). A: Screening of prognostic genes by univariate Cox; B: Least absolute shrinkage and selection operator (LASSO) regression model; C: Kaplan Meier curve shows that in The Cancer Immunome Atlas (TCGA) database, the prognosis of patients between high and low risk groups is different (Training group); D: Kaplan Meier curve shows the difference in prognosis between high and low risk groups in geo database (Test group); E: Nomograms were used to measure patient survival in The Cancer Immunome Atlas database.

high-risk and low-risk groups varied (Figure 5D). It is evident that the immune checkpoint inhibition may work better for the low-risk population.


Prediction of drug sensitivity

To distinguish between high and low risk groups in terms of medication sensitivity, we employed the prophetic package (R package). Interestingly, chemotherapeutic medication sensitivity was greater in the high-risk group. These chemotherapeutic medications include vinblastine, cisplatin, docetaxel, mitomycin, and adriamycin (Figure 6). These are all chemotherapeutic medications that are frequently used to treat bladder cancer.

DISCUSSION

Tregs, an immunosuppressive subset of CD4⁺ T cells, play a crucial role in maintaining self-tolerance, avoiding autoimmune disorders and transplant rejection, it has been proposed as candidates of biomarkers because of their capability to control alloimmune responses[10]. In tumor immunity, Tregs compromise immune surveillance against cancer in healthy individuals and damage the antitumor immune response in tumor-bearing hosts[11]. Among different types of tumor infiltrating lymphocytes, the infiltration of cytotoxic CD8⁺T cells plays a positive role in the prognosis of tumor patients and the clinical response of immunotherapy[12,13], but the infiltration of tumor infiltrating lymphocytes called Treg commonly indicates poor prognosis and poor effect of immunotherapy. Tregs have been reported to increase infiltration in various tumor tissues, which is related to tumor immune escape.

Tregs can secrete anti-inflammatory mediators including Interleukin (IL)-10, Transforming factor β (TGF- β), and IL-35, which can suppress the immune system. TGF- β lowers the lethality of NK cells and induces them to transform into type 1 innate lymphocytes, which has further effects on tumor development and metastasis[14].



DOI: 10.12998/wjcc.v11.i15.3444 Copyright ©The Author(s) 2023.

Figure 4 Differences in prognosis between high-risk and low-risk patients. A: Heat map of gene expression differences between high and low risk groups in The Cancer Genome Atlas database; B: Heat map of gene expression differences between high and low risk groups in Gene Expression Omnibus database; C: The ROC curve of the risk score and clinicopathological variables; D: Distribution heat map of model genes and clinicopathological variables; E: The ROC curve of the risk score and clinicopathological variables; F: Distribution heat map of model genes and clinicopathological variables.




Figure 6 Differences in IC50 values of different drugs between high and low risk groups. A: IC50 values of cisplatin between high and low risk groups; B: IC50 values of Docetaxel between high and low risk groups; C: IC50 values of vinblastine between high and low risk groups; D: IC50 values of Doxorubicin between high and low risk groups; E: IC50 values of Mitomycin. C between high and low risk groups.

lines to epirubicin. Moreover, the expression of *CMTM8* has been reported to be decreased in a variety of tumors, such as liver cancer, colon, lung adenocarcinoma and so on. At the same time, *CMTM8* is also associated with EMT (epithelial mesenchymal transformation) and immune infiltration[25,26]. The conclusions drawn from this article contradict our results. *DNASE2B* may be associated with chemotherapy sensitivity of gastric cancer[27]. Nerve guidance factor 4 (Netrin-4, *NTN4*) is a ligand of neo1 (a transmembrane receptor). Its expression in gastric cancer, breast cancer and neuroblasts is related to the prognosis of tumor patients, and EMT is closely related[28-30]. *Sort1* is involved in the occurrence, development, and drug resistance of a variety of tumors, such as colon cancer, breast cancer and gastric cancer[31,32]. The expression of *TCHH* was correlated with immune infiltration in Colon cancer and this result is based on TIMER database analysis[33]. Very little research has been done on the relationship between the genes included in the model and the infiltration of immune cells in tumor tissue. In addition, the risk score calculated by this model is positively correlated with PDCD2LG2 (PD-L1) and Cytotoxic T lymphocyte-4, suggesting that the model may be helpful to guide the treatment of immune examination sites.

Taha-Mehlitz *et al*[24] study showed that the mutation and overexpression of *FGFR3* were related to the low immune score of bladder cancer. It has been reported that Treg cells are highly infiltrated into tumors, which may lead to immunosuppression of tumor microenvironment and increase tumor immune escape[9]. Based on our data and previous studies, we have obtained a hypothesis that the increase of *FGFR3* mutation will lead to the increase of Treg infiltration (because the mutation frequency of *FGFR3* is significantly increased in the high-risk group), which will lead to the immune escape of tumor cells and lead to the poor prognosis of patients. This hypothesis may be a good research idea. The prediction of drug susceptibility may be an unexpected finding of this study.

Based on our data analysis, this model related to Tregs and NK cells can predict the prognosis of BUC, correlate with clinical data, and predict the sensitivity of bladder patients to chemotherapeutic drugs. However, it is not clear whether the changes of these genes in tumor tissue are the cause or result of Tregs and NK cells participating in the formation of TME. Here we merely provide some reference data for future researchers.

We constructed a prognosis-related model for patients with bladder tumors. This model is based on the infiltration of Treg cells and NK cells in tumor tissue. In addition to judging the prognosis of bladder cancer patients, it can also predict how sensitive patients are to chemotherapy and immunotherapy. Furthermore, based on this model, patients were divided into high and low risk groups, and variations in genetic alterations were found between the high and low risk groups.

In summary, Treg cell and NK cell percolation in this model provides new targets for targeted therapy of BUC. This study does have certain restrictions, though. Because there are many types of bladder cancer, yet, other research aspects may be overlooked, due to the pursuit of "accuracy" in the study of differentially expressed genes. In follow-up studies, we would consider looking at other subtypes to determine if there are any changes and investigating other immune-related processes for prognosis-related factors.

The role and function of Tregs and NK cells need to be confirmed using more sophisticated methodologies and approaches as our analysis is based on the TCGA database, which is openly accessible but has been mined.

CONCLUSION

Prognosis-related models of bladder tumor patients, based on Treg and NK cell percolation in tumor tissue. In addition to judging the prognosis of patients with bladder cancer, it can also predict the sensitivity of patients to chemotherapy and immunotherapy. At the same time, patients were divided into high and low risk groups based on this model, and differences in genetic mutations were found between the high and low risk groups.

ARTICLE HIGHLIGHTS

Research background

Regulatory T cells (Tregs) and natural killer (NK) cells play an essential role in the development of bladder urothelial carcinoma (BUC).

Research motivation

To identify genes associated with bladder cancer prognosis.

Research objectives

To construct a prognosis-related model to judge the prognosis of patients with bladder cancer, meanwhile, predict the sensitivity of patients to chemotherapy and immunotherapy.

Research methods

In this study, we analyzed publicly available datasets from two databases (<https://portal.gdc.cancer.gov/> and <https://www.ncbi.nlm.nih.gov/geo/>), both with authoritative data, to predict the diagnosis and prognosis of the disease by bioinformatic analysis.

Research results

The stage and risk scores are independent prognostic factors in patients with BUC. Mutations in *FGFR3* lead to an increase in Tregs percolation and affect the prognosis of the tumor, and additionally, *EMP1*, *TCHH* and *CNTNAP3B* in the model are mainly positively correlated with the expression of immune checkpoints, while *CMTM8*, *SORT1* and *IQSEC1* are negatively correlated with immune checkpoints and the high-risk group had higher sensitivity to chemotherapy drugs.

Research conclusions

The model can predict the sensitivity of patients to chemotherapy and immunotherapy in addition to their prognosis for bladder cancers.

Research perspectives

Prognosis-related models of bladder tumor patients, based on Treg and NK cell percolation in tumor tissue. In addition to judging the prognosis of patients with bladder cancer, it can also predict the sensitivity of patients to chemotherapy and immunotherapy. At the same time, patients were divided into high and low risk groups based on this model, and differences in genetic mutations were found between the high and low risk groups.

FOOTNOTES

Author contributions: Yang YJ and Xu XQ designed and coordinated the study; Yang YJ performed the experiments, acquired, and analyzed data; Xu XQ, Zhang YC, and Hu PC interpreted the data; Yang YJ, Xu XQ, and Yang WX wrote the manuscript; all authors approved the final version of the article; Yang YJ and Xu XQ contributed equally to this work.

Institutional review board statement: Nor ethical approval nor informed consent was required in this study due to the public available of data in the WGCNA, GEO, and TCIA databases (<https://portal.gdc.cancer.gov/> and <https://www.ncbi.nlm.nih.gov/geo/>).

Conflict-of-interest statement: There is no conflict of interest between the authors.

Data sharing statement: All authors agree to data sharing.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Yan-Jie Yang 0009-0002-1617-4374; Xiao-Qing Xu 0000-0001-7577-6362; Yi-Chao Zhang 0009-0005-7433-6791; Peng-Cheng Hu 0000-0002-3720-2488; Wu-Xia Yang 0000-0002-7967-9399.

S-Editor: Liu JH

L-Editor: A

P-Editor: Cai YX

REFERENCES

- Martinez Rodriguez RH, Buisan Rueda O, Ibarz L. Bladder cancer: Present and future. *Med Clin (Barc)* 2017; **149**: 449-455 [PMID: 28736063 DOI: 10.1016/j.medcli.2017.06.009]
- Rouanne M, Bajorin DF, Hannan R, Galsky MD, Williams SB, Necchi A, Sharma P, Powles T. Rationale and Outcomes for Neoadjuvant Immunotherapy in Urothelial Carcinoma of the Bladder. *Eur Urol Oncol* 2020; **3**: 728-738 [PMID: 33177001 DOI: 10.1016/j.euo.2020.06.009]
- Toor SM, Murshed K, Al-Dhaheeri M, Khawar M, Abu Nada M, Elkord E. Immune Checkpoints in Circulating and Tumor-Infiltrating CD4(+) T Cell Subsets in Colorectal Cancer Patients. *Front Immunol* 2019; **10**: 2936 [PMID: 31921188 DOI: 10.3389/fimmu.2019.02936]
- Phé V, Rouprêt M, Cussenot O, Chartier-Kastler E, Gamé X, Compérat E. Forkhead box protein P3 (Foxp3) expression serves as an early chronic inflammation marker of squamous cell differentiation and aggressive pathology of urothelial carcinomas in neurological patients. *BJU Int* 2015; **115** Suppl 6: 28-32 [PMID: 25600959 DOI: 10.1111/bju.13044]
- Guillerey C. NK Cells in the Tumor Microenvironment. *Adv Exp Med Biol* 2020; **1273**: 69-90 [PMID: 33119876 DOI: 10.1007/978-3-030-49270-0_4]
- Shimasaki N, Jain A, Campana D. NK cells for cancer immunotherapy. *Nat Rev Drug Discov* 2020; **19**: 200-218 [PMID: 31907401 DOI: 10.1038/s41573-019-0052-1]
- Hakimi AA, Voss MH, Kuo F, Sanchez A, Liu M, Nixon BG, Vuong L, Ostrovskaya I, Chen YB, Reuter V, Riaz N, Cheng Y, Patel P, Marker M, Reising A, Li MO, Chan TA, Motzer RJ. Transcriptomic Profiling of the Tumor Microenvironment Reveals Distinct Subgroups of Clear Cell Renal Cell Cancer: Data from a Randomized Phase III Trial. *Cancer Discov* 2019; **9**: 510-525 [PMID: 30622105 DOI: 10.1158/2159-8290.CD-18-0957]
- Fattahi S, Karimi M, Ghatreh-Samani M, Taheri F, Shirzad H, Mohammad Alibeigi F, Anjomshoa M, Bagheri N. Correlation between aryl hydrocarbon receptor and IL-17(+) and Foxp3(+) T-cell infiltration in bladder cancer. *Int J Exp Pathol* 2021; **102**: 249-259 [PMID: 34762773 DOI: 10.1111/ieip.12392]
- Hatzioannou A, Boumpas A, Papadopolou M, Papafragkos I, Varveri A, Alissafi T, Verginis P. Regulatory T Cells in Autoimmunity and Cancer: A Duplicious Lifestyle. *Front Immunol* 2021; **12**: 731947 [PMID: 34539668 DOI: 10.3389/fimmu.2021.731947]
- San Segundo D, Millán O, Muñoz-Cacho P, Boix F, Paz-Artal E, Talayero P, Morales JM, Muro M, De Cos MÁ, Guirado L, Llorente S, Pascual J, Arias M, Brunet M, López-Hoyos M. High proportion of pretransplantation activated regulatory T cells (CD4+CD25highCD62L+CD45RO+) predicts acute rejection in kidney transplantation: results of a multicenter study. *Transplantation* 2014; **98**: 1213-1218 [PMID: 25083613 DOI: 10.1097/TP.0000000000000202]
- Nishikawa H, Koyama S. Mechanisms of regulatory T cell infiltration in tumors: implications for innovative immune precision therapies. *J Immunother Cancer* 2021; **9** [PMID: 34330764 DOI: 10.1136/jitc-2021-002591]
- Saito T, Nishikawa H, Wada H, Nagano Y, Sugiyama D, Atarashi K, Maeda Y, Hamaguchi M, Ohkura N, Sato E, Nagase H, Nishimura J, Yamamoto H, Takiguchi S, Tanoue T, Suda W, Morita H, Hattori M, Honda K, Mori M, Doki Y, Sakaguchi S. Two FOXP3(+)CD4(+) T cell subpopulations distinctly control the prognosis of colorectal cancers. *Nat Med* 2016; **22**: 679-684 [PMID: 27111280 DOI: 10.1038/nm.4086]
- Saleh R, Elkord E. FoxP3(+) T regulatory cells in cancer: Prognostic biomarkers and therapeutic targets. *Cancer Lett* 2020; **490**: 174-185 [PMID: 32721551 DOI: 10.1016/j.canlet.2020.07.022]
- Gao Y, Souza-Fonseca-Guimaraes F, Bald T, Ng SS, Young A, Ngio SF, Rautela J, Straube J, Waddell N, Blake SJ, Yan J, Bartholin L, Lee JS, Vivier E, Takeda K, Messaoudene M, Zitvogel L, Teng MWL, Belz GT, Engwerda CR, Huntington ND, Nakamura K, Hölzel M, Smyth MJ. Tumor immunoevasion by the conversion of effector NK cells into type 1 innate

- lymphoid cells. *Nat Immunol* 2017; **18**: 1004-1015 [PMID: [28759001](#) DOI: [10.1038/ni.3800](#)]
- 15 **Marofi F**, Abdul-Rasheed OF, Rahman HS, Budi HS, Jalil AT, Yumashev AV, Hassanzadeh A, Yazdanifar M, Motavalli R, Chartrand MS, Ahmadi M, Cid-Arreguid A, Jarahian M. CAR-NK cell in cancer immunotherapy; A promising frontier. *Cancer Sci* 2021; **112**: 3427-3436 [PMID: [34050690](#) DOI: [10.1111/cas.14993](#)]
- 16 **Guillamón CF**, Martínez-Sánchez MV, Gimeno L, Campillo JA, Server-Pastor G, Martínez-García J, Martínez-Escribano J, Torroba A, Ferri B, Abellán DJ, Legaz I, López-Álvarez MR, Moya-Quiles MR, Muro M, Minguela A. Activating KIRs on Educated NK Cells Support Downregulation of CD226 and Inefficient Tumor Immunosurveillance. *Cancer Immunol Res* 2019; **7**: 1307-1317 [PMID: [31239317](#) DOI: [10.1158/2326-6066.CIR-18-0847](#)]
- 17 **Demaria O**, Gauthier L, Debroas G, Vivier E. Natural killer cell engagers in cancer immunotherapy: Next generation of immuno-oncology treatments. *Eur J Immunol* 2021; **51**: 1934-1942 [PMID: [34145579](#) DOI: [10.1002/eji.202048953](#)]
- 18 **Barry KC**, Hsu J, Broz ML, Cueto FJ, Binnewies M, Combes AJ, Nelson AE, Loo K, Kumar R, Rosenblum MD, Alvarado MD, Wolf DM, Bogunovic D, Bhardwaj N, Daud AI, Ha PK, Ryan WR, Pollack JL, Samad B, Asthana S, Chan V, Krummel MF. A natural killer-dendritic cell axis defines checkpoint therapy-responsive tumor microenvironments. *Nat Med* 2018; **24**: 1178-1191 [PMID: [29942093](#) DOI: [10.1038/s41591-018-0085-8](#)]
- 19 **Böttcher JP**, Bonavita E, Chakravarty P, Blees H, Cabeza-Cabrero M, Sammiceli S, Rogers NC, Sahai E, Zelenay S, Reis e Sousa C. NK Cells Stimulate Recruitment of cDC1 into the Tumor Microenvironment Promoting Cancer Immune Control. *Cell* 2018; **172**: 1022-1037.e14 [PMID: [29429633](#) DOI: [10.1016/j.cell.2018.01.004](#)]
- 20 **Miao L**, Jiang Z, Wang J, Yang N, Qi Q, Zhou W, Feng Z, Li W, Zhang Q, Huang B, Chen A, Zhang D, Zhao P, Li X. Epithelial membrane protein 1 promotes glioblastoma progression through the PI3K/AKT/mTOR signaling pathway. *Oncol Rep* 2019; **42**: 605-614 [PMID: [31233190](#) DOI: [10.3892/or.2019.7204](#)]
- 21 **Lai S**, Wang G, Cao X, Li Z, Hu J, Wang J. EMP-1 promotes tumorigenesis of NSCLC through PI3K/AKT pathway. *J Huazhong Univ Sci Technolog Med Sci* 2012; **32**: 834-838 [PMID: [23271282](#) DOI: [10.1007/s11596-012-1043-1](#)]
- 22 **Lin B**, Zhang T, Ye X, Yang H. High expression of EMP1 predicts a poor prognosis and correlates with immune infiltrates in bladder urothelial carcinoma. *Oncol Lett* 2020; **20**: 2840-2854 [PMID: [32782602](#) DOI: [10.3892/ol.2020.11841](#)]
- 23 **Taha-Mehlitz S**, Bianco G, Coto-Llerena M, Kancherla V, Bantug GR, Gallon J, Ercan C, Panebianco F, Eppenberger-Castori S, von Strauss M, Staubli S, Bolli M, Peterli R, Matter MS, Terracciano LM, von Flüe M, Ng CKY, Soysal SD, Kollmar O, Piscuoglio S. Adenylosuccinate lyase is oncogenic in colorectal cancer by causing mitochondrial dysfunction and independent activation of NRF2 and mTOR-MYC-axis. *Theranostics* 2021; **11**: 4011-4029 [PMID: [33754045](#) DOI: [10.7150/thno.50051](#)]
- 24 **Gao D**, Hu H, Wang Y, Yu W, Zhou J, Wang X, Wang W, Zhou C, Xu K. CMTM8 inhibits the carcinogenesis and progression of bladder cancer. *Oncol Rep* 2015; **34**: 2853-2863 [PMID: [26503336](#) DOI: [10.3892/or.2015.4310](#)]
- 25 **Kang N**, Xie X, Zhou X, Wang Y, Chen S, Qi R, Liu T, Jiang H. Identification and validation of EMT-immune-related prognostic biomarkers CDKN2A, CMTM8 and ILK in colon cancer. *BMC Gastroenterol* 2022; **22**: 190 [PMID: [35429970](#) DOI: [10.1186/s12876-022-02257-2](#)]
- 26 **Zhang W**, Qi H, Mo X, Sun Q, Li T, Song Q, Xu K, Hu H, Ma D, Wang Y. CMTM8 is Frequently Downregulated in Multiple Solid Tumors. *Appl Immunohistochem Mol Morphol* 2017; **25**: 122-128 [PMID: [26574634](#) DOI: [10.1097/PAI.0000000000000274](#)]
- 27 **Ha YJ**, Yoon SN, Jeon YJ, Cho DH, Roh SA, Kim BS, Kim HJ, Kim SY, Kim YS, Kim JC. Genome-wide identification of chemosensitive single nucleotide polymorphism markers in gastric cancer. *Anticancer Res* 2011; **31**: 4329-4338 [PMID: [22199298](#)]
- 28 **Lv B**, Song C, Wu L, Zhang Q, Hou D, Chen P, Yu S, Wang Z, Chu Y, Zhang J, Yang D, Liu J. Netrin-4 as a biomarker promotes cell proliferation and invasion in gastric cancer. *Oncotarget* 2015; **6**: 9794-9806 [PMID: [25909166](#) DOI: [10.18632/oncotarget.3400](#)]
- 29 **Villanueva AA**, Falcón P, Espinoza N, R LS, Milla LA, Hernandez-SanMiguel E, Torres VA, Sanchez-Gomez P, Palma V. The Netrin-4/ Neogenin-1 axis promotes neuroblastoma cell survival and migration. *Oncotarget* 2017; **8**: 9767-9782 [PMID: [28038459](#) DOI: [10.18632/oncotarget.14213](#)]
- 30 **Xu X**, Yan Q, Wang Y, Dong X. NTN4 is associated with breast cancer metastasis via regulation of EMT-related biomarkers. *Oncol Rep* 2017; **37**: 449-457 [PMID: [27840993](#) DOI: [10.3892/or.2016.5239](#)]
- 31 **Demeule M**, Charfi C, Currie JC, Larocque A, Zgheib A, Kozelko S, Béliveau R, Marsolais C, Annabi B. TH1902, a new docetaxel-peptide conjugate for the treatment of sortilin-positive triple-negative breast cancer. *Cancer Sci* 2021; **112**: 4317-4334 [PMID: [34314556](#) DOI: [10.1111/cas.15086](#)]
- 32 **Liang M**, Yao W, Shi B, Zhu X, Cai R, Yu Z, Guo W, Wang H, Dong Z, Lin M, Zhou X, Zheng Y. Circular RNA hsa_circ_0110389 promotes gastric cancer progression through upregulating SORT1 via sponging miR-127-5p and miR-136-5p. *Cell Death Dis* 2021; **12**: 639 [PMID: [34162830](#) DOI: [10.1038/s41419-021-03903-5](#)]
- 33 **Shi J**, Jiang D, Yang S, Sun Y, Wang J, Zhang X, Liu Y, Lu Y, Yang K. Molecular profile reveals immune-associated markers of lymphatic invasion in human colon adenocarcinoma. *Int Immunopharmacol* 2020; **83**: 106402 [PMID: [32200154](#) DOI: [10.1016/j.intimp.2020.106402](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

