

World Journal of *Clinical Cases*

World J Clin Cases 2024 January 6; 12(1): 1-235



Contents

Thrice Monthly Volume 12 Number 1 January 6, 2024

OPINION REVIEW

- 1 Gut-targeted therapies for type 2 diabetes mellitus: A review
Xu TC, Liu Y, Yu Z, Xu B

MINIREVIEWS

- 9 Honeymoon phase in type 1 diabetes mellitus: A window of opportunity for diabetes reversal?
Mittal M, Porchezian P, Kapoor N

ORIGINAL ARTICLE

Retrospective Cohort Study

- 15 Evaluating combined bevacizumab and XELOX in advanced colorectal cancer: Serum markers carcinoembryonic antigen, carbohydrate antigen 125, carbohydrate antigen 199 analysis
Zhou DB, Cheng J, Zhang XH
- 24 Clinical value of precise rehabilitation nursing in management of cerebral infarction
Xu YN, Wang XZ, Zhang XR

Retrospective Study

- 32 Marker Ki-67 is a potential biomarker for the diagnosis and prognosis of prostate cancer based on two cohorts
Song Z, Zhou Q, Zhang JL, Ouyang J, Zhang ZY
- 42 Natural history of asymptomatic gallbladder stones in clinic without beds: A long-term prognosis over 10 years
Sakai Y, Tsuyuguchi T, Ohyama H, Kumagai J, Kaiho T, Ohtsuka M, Kato N, Sakai T
- 51 Clinical nursing value of predictive nursing in reducing complications of pregnant women undergoing short-term massive blood transfusion during cesarean section
Cheng L, Li LP, Zhang YY, Deng F, Lan TT
- 59 Effect of cardiac rehabilitation care after coronary intervention on cardiac function recovery and negative mood in patients with myocardial infarction
Yang M, Huang YT, Hu XW, Wu CL
- 68 Efficacy and safety of Nafamostat mesylate in patients with end-stage renal failure
Liu K, Li ZH
- 76 Nursing effect of narrative nursing intervention on postoperative patients with severe lung cancer
Wen B, Liu Y, Min XX, Wang AQ

Observational Study

- 86 Interaction between adolescent sleep rhythms and gender in an obese population
Wu NN, Yan GL, Zhang HY, Sun L, Hou M, Xu GM

SYSTEMATIC REVIEWS

- 95 Endoscopic submucosal dissection *vs* transanal endoscopic surgery for rectal tumors: A systematic review and meta-analysis
Huang LW, Zhong Y
- 107 Impact of frailty on outcomes of elderly patients undergoing percutaneous coronary intervention: A systematic review and meta-analysis
Wang SS, Liu WH
- 119 Nasogastric tube syndrome: A Meta-summary of case reports
Juneja D, Nasa P, Chanchalani G, Jain R

CASE REPORT

- 130 Erythrodermic mycosis fungoides: A case report
Xu WB, Zhang YP, Zhou SP, Bai HY
- 136 Azacitidine maintenance therapy for blastic plasmacytoid dendritic cell neoplasm allograft: A case report
Tao LL, Wen HT, Wang ZY, Cheng J, Zhao L
- 142 Congestive ischemic colitis successfully treated with anti-inflammatory therapy: A case report
Lee GW, Park SB
- 148 Subarachnoid hemorrhage misdiagnosed as acute coronary syndrome leading to catastrophic neurologic injury: A case report
Lin JM, Yuan XJ, Li G, Gan XR, Xu WH
- 157 Successful management of severe hypoglycemia induced by total parenteral nutrition in patients with hepatocellular injury: Three cases reports
Fang LZ, Jin HX, Zhao N, Wu YP, Shi YQ
- 163 Endophthalmitis in silicone oil-filled eye: A case report
Yan HC, Wang ZL, Yu WZ, Zhao MW, Liang JH, Yin H, Shi X, Miao H
- 169 Lung imaging characteristics in a patient infected with *Elizabethkingia miricola* following cerebral hemorrhage surgery: A case report
Qi PQ, Zeng YJ, Peng W, Kuai J
- 176 Gastric IgG4-related disease mimicking a gastrointestinal stromal tumor in a child: A case report
Lin HCA, Lee KF, Huang TH
- 180 Labial inverse dilaceration of bilateral maxillary central incisors: A case report
Wang JM, Guo LF, Ma LQ, Zhang J

- 188 Changes in macrophage infiltration and podocyte injury in lupus nephritis patients with repeated renal biopsy: Report of three cases
Liu SY, Chen H, He LJ, Huang CK, Wang P, Rui ZR, Wu J, Yuan Y, Zhang Y, Wang WJ, Wang XD
- 196 Primary acinic cell carcinoma of the breast: A case report and review of literature
Ding JS, Zhang M, Zhou FF
- 204 Acupuncture for cervical dystonia associated with anxiety and depression: A case report
Zhang YT, Zhang JJ, Zha BX, Fan YQ, Xu YB, Yang J, Zhang QP
- 210 Intestinal malrotation complicated with gastric cancer: A case report
Jia XH, Kong S, Gao XX, Cong BC, Zheng CN
- 217 Addison's disease caused by adrenal tuberculosis may lead to misdiagnosis of major depressive disorder: A case report
Zhang TX, Xu HY, Ma W, Zheng JB
- 224 Pleural empyema with endobronchial mass due to *Rhodococcus equi* infection after renal transplantation: A case report and review of literature
Liang GF, Chao S, Sun Z, Zhu KJ, Chen Q, Jia L, Niu YL

LETTER TO THE EDITOR

- 232 Chronic venous insufficiency, could it be one of the missing pieces in the puzzle of treating pain?
Chang MC

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Woon-Man Kung, MD, Associate Professor, Surgeon, Department of Exercise and Health Promotion, College of Kinesiology and Health, Chinese Culture University, Taipei 11114, Taiwan. nskungwm@yahoo.com.tw

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJCC* as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Si Zhao*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Salim Surani, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

January 6, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Natural history of asymptomatic gallbladder stones in clinic without beds: A long-term prognosis over 10 years

Yuji Sakai, Toshio Tsuyuguchi, Hiroshi Ohyama, Junichiro Kumagai, Takashi Kaiho, Masayuki Ohtsuka, Naoya Kato, Tadao Sakai

Specialty type: Gastroenterology & hepatology

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Fan Y, China

Received: September 3, 2023

Peer-review started: September 3, 2023

First decision: October 24, 2023

Revised: October 31, 2023

Accepted: December 18, 2023

Article in press: December 18, 2023

Published online: January 6, 2024



Yuji Sakai, Tadao Sakai, Department of Gastroenterology, Sakai Clinic, Kimistu 299-1162, Japan

Yuji Sakai, Hiroshi Ohyama, Naoya Kato, Department of Gastroenterology, Graduate School of Medicine, Chiba University, Chiba 260-8677, Japan

Toshio Tsuyuguchi, Department of Gastroenterology, Chiba Prefectural Sawara Hospital, Chiba Prefectural Sawara Hospital, Sawara 287-0003, Japan

Junichiro Kumagai, Department of Gastroenterology, Kimitsu Central Hospital, Kisarazu 292-8535, Japan

Takashi Kaiho, Department of Surgery, Kimitsu Central Hospital, Kisarazu 292-8535, Japan

Masayuki Ohtsuka, Department of General Surgery, Graduate School of Medicine, Chiba University, Chiba 260-8677, Japan

Corresponding author: Yuji Sakai, MD, PhD, Doctor, Department of Gastroenterology, Sakai Clinic, 9-18-8 Minamikoyasu, Kimistu 299-1162, Japan. sakai4754@yahoo.co.jp

Abstract

BACKGROUND

Several studies have explored the long-term prognosis of patients with asymptomatic gallbladder stones. These reports were primarily conducted in facilities equipped with beds for addressing symptomatic cases.

AIM

To report the long-term prognosis of patients with asymptomatic gallbladder stones in clinics without bed facilities.

METHODS

We investigated the prognoses of 237 patients diagnosed with asymptomatic gallbladder stones in clinics without beds between March 2010 and October 2022. When symptoms developed, patients were transferred to hospitals where appropriate treatment was possible. We investigated the asymptomatic and survival periods during the follow-up.

RESULTS

Among the 237 patients, 214 (90.3%) remained asymptomatic, with a mean asymptomatic period of 3898.9279 ± 46.871 d (50-4111 d, 10.7 years on average). Biliary complications developed in 23 patients (9.7%), with a mean survival period of 4010.0285 ± 31.2788 d (53-4112 d, 10.9 years on average). No patient died of biliary complications.

CONCLUSION

The long-term prognosis of asymptomatic gallbladder stones in clinics without beds was favorable. When the condition became symptomatic, the patients were transferred to hospitals with beds that could address it; thus, no deaths related to biliary complications were reported. This finding suggests that follow-up care in clinics without beds is possible.

Key Words: Gallbladder stone; Acute cholangitis; Acute cholecystitis; Asymptomatic gallbladder stone; Symptomatic gallbladder stone

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: A long-term prognosis of asymptomatic gallbladder stones at a clinic without beds was favorable. Patients with asymptomatic gallbladder stones in a bedless clinic were well-cared for and had a long duration of no symptoms.

Citation: Sakai Y, Tsuyuguchi T, Ohyama H, Kumagai J, Kaiho T, Ohtsuka M, Kato N, Sakai T. Natural history of asymptomatic gallbladder stones in clinic without beds: A long-term prognosis over 10 years. *World J Clin Cases* 2024; 12(1): 42-50

URL: <https://www.wjgnet.com/2307-8960/full/v12/i1/42.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v12.i1.42>

INTRODUCTION

The reported prevalence of gallbladder stones is reported to be approximately 10%-25% in the United States and Europe [1-5]. The widespread use of abdominal ultrasonography has greatly facilitated the diagnosis of gallbladder stones [6-8]. For asymptomatic patients, clinical observation is typically sufficient. However, when symptoms, such as acute cholecystitis or acute cholangitis, manifest, appropriate treatments, including drainage, surgery, and administration of antibiotics, become essential [9-11].

Guidelines for acute cholecystitis and acute cholangitis clearly specify the criteria for transferring patients to facilities capable of providing appropriate treatment [12]. Diagnostic criteria have also been established for acute pancreatitis triggered by gallbladder stones [13]. Once patients are diagnosed with acute pancreatitis, it becomes imperative to transfer them to facilities with inpatient services, as untreated acute pancreatitis may lead to organ failure or life-threatening infections. Therefore, understanding the timing of transferring patients to facilities with inpatient services when gallbladder stones become symptomatic is crucial.

Although it is important for the attending physician to be aware of potential symptoms, asymptomatic gallbladder stones are frequently managed through periodic monitoring in clinics. To date, several reports have provided insights into the long-term prognosis of asymptomatic gallbladder stones mainly from facilities equipped to accommodate inpatients [14-19]. To the best of our knowledge, no studies have been reported from clinics that lack inpatient facilities that have observed clinical courses of asymptomatic gallbladder stones for more than a decade. Therefore, in this study, we explored the long-term prognosis of asymptomatic gallbladder stones at a clinic without beds.

MATERIALS AND METHODS

Between March 2010 and October 2022, we followed up 237 patients diagnosed with gallbladder stones among 3268 cases of abdominal ultrasonography performed during screening at Sakai Clinic, a bedless clinic. Patients with gallbladder stones visited the hospital once a month for evaluation of their condition. At the time of consultation, all 237 patients had asymptomatic gallbladder stones, and none of them had a history of seizures. Patients' demographic details are summarized in Table 1. The study included 123 males and 114 females, with a mean age of 47.675 ± 15.625 (range: 20-78) years. The mean number of gallbladder stones was 2.318 ± 3.500 (range: 1-20), and the mean stone diameter was 7.343 ± 4.298 (range: 1-22) mm. Ursodeoxycholic acid (UDCA) was administered based on patient's request after they were informed about its pharmacological effects. Patients who took UDCA ≥ 300 mg daily for 6 mo or more were categorized into the active treatment arm (UDCA group), and this arm included 68 patients. Comorbidity included hypertension in 125 (52.7%) patients, hyperlipidemia in 119 (50.2%) patients, diabetes mellitus in 110 (46.4%) patients, cerebrovascular disease in 42 (17.7%) patients, chronic respiratory disease in 37 (15.6%) patients, heart disease in 28 (11.8%) patients, chronic liver disease in 35 (14.8%) patients, chronic renal disease in 42 (17.7%) patients, and malignant disease in 10 (4.2%) patients. Patients diagnosed with gallbladder stones underwent abdominal ultrasonography once every 6-12 mo. Blood

Table 1 Patient characteristics

Characteristic	Value
Number of patients	237
Male	123
Female	114
Age in yr	47.675 ± 15.625 (20-78)
Gallbladder stone	
Number of stones	2,318 ± 3,500 (1-20)
Size in mm	7,343 ± 4,298 (1-22)
UDCA	68
Hypertension	125
Hyperlipidemia	119
Diabetes mellitus	110
Cerebrovascular disease	42
Chronic respiratory disease	37
Cardiac disease	28
Chronic liver disease	35
Chronic renal disease	42
Malignant disease	10

UDCA: Ursodeoxycholic acid.

samples were collected at least once every 6 mo or once a month depending on the comorbidity. Abdominal ultrasonography was routinely performed by changing the posture to depict the location of gallbladder and common bile duct. In cases where the visualization of the common bile duct was difficult with abdominal ultrasonography, abdominal computed tomography (CT) was performed. Abdominal ultrasonography was performed using one of the two models (TOSHIBA SSA-250A, Japan or Canon Xario 100, Japan). CT was performed using a 16-slice MDCT scanner (TOSHIBA Activion16, Japan), with a detector collimation of 1 mm × 16 mm, table feed of 24 mm per rotation, rotation time of 1 s, tube current of 200 effective mAs, and tube voltage of 120 kV. Axial images were reconstructed by 5 mm without overlap and coronal images by 3 mm without overlap. No contrast medium was used for the assessments.

Biliary complications, such as acute cholecystitis, acute cholangitis, obstructive jaundice, and gallstone pancreatitis, were considered symptomatic. Abdominal pain during the time-course observation caused by reasons other than gallbladder stones was not considered symptomatic. Emergency abdominal ultrasonography was performed when patients developed abdominal pain, fever, or jaundice during clinic visits. If additional tests were necessary, they were conducted at the discretion of physicians. When acute cholecystitis, acute cholangitis, or gallstone pancreatitis was suspected, patients were promptly transferred to hospitals equipped to provide appropriate care, following the Tokyo Guidelines (for acute cholecystitis and acute cholangitis)[20-25] and the revised Atlanta classification (for acute pancreatitis) that existed at the time of onset[13].

To assess the long-term prognosis, we directly communicated with the patient and transfer hospitals, in addition to utilizing data from the outpatient clinic. The study was conducted at this clinic after obtaining consent from all patients regarding the use of the data. We obtained approval from the institutional review board of Chiba Prefectural Sawara Hospital (IRB number 4-12). The authors declare no conflicts of interest.

Statistical analysis

For statistical analysis of categorical variables, the Pearson χ^2 test with Yates correction and Fisher's exact test were used, as deemed appropriate. Differences were considered statistically significant at *P* values < 0.05. Asymptomatic and survival periods were estimated using the Kaplan-Meier method. Data were analyzed using Excel software (Version 2020).

RESULTS

A total of 214 (90.3 %) patients remained asymptomatic during the observation period, lasting for an average duration of 3898.9279 ± 46.871 d (50-4111 d, 10.7 years on average) (Figure 1A). Among them, 23 (9.7%) patients experienced biliary




Figure 1 Long-term prognosis of asymptomatic gallbladder stones. A: Asymptomatic period; B: Survival period.

complications, including acute cholecystitis in 14 (5.9%) patients, acute cholangitis in 5 (2.1%) patients, obstructive jaundice in 2 (0.8%) patients, and gallstone pancreatitis in 2 (0.8%) patients (Table 2). Comparing the group with no biliary complications during long-term follow-up to the group with biliary complications revealed significantly higher rates of biliary complications in the group with multiple gallbladder stones, those with stones 10 mm or less in diameter, and the non-UDCA group (Table 3).

The mean survival period was 4010.0285 ± 31.2788 d (53-4112 d, 10.9 years on average) (Figure 1B). Eleven (4.6%) deaths occurred during the observation period, with the causes being lung cancer in 4 (1.7%) patients, cerebral infarction in 3 (1.3%) patients, acute myocardial infarction in 2 (0.8%) patients, gastric cancer in 1 (0.4%) patient, and colorectal cancer in 1 (0.4%) patient (Table 4). Notably, no deaths due to biliary complications were noted during the observation period for asymptomatic gallstones.

Prognoses were ascertained through outpatient visits alone in 211 (89%) patients and by inquiring hospitals to which patients were transferred in 13 (5.5%) patients. For the remaining 13 (5.5%) patients, the prognoses were determined through direct telephone conversation with them. Prognoses were determined for all 237 (100%) patients. Notably, 22 (9.3%) patients underwent abdominal CT due to the difficulty in visualizing the common bile duct using ultrasonography.

All 23 patients with biliary complications were transferred from the clinic to hospitals. All clinical and hospital diagnoses were consistent. Out of the 23 patients with biliary complications, 20 underwent cholecystectomies. Laparoscopic cholecystectomy and open cholecystectomy were performed in 18 patients and 2 patients, respectively. However, one of the patients, a 78-year-old man with lung cancer, who developed acute cholecystitis did not undergo surgery because his condition improved with conservative antibiotic therapy. When he provided informed consent regarding his prognosis, he did not wish to undergo further surgery. Instead, he was placed under a time-course observation, and no recurrence of acute cholecystitis was observed for 18 mo until he succumbed to cancer. Two other women who did not undergo surgery developed acute cholangitis, and their symptoms improved after endoscopic transpapillary removal of the common bile duct stones. Both the patients were 85 years old with multiple comorbidities. They were considered unfit for surgery; therefore, they were closely monitored without surgical intervention. In both cases, neither patient had experienced any seizures for over 2 years.

Postoperative biloma was noted in 1 patient (5%) who underwent laparoscopic cholecystectomy for acute cholecystitis due to surgery-related complications; however, the condition improved with conservative therapy (Figure 2). No accidental surgery-related symptoms were observed. In 1 case of acute cholecystitis, histopathological examination following laparoscopic cholecystectomy revealed carcinoma in situ. This case initially presented with numerous small stones, as observed on abdominal ultrasonography, which made evaluation of the gallbladder wall challenging. The patient has been followed up for 12 mo postoperatively, but no recurrence was noted. Gallbladder cancer was observed in only this patient (0.004%) during the observation period.

DISCUSSION

Several reports on the natural history of asymptomatic gallbladder stones have been published to date, primarily originating from institutions with inpatient facilities and research institutions[14-19]. Our study focused on reports from clinics without beds, but no major differences were observed compared to reports from clinics with beds and research institutions. In symptomatic patients, the diameter of stones was smaller and the number of stones was significantly higher than those of asymptomatic patients. Gallbladder stones with smaller diameters are more likely to affect the cystic duct than those with larger diameters. Similarly, gallbladder stones with smaller diameters are more likely to fall into the common bile duct than those with larger diameters. As the number of gallbladder stones increases, the probability of the

Table 2 Biliary complications noted during the observation period

Biliary complication	Value
Biliary event	23
Acute cholecystitis	14
Acute cholangitis	5
Obstructive jaundice	2
Gallstone pancreatitis	2

Table 3 Comparison of the groups with and without biliary complications

Characteristic	No symptom, <i>n</i> = 214	Biliary event, <i>n</i> = 23	<i>P</i> value
Male	108	15	0.1951
Female	106	8	0.1951
Gallstone			
Number ≤ 2	3	20	< 0.0010
Diameter < 10 mm	3	20	< 0.0010
UDCA	66	2	0.0277
Hypertension	115	10	0.6653
Hyperlipidemia	108	11	0.8293
Diabetes mellitus	102	8	0.5078
Cerebrovascular disease	36	6	0.1111
Chronic respiratory disease	33	4	0.5384
Cardiac disease	26	2	1.0000
Chronic liver disease	32	3	1.0000
Chronic renal disease	36	6	0.1111
Malignant disease	9	1	1.0000


UDCA: Ursodeoxycholic acid.

Table 4 Causes of deaths during the observation period

Causes of deaths	<i>n</i>
Total number	11
Lung cancer	4
Cerebral infarction	3
Acute myocardial infarction	2
Gastric cancer	1
Colorectal cancer	1

stones impacting the cystic duct or falling into the common bile duct is also expected to increase. A single large gallbladder stone is unlikely to fall into the common bile duct, which may explain our findings.

In addition, the UDCA group was significantly less likely to become symptomatic than the non-UDCA group. UDCA, known for promoting bile outflow, may suppress the increase in gallbladder stones by promoting bile outflow. Therefore, it may prevent gallbladder stones from becoming impacted in the cystic duct or falling into the common bile duct, which may reduce the likelihood of symptoms. Previous studies have reported that long-term administration of UDCA has a prophylactic effect against the development of symptomatic gallbladder stones[26]. Some reports have suggested that short-term UDCA administration during the waiting period for cholecystectomy does not prevent the condition from



DOI: 10.12998/wjcc.v12.i1.42 Copyright ©The Author(s) 2024.

Figure 2 A 68-year-old female. A: Computed tomography revealed gallbladder stones; B: Onset of acute cholecystitis 1215 d after the diagnosis of gallbladder stones; C: Biloma was noted on the computed tomography scan after laparoscopic cholecystectomy (arrow).

becoming symptomatic[27]. Therefore, it is unlikely that UDCA has little preventive effect against symptomatic gallbladder stones when administered for a brief period.

In this study, 1 case of gallbladder cancer was identified, demonstrating carcinoma in situ upon pathological examination of resected specimen following laparoscopic cholecystectomy for acute cholecystitis. Abdominal ultrasonography revealed a number of small stones, posing challenges in the evaluation of the gallbladder. Preoperative examination at the hospital to which the patient was transferred did not detect the presence of gallbladder cancer. A meta-analysis of three cohort studies and seven case-control studies revealed that gallbladder stones present the strongest risk factor for gallbladder cancer, with a relative risk of 4.9[28]. Meanwhile, a cohort study spanning approximately 11 years, including more than 110000 patients, reported a hazard ratio of 1.07 for gallbladder stones and a history of cholecystitis in relation to carcinogenesis, showing a negative association between gallbladder stones and the development of gallbladder cancer[29]. The incidence of gallbladder cancer in patients with gallbladder stones is as low as 0.01%-0.02% annually. Observational studies over 5 years have also shown an extremely limited incidence of approximately 0.3%[30]. Although the causal relationship between gallbladder stones and gallbladder cancer is well known, concrete evidence directly linking gallbladder stones to the development of gallbladder cancer is currently lacking. In this study, the incidence of gallbladder cancer was extremely low (0.004%). Preventive cholecystectomy can help obviate symptoms or hedge the risk of gallbladder cancer.

Cholecystectomy may cause life-threatening complications, such as vascular and bile duct injuries, albeit these are extremely rare[10]. In our study, biloma was noted after the procedure, although it was not a serious complication. Notably, these complications occur at a certain rate after cholecystectomy. Although no prospective studies have been conducted concerning cholecystectomy *vs* time-course observation, the therapeutic intervention for asymptomatic cases offers little benefits. Cholecystectomy should be considered only when its benefits outweigh its risks. In cases where evaluation of the gallbladder wall is difficult because of chronic cholecystitis or gallstones, surgery should be considered even in the absence of symptoms.

The long-term prognosis of asymptomatic gallstones in clinics without beds was favorable, with no deaths due to biliary complications when medical care was provided according to the criteria for diagnosis and transportation. The follow-up of these patients at the outpatient clinic was satisfactory. However, this favorable long-term prognosis is only for asymptomatic patients and not for symptomatic gallbladder stones. Symptomatic gallstones have a higher incidence of biliary complications than asymptomatic gallstones[16-19]. We cannot rule out the possibility that different results could be obtained in a long-term prognostic study. Notably, our study was retrospective and conducted at a single institution. A prospective study with a larger number of bedless clinics is necessary in the future to further investigate this matter.

CONCLUSION

The long-term prognosis of asymptomatic gallbladder stones in clinics without beds is favorable. When the condition became symptomatic, the patients were transported to hospitals equipped with beds to manage the condition, contributing to the absence of reported deaths related to biliary complications. The follow-up of asymptomatic gallbladder stones in a clinic without inpatient beds was deemed effective.

ARTICLE HIGHLIGHTS

Research background

Several reports on a long-term prognosis of asymptomatic gallbladder stones that have been available to date are mainly from facilities that can accommodate inpatients. As far as we know, there has been no report to date from a clinic without

beds that observed clinical courses of more than a decade.

Research motivation

There is currently no study of the long-term prognosis of asymptomatic gallbladder stones from a clinic without beds.

Research objectives

To determine the long-term prognosis of asymptomatic gallbladder stones in a clinic without beds.

Research methods

We followed up 237 cases diagnosed with asymptomatic gallbladder stones.

Research results

Those patients whose condition was asymptomatic during the observation period accounted for 214 cases (90.3%). The asymptomatic period was 3898.9279 ± 46.871 d (50–4111 d, 10.7 years on average).

Research conclusions

When the condition turned symptomatic, patients were transported to hospitals with beds that could address the condition, and no deaths related to biliary complications was reported. Asymptomatic gallbladder stones were considered to be well followed in a bedless clinic.

Research perspectives

Our study was a retrospective study in a single institution. A prospective study in a large number of bedless clinics is considered necessary in the future.

FOOTNOTES

Author contributions: Sakai Y, Ohtsuka M, Tsuyuguchi T, and Kato N were responsible for manuscript preparation; Sakai Y wrote the paper; Tsuyuguchi T and Sakai T revised the manuscript; Ohyama H, Kumagai J, and Kaiho T collected the references.

Institutional review board statement: A study at this clinic was conducted after obtaining approval from all patients about the use of data. We requested the Institutional Review Board at Chiba Prefectural Sawara Hospital to review data and obtained their approval (IRB No. 4-12).

Informed consent statement: In order to know the long-term prognosis, we confirmed the prognosis by directly calling the patient and the transportation hospitals in addition to the information at the outpatient clinic.

Conflict-of-interest statement: All authors declare having no conflicts of interest.

Data sharing statement: The data that support the findings of this study are available from the corresponding author, upon reasonable request.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Japan

ORCID number: Yuji Sakai 0000-0002-1782-4829; Toshio Tsuyuguchi 0000-0003-4897-5635; Hiroshi Ohyama 0000-0003-0614-581X; Junichiro Kumagai 0000-0002-9031-0103; Masayuki Ohtsuka 0000-0002-5091-1078.

S-Editor: Yan JP

L-Editor: Filipodia

P-Editor: Yan JP

REFERENCES

- 1 Everhart JE, Khare M, Hill M, Maurer KR. Prevalence and ethnic differences in gallbladder disease in the United States. *Gastroenterology* 1999; **117**: 632-639 [PMID: 10464139 DOI: 10.1016/s0016-5085(99)70456-7]
- 2 Glambek I, Kvaale G, Arnesjö B, Søreide O. Prevalence of gallstones in a Norwegian population. *Scand J Gastroenterol* 1987; **22**: 1089-1094 [PMID: 3321395 DOI: 10.3109/00365528708991963]
- 3 Muhrbeck O, Ahlberg J. Prevalence of gallstone disease in a Swedish population. *Scand J Gastroenterol* 1995; **30**: 1125-1128 [PMID: 8540000]

- 8578175 DOI: [10.3109/00365529509101618](https://doi.org/10.3109/00365529509101618)]
- 4 Shaffer EA. Gallstone disease: Epidemiology of gallbladder stone disease. *Best Pract Res Clin Gastroenterol* 2006; **20**: 981-996 [PMID: [17127183](https://pubmed.ncbi.nlm.nih.gov/17127183/) DOI: [10.1016/j.bpg.2006.05.004](https://doi.org/10.1016/j.bpg.2006.05.004)]
- 5 Portincasa P, Moschetta A, Palasciano G. Cholesterol gallstone disease. *Lancet* 2006; **368**: 230-239 [PMID: [16844493](https://pubmed.ncbi.nlm.nih.gov/16844493/) DOI: [10.1016/S0140-6736\(06\)69044-2](https://doi.org/10.1016/S0140-6736(06)69044-2)]
- 6 Gustafsson C, McNicholas A, Sondén A, Törmgren S, Järnbert-Pettersson H, Lindelius A. Accuracy of Surgeon-Performed Ultrasound in Detecting Gallstones: A Validation Study. *World J Surg* 2016; **40**: 1688-1694 [PMID: [26935564](https://pubmed.ncbi.nlm.nih.gov/26935564/) DOI: [10.1007/s00268-016-3468-3](https://doi.org/10.1007/s00268-016-3468-3)]
- 7 Carroll PJ, Gibson D, El-Faedy O, Dunne C, Coffey C, Hannigan A, Walsh SR. Surgeon-performed ultrasound at the bedside for the detection of appendicitis and gallstones: systematic review and meta-analysis. *Am J Surg* 2013; **205**: 102-108 [PMID: [22748292](https://pubmed.ncbi.nlm.nih.gov/22748292/) DOI: [10.1016/j.amjsurg.2012.02.017](https://doi.org/10.1016/j.amjsurg.2012.02.017)]
- 8 Fang R, Pilcher JA, Putnam AT, Smith T, Smith DL. Accuracy of surgeon-performed gallbladder ultrasound. *Am J Surg* 1999; **178**: 475-479 [PMID: [10670856](https://pubmed.ncbi.nlm.nih.gov/10670856/) DOI: [10.1016/s0002-9610\(99\)00225-1](https://doi.org/10.1016/s0002-9610(99)00225-1)]
- 9 Sakai Y, Tsuyuguchi T, Kato K, Sugiyama H, Nishikawa T, Takahashi M, Yokosuka O. Clinical usefulness of doripenem (DRPM), a carbapenem antimicrobial drug, for the treatment of patients with acute cholangitis: retrospective study. *Int J Clin Pharmacol Ther* 2013; **51**: 19-25 [PMID: [23073141](https://pubmed.ncbi.nlm.nih.gov/23073141/) DOI: [10.5414/CP201786](https://doi.org/10.5414/CP201786)]
- 10 Gallaheer JR, Charles A. Acute Cholecystitis: A Review. *JAMA* 2022; **327**: 965-975 [PMID: [35258527](https://pubmed.ncbi.nlm.nih.gov/35258527/) DOI: [10.1001/jama.2022.2350](https://doi.org/10.1001/jama.2022.2350)]
- 11 Ito K, Fujita N, Noda Y, Kobayashi G, Kimura K, Sugawara T, Horaguchi J. Percutaneous cholecystostomy vs gallbladder aspiration for acute cholecystitis: a prospective randomized controlled trial. *AJR Am J Roentgenol* 2004; **183**: 193-196 [PMID: [15208137](https://pubmed.ncbi.nlm.nih.gov/15208137/) DOI: [10.2214/ajr.183.1.1830193](https://doi.org/10.2214/ajr.183.1.1830193)]
- 12 Miura F, Okamoto K, Takada T, Strasberg SM, Asbun HJ, Pitt HA, Gomi H, Solomkin JS, Schlossberg D, Han HS, Kim MH, Hwang TL, Chen MF, Huang WS, Kiriyaama S, Itoi T, Garden OJ, Liau KH, Horiguchi A, Liu KH, Su CH, Gouma DJ, Belli G, Dervenis C, Jagannath P, Chan ACW, Lau WY, Endo I, Suzuki K, Yoon YS, de Santibañes E, Giménez ME, Jonas E, Singh H, Honda G, Asai K, Mori Y, Wada K, Higuchi R, Watanabe M, Rikiyama T, Sata N, Kano N, Umezawa A, Mukai S, Tokumura H, Hata J, Kozaka K, Iwashita Y, Hibi T, Yokoe M, Kimura T, Kitano S, Inomata M, Hirata K, Sumiyama Y, Inui K, Yamamoto M. Tokyo Guidelines 2018: initial management of acute biliary infection and flowchart for acute cholangitis. *J Hepatobiliary Pancreat Sci* 2018; **25**: 31-40 [PMID: [28941329](https://pubmed.ncbi.nlm.nih.gov/28941329/) DOI: [10.1002/jhbp.509](https://doi.org/10.1002/jhbp.509)]
- 13 Banks PA, Bollen TL, Dervenis C, Gooszen HG, Johnson CD, Sarr MG, Tsiotos GG, Vege SS; Acute Pancreatitis Classification Working Group. Classification of acute pancreatitis--2012: revision of the Atlanta classification and definitions by international consensus. *Gut* 2013; **62**: 102-111 [PMID: [23100216](https://pubmed.ncbi.nlm.nih.gov/23100216/) DOI: [10.1136/gutjnl-2012-302779](https://doi.org/10.1136/gutjnl-2012-302779)]
- 14 Gracie WA, Ransohoff DF. The natural history of silent gallstones: the innocent gallstone is not a myth. *N Engl J Med* 1982; **307**: 798-800 [PMID: [7110244](https://pubmed.ncbi.nlm.nih.gov/7110244/) DOI: [10.1056/NEJM198209233071305](https://doi.org/10.1056/NEJM198209233071305)]
- 15 Morris-Stiff G, Sarvepalli S, Hu B, Gupta N, Lal P, Burke CA, Garber A, McMichael J, Rizk MK, Vargo JJ, Ibrahim M, Rothberg MB. The Natural History of Asymptomatic Gallstones: A Longitudinal Study and Prediction Model. *Clin Gastroenterol Hepatol* 2023; **21**: 319-327.e4 [PMID: [35513234](https://pubmed.ncbi.nlm.nih.gov/35513234/) DOI: [10.1016/j.cgh.2022.04.010](https://doi.org/10.1016/j.cgh.2022.04.010)]
- 16 Festi D, Reggiani ML, Attili AF, Loria P, Pazzi P, Scaioi E, Capodicasa S, Romano F, Roda E, Colecchia A. Natural history of gallstone disease: Expectant management or active treatment? Results from a population-based cohort study. *J Gastroenterol Hepatol* 2010; **25**: 719-724 [PMID: [20492328](https://pubmed.ncbi.nlm.nih.gov/20492328/) DOI: [10.1111/j.1440-1746.2009.06146.x](https://doi.org/10.1111/j.1440-1746.2009.06146.x)]
- 17 McSherry CK, Ferstenberg H, Calhoun WF, Lahman E, Virshup M. The natural history of diagnosed gallstone disease in symptomatic and asymptomatic patients. *Ann Surg* 1985; **202**: 59-63 [PMID: [4015212](https://pubmed.ncbi.nlm.nih.gov/4015212/) DOI: [10.1097/0000658-198507000-00009](https://doi.org/10.1097/0000658-198507000-00009)]
- 18 Friedman GD, Raviola CA, Fireman B. Prognosis of gallstones with mild or no symptoms: 25 years of follow-up in a health maintenance organization. *J Clin Epidemiol* 1989; **42**: 127-136 [PMID: [2918322](https://pubmed.ncbi.nlm.nih.gov/2918322/) DOI: [10.1016/0895-4356\(89\)90086-3](https://doi.org/10.1016/0895-4356(89)90086-3)]
- 19 Persson GE. Expectant management of patients with gallbladder stones diagnosed at planned investigation. A prospective 5- to 7-year follow-up study of 153 patients. *Scand J Gastroenterol* 1996; **31**: 191-199 [PMID: [8658043](https://pubmed.ncbi.nlm.nih.gov/8658043/) DOI: [10.3109/00365529609031985](https://doi.org/10.3109/00365529609031985)]
- 20 Hirota M, Takada T, Kawarada Y, Nimura Y, Miura F, Hirata K, Mayumi T, Yoshida M, Strasberg S, Pitt H, Gadacz TR, de Santibanes E, Gouma DJ, Solomkin JS, Belghiti J, Neuhaus H, Büchler MW, Fan ST, Ker CG, Padbury RT, Liau KH, Hilvano SC, Belli G, Windsor JA, Dervenis C. Diagnostic criteria and severity assessment of acute cholecystitis: Tokyo Guidelines. *J Hepatobiliary Pancreat Surg* 2007; **14**: 78-82 [PMID: [17252300](https://pubmed.ncbi.nlm.nih.gov/17252300/) DOI: [10.1007/s00534-006-1159-4](https://doi.org/10.1007/s00534-006-1159-4)]
- 21 Wada K, Takada T, Kawarada Y, Nimura Y, Miura F, Yoshida M, Mayumi T, Strasberg S, Pitt HA, Gadacz TR, Büchler MW, Belghiti J, de Santibanes E, Gouma DJ, Neuhaus H, Dervenis C, Fan ST, Chen MF, Ker CG, Bornman PC, Hilvano SC, Kim SW, Liau KH, Kim MH. Diagnostic criteria and severity assessment of acute cholangitis: Tokyo Guidelines. *J Hepatobiliary Pancreat Surg* 2007; **14**: 52-58 [PMID: [17252297](https://pubmed.ncbi.nlm.nih.gov/17252297/) DOI: [10.1007/s00534-006-1156-7](https://doi.org/10.1007/s00534-006-1156-7)]
- 22 Yokoe M, Takada T, Strasberg SM, Solomkin JS, Mayumi T, Gomi H, Pitt HA, Gouma DJ, Garden OJ, Büchler MW, Kiriyaama S, Kimura Y, Tsuyuguchi T, Itoi T, Yoshida M, Miura F, Yamashita Y, Okamoto K, Gabata T, Hata J, Higuchi R, Windsor JA, Bornman PC, Fan ST, Singh H, de Santibanes E, Kusachi S, Murata A, Chen XP, Jagannath P, Lee S, Padbury R, Chen MF; Tokyo Guidelines Revision Committee. New diagnostic criteria and severity assessment of acute cholecystitis in revised Tokyo Guidelines. *J Hepatobiliary Pancreat Sci* 2012; **19**: 578-585 [PMID: [22872303](https://pubmed.ncbi.nlm.nih.gov/22872303/) DOI: [10.1007/s00534-012-0548-0](https://doi.org/10.1007/s00534-012-0548-0)]
- 23 Kiriyaama S, Takada T, Strasberg SM, Solomkin JS, Mayumi T, Pitt HA, Gouma DJ, Garden OJ, Büchler MW, Yokoe M, Kimura Y, Tsuyuguchi T, Itoi T, Yoshida M, Miura F, Yamashita Y, Okamoto K, Gabata T, Hata J, Higuchi R, Windsor JA, Bornman PC, Fan ST, Singh H, de Santibanes E, Gomi H, Kusachi S, Murata A, Chen XP, Jagannath P, Lee S, Padbury R, Chen MF; Tokyo Guidelines Revision Committee. New diagnostic criteria and severity assessment of acute cholangitis in revised Tokyo Guidelines. *J Hepatobiliary Pancreat Sci* 2012; **19**: 548-556 [PMID: [22825491](https://pubmed.ncbi.nlm.nih.gov/22825491/) DOI: [10.1007/s00534-012-0537-3](https://doi.org/10.1007/s00534-012-0537-3)]
- 24 Yokoe M, Hata J, Takada T, Strasberg SM, Asbun HJ, Wakabayashi G, Kozaka K, Endo I, Deziel DJ, Miura F, Okamoto K, Hwang TL, Huang WS, Ker CG, Chen MF, Han HS, Yoon YS, Choi IS, Yoon DS, Noguchi Y, Shikata S, Ukai T, Higuchi R, Gabata T, Mori Y, Iwashita Y, Hibi T, Jagannath P, Jonas E, Liau KH, Dervenis C, Gouma DJ, Cherqui D, Belli G, Garden OJ, Giménez ME, de Santibañes E, Suzuki K, Umezawa A, Supe AN, Pitt HA, Singh H, Chan ACW, Lau WY, Teoh AYW, Honda G, Sugioka A, Asai K, Gomi H, Itoi T, Kiriyaama S, Yoshida M, Mayumi T, Matsumura N, Tokumura H, Kitano S, Hirata K, Inui K, Sumiyama Y, Yamamoto M. Tokyo Guidelines 2018: diagnostic criteria and severity grading of acute cholecystitis (with videos). *J Hepatobiliary Pancreat Sci* 2018; **25**: 41-54 [PMID: [29032636](https://pubmed.ncbi.nlm.nih.gov/29032636/) DOI: [10.1002/jhbp.515](https://doi.org/10.1002/jhbp.515)]
- 25 Kiriyaama S, Kozaka K, Takada T, Strasberg SM, Pitt HA, Gabata T, Hata J, Liau KH, Miura F, Horiguchi A, Liu KH, Su CH, Wada K, Jagannath P, Itoi T, Gouma DJ, Mori Y, Mukai S, Giménez ME, Huang WS, Kim MH, Okamoto K, Belli G, Dervenis C, Chan ACW, Lau

- WY, Endo I, Gomi H, Yoshida M, Mayumi T, Baron TH, de Santibañes E, Teoh AYB, Hwang TL, Ker CG, Chen MF, Han HS, Yoon YS, Choi IS, Yoon DS, Higuchi R, Kitano S, Inomata M, Deziel DJ, Jonas E, Hirata K, Sumiyama Y, Inui K, Yamamoto M. Tokyo Guidelines 2018: diagnostic criteria and severity grading of acute cholangitis (with videos). *J Hepatobiliary Pancreat Sci* 2018; **25**: 17-30 [PMID: 29032610 DOI: 10.1002/jhbp.512]
- 26 **Tomida S**, Abei M, Yamaguchi T, Matsuzaki Y, Shoda J, Tanaka N, Osuga T. Long-term ursodeoxycholic acid therapy is associated with reduced risk of biliary pain and acute cholecystitis in patients with gallbladder stones: a cohort analysis. *Hepatology* 1999; **30**: 6-13 [PMID: 10385632 DOI: 10.1002/hep.510300108]
- 27 **Venneman NG**, Besselink MG, Keulemans YC, Vanberge-Henegouwen GP, Boermeester MA, Broeders IA, Go PM, van Erpecum KJ. Ursodeoxycholic acid exerts no beneficial effect in patients with symptomatic gallstones awaiting cholecystectomy. *Hepatology* 2006; **43**: 1276-1283 [PMID: 16729326 DOI: 10.1002/hep.21182]
- 28 **Randi G**, Franceschi S, La Vecchia C. Gallbladder cancer worldwide: geographical distribution and risk factors. *Int J Cancer* 2006; **118**: 1591-1602 [PMID: 16397865 DOI: 10.1002/ijc.21683]
- 29 **Yagyu K**, Lin Y, Obata Y, Kikuchi S, Ishibashi T, Kurosawa M, Inaba Y, Tamakoshi A; JACC Study Group. Bowel movement frequency, medical history and the risk of gallbladder cancer death: a cohort study in Japan. *Cancer Sci* 2004; **95**: 674-678 [PMID: 15298731 DOI: 10.1111/j.1349-7006.2004.tb03328.x]
- 30 **Miyazaki M**, Takada T, Miyakawa S, Tsukada K, Nagino M, Kondo S, Furuse J, Saito H, Tsuyuguchi T, Chijiwa K, Kimura F, Yoshitomi H, Nozawa S, Yoshida M, Wada K, Amano H, Miura F; Japanese Association of Biliary Surgery; Japanese Society of Hepato-Pancreatic Surgery; Japan Society of Clinical Oncology. Risk factors for biliary tract and ampullary carcinomas and prophylactic surgery for these factors. *J Hepatobiliary Pancreat Surg* 2008; **15**: 15-24 [PMID: 18274840 DOI: 10.1007/s00534-007-1276-8]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

