

World Journal of *Clinical Pediatrics*

Quarterly Volume 13 Number 1 March 9, 2024



EDITORIAL

Jain S. 'Prediabetes' as a practical distinctive window for workable fruitful wonders: Prevention and progression alert as advanced professionalism. *World J Clin Pediatr* 2024; 13(1): 89201 [DOI: [10.5409/wjcp.v13.i1.89201](https://doi.org/10.5409/wjcp.v13.i1.89201)]

MINIREVIEWS

Hudson AS, Wahbeh GT, Zheng HB. Imaging and endoscopic tools in pediatric inflammatory bowel disease: What's new? *World J Clin Pediatr* 2024; 13(1): 89091 [DOI: [10.5409/wjcp.v13.i1.89091](https://doi.org/10.5409/wjcp.v13.i1.89091)]

Elghoudi A, Zourob D, Al Atrash E, Alshamsi F, Alkatheeri M, Narchi H, Bitar R. Evolving strategies: Enhancements in managing eosinophilic esophagitis in pediatric patients. *World J Clin Pediatr* 2024; 13(1): 89580 [DOI: [10.5409/wjcp.v13.i1.89580](https://doi.org/10.5409/wjcp.v13.i1.89580)]

ORIGINAL ARTICLE**Case Control Study**

Murillo Galvis M, Ortegón Ochoa S, Plata García CE, Valderrama Junca MP, Inga Ceballos DA, Mora Gómez DM, Granados CM, Rondón M. Exclusive breastfeeding greater than 50%, success of education in a university hospital in Bogotá: Case-control study. *World J Clin Pediatr* 2024; 13(1): 87713 [DOI: [10.5409/wjcp.v13.i1.87713](https://doi.org/10.5409/wjcp.v13.i1.87713)]

Atef Abdelsattar Ibrahim H, Mohsen M, Salep Aziz Hanna B, Mahmoud D, Mohamed Abdelhamid El-Khashab K. Childhood asthma biomarkers including zinc: An exploratory cross-sectional study. *World J Clin Pediatr* 2024; 13(1): 87866 [DOI: [10.5409/wjcp.v13.i1.87866](https://doi.org/10.5409/wjcp.v13.i1.87866)]

Metwali WA, Elmashad AM, Hazzaa SME, Al-Beltagi M, Hamza MB. Salivary C-reactive protein and mean platelet volume as possible diagnostic markers for late-onset neonatal pneumonia. *World J Clin Pediatr* 2024; 13(1): 88645 [DOI: [10.5409/wjcp.v13.i1.88645](https://doi.org/10.5409/wjcp.v13.i1.88645)]

Kalashnikova E, Isupova E, Gaidar E, Sorokina L, Kaneva M, Masalova V, Dubko M, Kornishina T, Lubimova N, Kuchinskaya E, Chikova I, Raupov R, Kalashnikova O, Kostik M. BCD020 rituximab bioanalogue compared to standard treatment in juvenile systemic lupus erythematosus: The data of 12 months case-control study. *World J Clin Pediatr* 2024; 13(1): 89049 [DOI: [10.5409/wjcp.v13.i1.89049](https://doi.org/10.5409/wjcp.v13.i1.89049)]

Retrospective Cohort Study

Isa HM, Isa AJ, Alnasheet MA, Mansoor MM. Fever assessment in children under five: Are we following the guidelines? *World J Clin Pediatr* 2024; 13(1): 88864 [DOI: [10.5409/wjcp.v13.i1.88864](https://doi.org/10.5409/wjcp.v13.i1.88864)]

Belozherov KE, Solomatina NM, Isupova EA, Kuznetsova AA, Kostik MM. Systemic juvenile idiopathic arthritis-associated lung disease: A retrospective cohort study. *World J Clin Pediatr* 2024; 13(1): 88912 [DOI: [10.5409/wjcp.v13.i1.88912](https://doi.org/10.5409/wjcp.v13.i1.88912)]

Observational Study

Suksantilerd S, Thawatchai R, Rungrojjananon N. Prevalence of vitamin D deficiency in exclusively breastfed infants at Charoenkrung Pracharak Hospital. *World J Clin Pediatr* 2024; 13(1): 86693 [DOI: [10.5409/wjcp.v13.i1.86693](https://doi.org/10.5409/wjcp.v13.i1.86693)]

Zein MM, Arafa N, El-Shabrawi MHF, El-Koofy NM. Effect of nutrition-related infodemics and social media on maternal experience: A nationwide survey in a low/middle income country. *World J Clin Pediatr* 2024; 13(1): 89139 [DOI: [10.5409/wjcp.v13.i1.89139](https://doi.org/10.5409/wjcp.v13.i1.89139)]

Manokaran K, Spaan J, Cataldo G, Lyons C, Mitchell PD, Sare T, Zimmerman LA, Rufo PA. Inpatient management of iron deficiency anemia in pediatric patients with inflammatory bowel disease: A single center experience. *World J Clin Pediatr* 2024; 13(1): 89318 [DOI: [10.5409/wjcp.v13.i1.89318](https://doi.org/10.5409/wjcp.v13.i1.89318)]

El Mouzan M, Al Sarkhy A, Assiri A. Gut microbiota predicts the diagnosis of ulcerative colitis in Saudi children. *World J Clin Pediatr* 2024; 13(1): 90755 [DOI: [10.5409/wjcp.v13.i1.90755](https://doi.org/10.5409/wjcp.v13.i1.90755)]

SYSTEMATIC REVIEWS

Al-Beltagi M, Saeed NK, Bediwy AS, Elbeltagi R, Hamza MB. Gastrointestinal tolerability of organic infant formula compared to traditional infant formula: A systematic review. *World J Clin Pediatr* 2024; 13(1): 88783 [DOI: [10.5409/wjcp.v13.i1.88783](https://doi.org/10.5409/wjcp.v13.i1.88783)]

Avendaño-Vásquez CJ, Villamizar-Osorio ML, Niño-Peñaranda CJ, Medellín-Olaya J, Reina-Gamba NC. Sociodemographic determinants associated with breastfeeding in term infants with low birth weight in Latin American countries. *World J Clin Pediatr* 2024; 13(1): 89086 [DOI: [10.5409/wjcp.v13.i1.89086](https://doi.org/10.5409/wjcp.v13.i1.89086)]

LETTER TO THE EDITOR

Grauslund AC, Lindkvist EB, Thorsen SU, Ballegaard S, Faber J, Svensson J, Berg AK. Pressure pain sensitivity: A new stress measure in children and adolescents with type 1 diabetes? *World J Clin Pediatr* 2024; 13(1): 89619 [DOI: [10.5409/wjcp.v13.i1.89619](https://doi.org/10.5409/wjcp.v13.i1.89619)]

ABOUT COVER

Peer Reviewer of *World Journal of Clinical Pediatrics*, Fateen Ata, MSc, MBBS, ABHS (Internal Medicine) Academic Editor, Academic Fellow, Academic Research, Doctor, Endocrinology, Hamad Medical Corporation, Qatar. docfateenata@gmail.com

AIMS AND SCOPE

The primary aim of the *World Journal of Clinical Pediatrics (WJCP, World J Clin Pediatr)* is to provide scholars and readers from various fields of pediatrics with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCP mainly publishes articles reporting research results and findings obtained in the field of pediatrics and covering a wide range of topics including anesthesiology, cardiology, endocrinology, gastroenterology, hematology, immunology, infections and infectious diseases, medical imaging, neonatology, nephrology, neurosurgery, nursing medicine, perinatology, pharmacology, respiratory medicine, and urology.

INDEXING/ABSTRACTING

The *WJCP* is now abstracted and indexed in PubMed, PubMed Central, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The *WJCP*'s CiteScore for 2022 is 1.7 and Scopus CiteScore rank 2022: Pediatrics, perinatology and child health is 176/306.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Yu-Qing Zhao*; Production Department Director: *Xiang Li*; Cover Editor: *Xu Guo*.

NAME OF JOURNAL

World Journal of Clinical Pediatrics

ISSN

ISSN 2219-2808 (online)

LAUNCH DATE

June 8, 2012

FREQUENCY

Quarterly

EDITORS-IN-CHIEF

Toru Watanabe, Elena Daniela Serban, Surjit Singh, Consolato M Sergi

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2219-2808/editorialboard.htm>

PUBLICATION DATE

March 9, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Case Control Study

Salivary C-reactive protein and mean platelet volume as possible diagnostic markers for late-onset neonatal pneumonia

Wafaa Ahmed Metwali, Abdelrahman Mohamed Elmashad, Sahar Mohey Eldin Hazzaa, Mohammed Al-Beltagi, Mohamed Basiony Hamza

Specialty type: Pediatrics

Provenance and peer review:

Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): B

Grade C (Good): 0

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Rigante D, Italy

Received: October 3, 2023

Peer-review started: October 3, 2023

First decision: November 2, 2023

Revised: November 3, 2023

Accepted: December 11, 2023

Article in press: December 11, 2023

Published online: March 9, 2024



Wafaa Ahmed Metwali, Abdelrahman Mohamed Elmashad, Mohammed Al-Beltagi, Mohamed Basiony Hamza, Department of Pediatric, Faculty of Medicine, Tanta University, Tanta 31511, Algharbia, Egypt

Sahar Mohey Eldin Hazzaa, Department of Clinical Pathology, Faculty of Medicine, Tanta University, Tanta 31511, Algharbia, Egypt

Mohammed Al-Beltagi, Department of Pediatric, University Medical Center, Dr. Suliaman Al Habib Medical Group, Manama 26671, Manama, Bahrain

Mohammed Al-Beltagi, Department of Pediatric, University Medical Center, King Abdulla Medical City, Arabian Gulf University, Manama 26671, Manama, Bahrain

Corresponding author: Mohammed Al-Beltagi, MBChB, MD, PhD, Academic Editor, Chairman, Professor, Research Scientist, Department of Pediatric, Faculty of Medicine, Tanta University, Al-Bahr Street, Tanta 31511, Algharbia, Egypt. mbelrem@hotmail.com

Abstract

BACKGROUND

Neonatal sepsis, a formidable threat to newborns, is a leading cause of neonatal mortality, with late-onset sepsis manifesting after 72 hours post-birth being particularly concerning. Pneumonia, a prevalent sepsis presentation, poses a significant risk, especially during the neonatal phase when lung defenses are compromised. Accurate diagnosis of pneumonia is imperative for timely and effective interventions. Saliva, a minimally invasive diagnostic medium, holds great promise for evaluating infections, especially in infants.

AIM

To investigate the potential of serum C-reactive protein (CRP), salivary CRP (sCRP), and mean platelet volume (MPV) as diagnostic markers for late-onset neonatal pneumonia (LONP).

METHODS

Eighty full-term neonates were systematically examined, considering anthropometric measurements, clinical manifestations, radiology findings, and essential biomarkers, including serum CRP, sCRP, and MPV.

RESULTS

The study reveals noteworthy distinctions in serum CRP levels, MPV, and the serum CRP/MPV ratio between neonates with LONP and healthy controls. MPV exhibited a robust discriminatory ability [area under the curve (AUC) = 0.87] with high sensitivity and specificity at a cutoff value of > 8.8. Correlations between serum CRP, sCRP, and MPV were also identified. Notably, sCRP demonstrated excellent predictive value for serum CRP levels (AUC = 0.89), underscoring its potential as a diagnostic tool.

CONCLUSION

This study underscores the diagnostic promise of salivary and serum biomarkers, specifically MPV and CRP, in identifying and predicting LONP among neonates. These findings advocate for further research to validate their clinical utility in larger neonatal cohorts.

Key Words: Neonatal sepsis; Late-onset pneumonia; Salivary C-reactive protein; Mean platelet volume; Diagnostic markers; Newborn infections

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: This prospective study explores the potential of salivary C-reactive protein (CRP) (sCRP) and mean platelet volume (MPV) as diagnostic markers for late-onset neonatal pneumonia (LONP). Analyzing 80 neonates, significant differences in serum CRP levels, MPV, and the serum CRP/MPV ratio were observed between LONP cases and healthy controls. MPV demonstrated strong discriminatory ability with high sensitivity and specificity at a cutoff value of > 8.8. sCRP displayed notable predictive value for serum CRP levels. These findings highlight the diagnostic potential of salivary and serum biomarkers in identifying and predicting LONP among neonates.

Citation: Metwali WA, Elmashad AM, Hazzaa SME, Al-Beltagi M, Hamza MB. Salivary C-reactive protein and mean platelet volume as possible diagnostic markers for late-onset neonatal pneumonia. *World J Clin Pediatr* 2024; 13(1): 88645

URL: <https://www.wjgnet.com/2219-2808/full/v13/i1/88645.htm>

DOI: <https://dx.doi.org/10.5409/wjcp.v13.i1.88645>

INTRODUCTION

Neonatal sepsis is a serious infection in newborns with a very high risk of neonatal death and occupies the third rank among the causes of neonatal death[1]. It can manifest in various forms, like septicemia, pneumonia, meningitis, osteomyelitis, arthritis, and urinary tract infections. Late-onset sepsis occurs after 72 hours of birth and is a significant cause of infant mortality[2]. Despite medical advances, diagnosing and managing neonatal infections remains challenging. Childhood mortality due to pneumonia carries the highest risk during the neonatal phase since the fetus and neonate have compromised lung defenses, making them more prone to infections[3]. Neonatal pneumonia can be classified according to its onset into early-onset (within the 1st wk of life) and late-onset (onset of symptoms after the 1st wk of life within the first 28 d). Late-onset neonatal pneumonia is further classified into hospital- or community-acquired. The community-acquired neonatal pneumonia occurs in term and near-term neonates who were discharged home after the initial birth hospitalization. Hospital-acquired late-onset pneumonia (LOP) occurs in newborns who remain hospitalized since birth (*e.g.*, preterm infants)[4,5].

It is crucial to accurately diagnose pneumonia to assess the disease's impact, implement suitable preventive or treatment measures, and develop more efficient interventions[6]. Saliva has been found to have excellent potential as a diagnostic fluid over the years. It's easy and non-invasive collection method makes it the most attractive diagnostic medium to examine vulnerable populations such as infants, toddlers and children[7]. C-reactive protein (CRP), which is a major acute phase protein, is a member of the pentraxin family and plays a central role in innate and adaptive immunity. It takes 10-12 h for CRP to rise significantly after the onset of an infection[8]. Since CRP shows an increase in several conditions, it is better to use it in combination with other biomarkers.

Platelets, small non-nucleated cells derived from precursor megakaryocytes, have multiple functions and play a vital role in hemostasis by forming blood clots. They are a natural source of growth factors, including platelet-derived growth factor and transforming growth factor- β , which are essential for connective tissue repair and regeneration[9]. Platelet-rich plasma has been used to increase the concentration of these growth factors and aid wound healing. Thrombopoiesis, the production of platelets, is driven by thrombopoietin and several transcription factors. In inflammatory states, interleukin-6 enhances the process of proplatelet formation by increasing thrombopoietin levels[10]. Mean platelet volume (MPV) is one of the hemogram parameters that is affected by many inflammatory conditions. In neonates, MPV can predict the development of sepsis and its severity[11]. Therefore, the combined measurement of CRP and MPV can be used to diagnose bacterial *vs* viral pneumonia and predict its complications[12]. This study aims to assess the effectiveness of salivary CRP (sCRP) and MPV in identifying LOP in newborns.

MATERIALS AND METHODS

The present research was a prospective case-control study conducted on eighty full-term neonates recruited serially from the Neonatal Intensive Care Unit (NICU) and Clinic, Pediatric Department, the tertiary care hospital of Tanta University between June 2021 and May 2022, to evaluate the usefulness of sCRP and MPV in identifying Late Onset Neonatal Pneumonia (LONP). The recruited neonates were divided into two comparable groups: Group I included neonates who developed late-onset neonatal pneumonia (who developed pneumonia after 3rd d and before 28th d of life). Group II included healthy neonates with no clinical manifestation of infection or other systemic diseases.

We included full-term neonates (gestational age ≥ 37 wk and birth weight ≥ 2.5 kg) with post-natal age between 7 and 28 d with clinical suspension of LOP. All eligible neonates underwent comprehensive assessments of their prenatal, perinatal, and postnatal history, thorough clinical examinations, a complete blood cell count including differential, evaluation of CRP levels, urine analysis and culture, blood culture, cerebrospinal fluid analysis and culture, and relevant infection markers. Blood gases, chest imaging (plain X-ray and/or ultrasonography), echocardiography, and abdominal X-ray were conducted based on specific clinical indications. We followed Strengthening the Reporting of Observational Studies in Epidemiology for Newborn Infection for reporting neonatal infection[13] and Gerdes' sepsis screen (> 2) to screen for neonatal sepsis, including pneumonia[14]. Pneumonia was suspected in the presence of fever or temperature instability, irritability, lethargy, feeding difficulty, apnea, or respiratory distress. Other systemic manifestations, such as hepatomegaly, abdominal distention, convulsion, hypotonia, hemodynamic instability, and bleeding diathesis, were considered general manifestations of neonatal sepsis. We classified the patients as mild [degree of respiratory distress (RD) 1: Tachypnea > 60 /min and flaring nostrils], moderate (RD 2: RD2 + intercostal and subcostal retractions), severe (RD2 + expiratory grunting), and advanced (RD 3 + central cyanosis); according to the degree of respiratory distress. Chest X-ray findings varied from normal chest X-ray to localized or diffuse alveolar densities, reticular opacities, homogenous ground glass opacities, and dense bilateral air space-filling process with air bronchograms. Any pneumonia complication findings were also recorded, such as interstitial emphysema, pleural effusion, pneumomediastinum, or pneumothorax.

We excluded premature infants and neonates with inflammation other than pneumonia, congenital heart conditions, hypoxic-ischemic encephalopathy, liver or kidney issues, hereditary coagulopathies, or any other systemic disorders unrelated to pneumonia that could impact CRP or platelet size levels. We also excluded neonates exposed to antibiotics before admission, neonates younger than 7 d, or infants older than 28 d of life. According to NICU protocol, all children with suspected pneumonia receive the appropriate management. All parents, guardians, or next of kin signed informed consent for the minors to participate in this study. The Institutional Ethical and Research Review Board of the Faculty of Medicine, Tanta University, approved the study.

Laboratory investigations included salivary and serum CRP measurement and MPV measurement. We collected salivary samples just before feeding to avoid milk contamination. With gentle handling of the baby, we stimulated the saliva secretion by allowing the baby to suck on a clean, sterilized pacifier for a few minutes. The head of the baby was elevated to allow the saliva to collect on the floor of the mouth, under the neonates' tongues, for accessible collection. The saliva samples were collected using a one-ml syringe without a needle with the suction pressure applied manually for about 10–15 s, collecting about 0.5 mL of saliva. Then, the samples were transferred to sterile polypropylene tubes to avoid contamination and stored at -20 °C until CRP analysis and measuring using ELISA. Serum CRP levels were determined using a fully automated auto-analyzer Cobas c501 (Roche Diagnostics, Mannheim, Germany).

A peripheral blood sample was collected just before feeding (to avoid the effect of feeding on the platelet volume) into a clean, sterile EDTA vacutainer tube to measure MPV value. The sample was handled gently without unnecessary agitation to minimize platelet activation and analyzed using an automated blood cell counter (Cell-Dyn 3700, Abbott Laboratories, IL, United States) within 60 min of collection to avoid platelet swelling and pseudo increase in MPV value. The analyzer calculates the MPV by dividing the total platelet volume by the number of platelets in the blood sample.

Statistical analysis

We used the Power and Precision V3 program (<http://www.Power-Analysis.com>, Englewood, New Jersey) to determine the study's power level. The collected data were organized, tabulated, and subjected to statistical analysis using the SPSS version 20 (SPSS, Chicago, IL, United States) to determine the sensitivity, specificity, and predictive value of MPV, sCRP, serum CRP, and the serum CRP/MPV ratio for diagnosing LONP cases. We used the Shapiro-Wilk test to test the normality of data distribution. Mann-Whitney *U*-test assessed the differences between groups regarding nonparametric quantitative data. Receiver operating characteristic curves were used to identify optimal cutoff values for differentiating patients with LONP from healthy controls. We used the mean and standard deviation to characterize the quantitative data. We considered the findings to be statistically significant when the *P* was < 0.05 .

RESULTS

In this study, we compared two groups of neonates: One with LOP comprising 40 neonates and a control group of 40 healthy neonates. Our analysis in Table 1 revealed no significant differences in sex distribution, age, mode of delivery, Apgar score, and the presence or absence of maternal illness during pregnancy between the neonates with LOP and the control group. However, notable variations in anthropometric measurements were observed, with the neonates with LOP neonates exhibiting lower weight, length, and head circumference than the control group. Additionally, the LOP group demonstrated substantially elevated levels of serum CRP, MPV, and the CRP/MPV ratio, indicating the potential

Table 1 Demographic and laboratory findings of neonates with late-onset pneumonia and the control group

Parameter		Neonates with LOP, n = 40	Control group, n = 40	P value
Sex	M	21	22	> 0.05
	F	19	18	> 0.05
	M:F	1.1:1	1.2:1	> 0.05
Age in d		15 ± 4.6	14 ± 5	> 0.05
Anthropometric measurements at admission	Weight in g	3454 ± 360	3728 ± 316	< 0.0001 ^a
	Length in cm	51.42 ± 1.0	51.87 ± 0.57	< 0.05 ^a
	Head circumference	36.21 ± 0.8	36.7 ± 0.60	< 0.05 ^a
Type of delivery	Normal	14 (35%)	16 (40%)	> 0.05
	CS	26 (65%)	24 (60%)	> 0.05
APGAR score	At 1 min	7.1 ± 2.5	7.4 ± 2.3	> 0.05
	At 5 min	8.5 ± 3.2	9 ± 2.9	> 0.05
Maternal illness	Non	31 (77.5%)	32 (80%)	> 0.05
	DM	3 (7.5%)	2 (5%)	> 0.05
	hypertension	2 (5%)	5 (12.5%)	> 0.05
	UTI	4 (10%)	3 (7.5%)	> 0.05
Serum CRP in mg/L		38.58 ± 24.9	3.60 ± 2.25	< 0.0001 ^a
sCRP in mg/L		6.17 ± 3.38	3.07 ± 1.24	< 0.0001 ^a
MPV		9.99 ± 0.94	8.42 ± 0.83	< 0.0001 ^a
Serum CRP/MPV		3.86 ± 2.29	0.42 ± 0.39	< 0.0001 ^a

^aP < 0.05.

Data are n, n (%), or mean ± SD. SCRP: C-reactive protein; CS: Caesarean section; DM: Diabetes mellitus; F: Female; LOP: Late-onset pneumonia; M: Male; MPV: Mean platelet volume; sCRP: Salivary C-reactive protein; UTI: Urinary tract infection.

diagnostic value of these markers for LOP among neonates.

Table 2 shows LOP's clinical and radiological features in a cohort of 40 neonates. Clinical manifestations were prominent, with 65% and 70% of neonates exhibiting fever and cough, respectively. Various degrees of respiratory distress were observed, ranging from RD 1 to RD 4 in 10% to 40% of neonates. Abnormal auscultatory findings were prevalent, including decreased air entry (82.5%) and fine crepitations (87.5%). Radiologically, pneumonic patches with an air bronchogram were the most common pattern (75%), followed by homogenous ground glass shadows (10.0%) and interstitial pneumonia (10.0%). A smaller percentage showed a complete white lung (2.5%), and complications were noted in 2.5% of cases. Oxygen support was crucial, with 77.5% requiring a nasal cannula, 12.5% supported by nasal continuous positive airway pressure, and 10% necessitating mechanical ventilation. These comprehensive findings offer valuable insights into LOP's clinical and radiological spectrum, enabling more precise diagnoses and tailored treatment approaches for affected neonates.

In addition to the clinical and radiological findings, the study evaluated the diagnostic validity of key markers in discriminating patients from controls, as shown in Table 3. The area under the curve (AUC), sensitivity, specificity, positive predictive values (PPV), and negative predictive values (NPV) were calculated for MPV, Serum CRP, and sCRP. MPV exhibited a high discriminatory ability (AUC = 0.87) with a sensitivity of 86.67% and specificity of 80.0% at a cutoff value of > 8.8. Serum CRP also showed good discriminative power (AUC = 0.81), with a sensitivity of 76.67% and specificity of 60.0% at a cutoff value of > 6. Similarly, sCRP demonstrated notable discriminatory ability (AUC = 0.80) with a sensitivity of 76.67% and specificity of 83.33% at a cutoff value of > 3.5. Furthermore, correlations were explored between serum CRP, sCRP, and MPV in Table 4, revealing significant positive correlations. The study also assessed the diagnostic validity of sCRP in predicting serum CRP levels, demonstrating a high AUC of 0.89 with a sensitivity of 91.3% and specificity of 71.4% at a cutoff value of >3.2, as shown in Table 5. These comprehensive assessments highlight the potential diagnostic utility of MPV and CRP markers, both serum and salivary, in discriminating and predicting disease severity in neonates with LOP.

Table 2 Clinical and radiological findings of neonates with late onset pneumonia

Finding	Number of 40 total	Percentage (%)
Clinical findings		
Fever	26	65.0
Cough	28	70.0
Degree of respiratory distress		
1	7	17.5
2	16	40.0
3	13	32.5
4	4	10.0
Decreases air entry	33	82.5
Fine crepitations	35	87.5
Radiological findings		
Homogenous ground glass shadow	4	10.0
Pneumonic patches and air bronchogram	30	75.0
Interstitial pneumonia	4	10.0
Complete white lung	1	2.5
Complications	1	2.5
Need for oxygen support		
Nasal canula	31	77.5
nCPAP	5	12.5
Mechanical ventilation	4	10.0

nCPAP: Nasal continuous positive airway pressure.

Table 3 Validity (area under the curve, sensitivity, specificity) of mean platelet volume, salivary C-reactive protein, serum C-reactive protein, and serum C-reactive protein/mean platelet volume to discriminate patients from controls

Parameter	AUC	95%CI	P value	Cut-off value	Sensitivity	Specificity	PPV	NPV
MPV	0.87	0.77–0.97	< 0.001 ^a	> 8.8	86.67	80.0	81.2	85.7
Serum CRP	0.81	0.70–0.92	< 0.001 ^a	> 6 mg/dL	76.67	60.0	65.7	72.0
sCRP	0.80	0.68–0.92	< 0.001 ^a	> 3.5 mg/dL	76.67	83.33	82.1	78.1

^a*P* < 0.05.

AUC: Area under the curve; CI: Confidence interval; CRP: C-reactive protein; MPV: Mean platelet volume; NPV: Negative predictive value; PPV: Positive predictive value; sCRP: Salivary C-reactive protein.

DISCUSSION

LOP in neonates is a critical condition, presenting unique challenges in diagnosis and management that demands a thorough understanding of its clinical, radiological, and biochemical features to enhance diagnostic precision, facilitate tailored therapeutic interventions and improve clinical outcomes[15]. This study aimed to comprehensively explore these aspects and evaluate the diagnostic potential of specific biomarkers in discriminating between neonates with LOP and healthy controls.

The clinical manifestations observed in neonates with LOP encompassed fever, cough, varying degrees of respiratory distress, decreased air entry and fine crepitations. These clinical findings align with the well-documented respiratory symptoms associated with pneumonia. These findings agreed with Omran *et al*[12], who studied 35 full-term neonates diagnosed with LOP and 35 controls. They found fine crepitation in 32 (91.4%), decreased air entrance in 24 (74.3%) and intercostal retractions in 25 (71.4%). Furthermore, the varying degrees of respiratory distress, observed as thoracic retractions in Omran *et al*'s[12] study and as RD1 to RD4 in our study, indicate the diverse respiratory involvement in

Table 4 Correlation of serum C-reactive protein with each salivary C-reactive protein and mean platelet volume, neonates with late-onset pneumonia (n = 40)

Parameter	r value	P value
Serum CRP vs sCRP	0.59	0.001 ^a
Serum CRP vs MPV	0.66	< 0.001 ^a
Salivary CRP vs MPV	0.54	0.01 ^a

^aP < 0.05.

CRP: Salivary C-reactive protein; MPV: Mean platelet volume; sCRP: Salivary C-reactive protein.

Table 5 Diagnostic validity (area under the curve, sensitivity, specificity) for salivary C-reactive protein to predict serum C-reactive protein > 6 (n = 31) from serum C-reactive protein ≤ 6 (n = 9) in the patient group (n = 40)

Parameter	AUC	95%CI	P value	Cut-off Value	Sensitivity	Specificity	PPV	NPV
sCRP	0.89	0.77-1.1	< 0.001 ^a	> 3.2	91.3	71.4	91.3	71.4

^aP < 0.05.

AUC: Area under the curve; CI: Confidence interval; NPV: negative predictive value; PPV: Positive predictive value; sCRP: Salivary C-reactive protein.

LOP. Additionally, radiological examinations provided valuable insights into the lung pathologies present in LOP, notably pneumonic patches with air bronchograms, ground glass shadows, and interstitial pneumonia. These findings agree with that of Haney *et al*[16], who found that bilateral alveolar densities were the most commonly identified X-ray abnormality, noted in 77% of cases. One-third of their patients had typically dense and extensive alveolar changes with frequent air bronchograms. Our findings resonate with existing literature, further validating the robustness of our study.

Intriguingly, our study delved deeper into the potential diagnostic value of biomarkers. MPV, serum CRP, and sCRP were identified as promising, readily available candidates for discriminating patients from controls. MPV, a marker often associated with inflammatory conditions, demonstrated high discriminative ability (AUC = 0.87) with significant sensitivity and specificity, suggesting its potential as a diagnostic tool for LOP.

Platelets play a significant role in neonatal sepsis-induced coagulopathy. During systemic inflammation, P-selectin is expressed on the surface of platelets, enhancing platelet adherence to leukocytes, platelet aggregation, and expression of tissue factor on monocytes[17]. MPV is considered a marker of platelet function and activation, associated with larger and more reactive platelets. During inflammatory states, platelets become activated, leading to an increase in MPV. This inflammation observed in LOP triggers platelet activation, resulting in larger platelets (higher MPV). Therefore, MPV could be used as a marker that indicates systemic inflammation and infection, including LOP[18]. It is being studied in various conditions in adults and children, including acute coronary syndrome and acute appendicitis[19,20]. In the current study, MPV was shown to have a high level of discriminative ability (AUC = 0.87), with a sensitivity of 86.67% and specificity of 80.0% at a cutoff value of > 8.8 for neonates with LOP. The high AUC score indicates that MPV is an excellent diagnostic tool for distinguishing neonates with LOP from healthy controls. An AUC of 0.87 signifies a strong ability to correctly classify patients and controls based on MPV levels. In addition, MPV has a sensitivity of 86.67%, which means it accurately identified the majority (86.67%) of neonates with LOP, reducing false negatives. On the other hand, the specificity of 80.0% suggests that MPV effectively excluded a significant portion (80.0%) of healthy neonates, reducing false positives. With a cutoff value of > 8.8 for MPV, neonates with MPV levels exceeding this threshold are likelier to have LOP.

Our results agree with many previous works. Omran and colleagues found a significant difference in MPV levels between neonates suffering from pneumonia and those who didn't and established a noteworthy association between MPV and CRP in both serum and saliva[21]. We have confirmed these findings. MPV, with a cut-off value of 9.0 fl exhibited an excellent diagnostic accuracy of 80% in identifying infants with pneumonia. Similarly, Pamudji and Kardana [22] reported that an MPV of 7.44 fl had 80% sensitivity and 84.2% specificity in diagnosing neonatal sepsis. In addition, Wang *et al*[23] conducted a meta-analysis which found that MPV was significantly higher in patients with neonatal sepsis than in the control group, suggesting that MPV could be used as an early indicator for diagnosing neonatal sepsis in clinical practice. This can help clinicians make diagnostic decisions based on MPV levels. With a high AUC and its balanced sensitivity and specificity, MPV holds promise as a diagnostic marker for neonatal LOP. It may offer a relatively simple and cost-effective method to diagnose neonatal pneumonia[24]. However, further research and validation in larger and more diverse cohorts are needed to definitively establish MPV's diagnostic accuracy and clinical utility.

CRP is one of the most utilized biomarkers to monitor infection and inflammation in the pediatric and neonatal populations[25]. Serum CRP, a well-established inflammation marker, exhibited a substantial discriminative ability (AUC = 0.81) and meaningful sensitivity and specificity, as observed in the current study. These findings are consistent with previous research indicating the diagnostic significance of serum CRP in respiratory infections. Kumar *et al*[26] found higher overall serum CRP accuracy in diagnosing late-onset neonatal sepsis, ranging from 96.5% in proven sepsis to

99.1% in probable sepsis with a specificity of 85.3 %, using a CRP cut-off value of 5 mg/L. This suggests that CRP has a high diagnostic accuracy in identifying neonates at risk of sepsis. In addition, Omran *et al*[27] found a significant increase in serum CRP with a mean of 29.4 ± 13 mg/L neonates with late-onset sepsis.

Serum CRP also exhibited considerable discriminative ability, with notable sensitivity and specificity for neonatal infection, including pneumonia. Its diagnostic significance in respiratory infections aligns with the observed potential in discriminating LOP. A meta-analysis by Xiao *et al*[28] revealed that infants with pneumonia exhibited higher serum CRP levels than healthy infants. Therefore, serum CRP levels might be an independent diagnostic tool for pneumonia in children. In addition, Li and Chen[29] found a close correlation between higher serum CRP levels and the progression of neonatal pneumonia. Several studies have reported different cut-off values for serum CRP levels in diagnosing neonatal sepsis. These values range widely, from 1.5 to 20 mg/L, and are associated with varying sensitivities and specificities. For instance, sensitivity values range from 74% to 98%, while specificities range from 71% to 94%, whether using a single measurement at least 12 h after the onset of symptoms or serial CRP determinations[30-33].

The detection of CRP in saliva is a new and promising diagnostic method that has recently gained attention as an emerging biomarker. It shows potential in diagnosing various medical conditions, such as pneumonia. One of the significant advantages of using sCRP in diagnosing neonatal pneumonia is its non-invasive nature. Saliva collection is less intrusive and more feasible than obtaining blood samples, especially in neonates, where it can be challenging to draw blood[34,35]. Iyengar *et al*[36] conducted a study on sCRP detection and its usefulness in neonates. It is considered the first study to detect, quantify, and demonstrate that sCRP is a good measure of discrimination for clinically relevant serum CRP thresholds. The study included the most salivary samples obtained from neonates suffering from necrotizing enterocolitis or spontaneous intestinal perforation, infectious diseases, and post-operative monitoring. The median sCRP concentration was found to be 3.1 ng/mL, whereas the median serum CRP concentration was 106.1 mg/L[36]. Interestingly, we used sCRP in our study as a non-invasive alternative that displayed noteworthy discriminatory ability (AUC = 0.80) and strong correlations with serum CRP and MPV. Our results emphasize the potential of sCRP as a viable diagnostic marker in neonates with LOP and agree with many previous studies. A study conducted by Omran *et al*[27] found a result similar to ours. They observed a significant difference in the mean level of sCRP between septic neonates (12.0 ± 4.6 ng/L) and the control group (2.8 ± 1.2 ng/L). The sensitivity of sCRP was 94.3%, and specificity was 80% at a cut-off point of 3.48 ng/L[27]. Barekatin *et al*[25] also reported a significant increase in sCRP levels in neonates with sepsis compared to healthy controls. The AUC value was 0.63, with a sensitivity of 44.9%, specificity of 80%, PPV of 73.3%, NPV of 54.2%, and diagnostic accuracy of 61% at a cutoff of 4.55 ng/L[25].

The current study has shown a significant correlation ($r = 0.59$, $P < 0.001$) between serum CRP and sCRP levels in neonates with LOP. This suggests that sCRP levels reflect those in serum and can serve as a non-invasive diagnostic biomarker for neonatal LOP. sCRP can potentially be used as a proxy for serum CRP, indicating the systemic inflammatory response associated with pneumonia[37]. The positive correlation strengthens the case for considering sCRP as a reliable diagnostic tool for neonatal LOP. In a study by Iyengar *et al*[36], an sCRP concentration of 4.84 ng/L was found to have 64% sensitivity and 94% specificity for predicting a serum CRP of 5 mg/L. It was also found to have 54% sensitivity and 95% specificity for predicting a serum CRP of 10 mg/L. On the other hand, Tosson *et al*[38] found no significant correlation between sCRP and serum CRP levels in neonates with late-onset sepsis. To ensure consistent and accurate sCRP measurement across different studies, it is important to address various challenges that may affect its accuracy. For instance, screening for oral trauma and controlling the salivary flow rate is necessary to account for the salivary dilution effect. However, there are no available reliable strategies to make sCRP an accurate quantitative measure of serum CRP, which limits the use of point-of-care systemic inflammation testing[39].

The current study found a significant positive correlation ($r = 0.66$, $P < 0.001$) between serum CRP and MPV in neonates with LOP. This means that when serum CRP levels increase, MPV also tends to increase. The positive correlation between MPV and serum CRP suggests that MPV is associated with the inflammatory response and may reflect the level of inflammation present in pneumonia. In addition to this positive correlation between CRP and MPV in the current study, the serum CRP/MPV ratio in neonates with LOP (3.86 ± 2.29) was significantly higher than in the control group (0.42 ± 0.39) with a P value of less than 0.0001. The elevated serum CRP/MPV ratio in neonates with LOP signifies a higher inflammatory state. CRP is a key acute-phase protein that increases during inflammation. Concurrently, as a marker of platelet activation, MPV is influenced by the inflammatory response. The higher ratio suggests a greater inflammatory burden and platelet activation in neonates with LOP[40]. Using MPV as a diagnostic marker in combination with serum CRP could provide a more comprehensive understanding of the inflammatory status in neonates with LOP. Omran *et al*[12] found a significant increase of CRP/MPV in neonates with late-onset sepsis than in the control.

The current study reveals a statistically significant positive correlation between MPV and sCRP in neonates with LOP ($r = 0.54$, $P = 0.01$). This correlation indicates that as sCRP levels increase, MPV also tends to increase. Our findings agree with the work of Omran *et al*[27], who found a significant positive correlation between sCRP and MPV ($P < 0.001$). The positive correlation between MPV and sCRP suggests that both biomarkers are associated with the inflammatory response seen in LOP.

Elevated sCRP levels indicate systemic inflammation, and this correlation implies that higher levels of inflammation are associated with an increase in MPV. Understanding the correlation between sCRP and MPV is clinically valuable[41]. Monitoring both sCRP and MPV in neonates with LOP can provide complementary information about the severity of the inflammatory response[23,36]. If both sCRP and MPV are elevated, it could indicate a more pronounced inflammatory state, prompting close monitoring and possibly more aggressive treatment. The correlation between sCRP and MPV suggests that MPV, an easily measurable parameter, could serve as a supplementary diagnostic marker alongside sCRP. It may enhance the accuracy of diagnosing and monitoring the inflammatory status in neonates with LOP[42].

Moreover, the correlations between serum CRP, sCRP, and MPV underlined their interrelated nature, indicating the potential for a multi-marker approach in diagnosis and disease monitoring. The high AUC of sCRP in predicting serum CRP levels further advocates for its utility in disease assessment[38]. Understanding these correlations is critical for establishing the diagnostic potential of these markers. The strong correlations between serum CRP, sCRP, and MPV suggest their interlinked roles in reflecting the inflammatory response associated with LOP. Incorporating these markers collectively in diagnostic algorithms could enhance accuracy and offer a more holistic assessment of the inflammatory status in neonates with LOP, potentially leading to improved diagnostic and therapeutic approaches[43]. Further research should delve deeper into these correlations, exploring their clinical implications and potential for diagnostic integration.

Limitations of a study

It is crucial to provide possible limitation of the current study that limits the generalizability of its findings. The study's sample size is relatively modest, which may limit the generalizability of the findings to a broader neonatal population. The study being conducted in a single center may introduce institutional biases and limit the external validity of the results. The variability in the clinical presentation of LOP among neonates may introduce heterogeneity in the study population, potentially affecting the consistency and reliability of the observed clinical and radiological features. The study's cross-sectional design limits the assessment of temporal relationships and trends over time, which could be valuable for understanding the progression and outcomes of LOP in neonates. While MPV, serum CRP, and sCRP were evaluated as potential diagnostic markers, other relevant biomarkers that could contribute to a more comprehensive assessment were omitted. We also should consider the factors that could affect the sCRP levels, such as salivary flow rate, circadian rhythm, age, sex, type of salivary gland, salivary stimulation, feeding, and collection method. The study's exclusion of preterm neonates might limit the generalizability of the findings to the entire neonatal population, as preterm infants often have unique healthcare needs and susceptibilities. The study also employed specific radiological techniques (plain Chest X-rays); however, using advanced imaging modalities such as high-resolution computed tomography or other advanced imaging methods could have provided additional valuable insights into the lung pathology of neonates with LOP. In addition, the study's findings might be specific to a particular ethnic or geographical population. Caution should be exercised when generalizing the results to a more diverse or different population.

Suggestions for future research

Including a larger and more diverse cohort of neonates could provide a more comprehensive representation of LOP cases. Multicenter studies involving diverse healthcare settings could offer a more comprehensive view of LOP cases. We also must explore a broader array of biomarkers for a more accurate diagnosis.

CONCLUSION

This study offers a comprehensive understanding of the clinical, radiological, and biomarker profiles in neonates with LOP, aligning with the observations made by previous studies. The potential diagnostic utility of MPV, serum CRP, and sCRP was evident, opening new avenues for non-invasive diagnostic approaches. Integrating these biomarkers into clinical practice may enhance diagnostic accuracy and subsequently improve outcomes for neonates with LOP. Future research should focus on validating these findings in larger cohorts and exploring the prognostic implications of these biomarkers in guiding therapeutic strategies.

ARTICLE HIGHLIGHTS

Research background

Neonatal sepsis is a significant cause of neonatal mortality, and late-onset pneumonia (LOP) is a challenging form of sepsis to diagnose. Saliva has been identified as a potential diagnostic fluid for neonates. C-reactive protein (CRP) and mean platelet volume (MPV) are biomarkers that can indicate inflammation and are of interest in diagnosing neonatal infections.

Research motivation

The research is motivated by the need to improve the diagnosis of LOP in newborns, a serious condition that can lead to high mortality rates. Current diagnostic methods for neonatal infections, including pneumonia, can be challenging. The motivation is to find non-invasive and effective diagnostic tools that can aid in the early and accurate identification of LOP. Salivary CRP (sCRP) and MPV are being investigated as potential biomarkers to enhance the diagnosis and management of LOP, aiming to improve clinical outcomes for affected newborns.

Research objectives

We aimed to assess the diagnostic accuracy of sCRP and MPV biomarkers, analyzing their temporal trends, considering demographic factors, and exploring their clinical implications in diagnosing LOP in newborns.

Research methods

The study involved 80 full-term neonates divided into a group with LOP and a control group. Clinical assessments, blood tests, and imaging were conducted to diagnose LOP. Salivary and serum CRP levels, as well as MPV, were measured. Statistical analysis was performed to determine the diagnostic validity of these markers.

Research results

Neonates with LOP showed differences in weight, length, head circumference, serum CRP, MPV, and the CRP/MPV ratio compared to the control group. Clinical and radiological features of LOP were observed, including fever, cough, respiratory distress, and abnormal auscultatory findings. MPV, serum CRP, and sCRP exhibited good discriminative power for diagnosing LOP. Positive correlations were found between serum CRP, sCRP, and MPV.

Research conclusions

The study provides insights into the clinical, radiological, and biomarker profiles in neonates with LOP. MPV, serum CRP, and sCRP show potential for non-invasive diagnostic approaches. Integrating these biomarkers into clinical practice may enhance diagnostic accuracy and improve outcomes for neonates with LOP.

Research perspectives

Further research in neonatal LOP is needed to validate findings and assess generalizability across different populations and healthcare settings, investigate temporal trends and longitudinal studies, explore multi-marker approaches, assess ethnic and geographic variations, analyze the kinetics of sCRP and MPV, conduct studies on preterm neonates, compare diagnostic performance with other modalities, examine clinical implications, develop point-of-care testing methods, and investigate therapeutic implications. These research perspectives can lead to improved clinical practices and outcomes for affected neonates.

ACKNOWLEDGEMENTS

We thank the anonymous referees and editors for their valuable suggestions.

FOOTNOTES

Author contributions: El-Mashad MA and Hamza MB provided the research idea and initiated the study design; Metwali WA and El-Mashad MA collected the patients and their information; Hamza M and El-Mashad MA were responsible for statistical analysis; Metwali WA and Hazzaa HM were responsible for the technical part of the study; Also, they performed data analysis; Al-Biltagi M analyzed the data and wrote the final manuscript; all authors revised and agreed on the final version of the manuscript.

Institutional review board statement: We performed the study according to the latest version of Helsinki's Declaration. The Institutional Ethical and Research Review Board of the Faculty of Medicine, Tanta University, approved the study.

Informed consent statement: All parents, guardians, or next of kin signed informed consent for the minors to participate in this study.

Conflict-of-interest statement: All the authors declare that they have no potential nor real conflicts to disclose.

Data sharing statement: Data are available upon reasonable request.

STROBE statement: The authors have read the STROBE Statement – checklist of items, and the manuscript was prepared and revised according to the STROBE Statement – checklist of items.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Egypt

ORCID number: Mohammed Al-Beltagi 0000-0002-7761-9536.

S-Editor: Lin C

L-Editor: Filipodia

P-Editor: Cai YX

REFERENCES

- 1 **Mezgebu T**, Ossabo G, Zekiwas A, Mohammed H, Demisse Z. Neonatal sepsis and its associated factors among neonates admitted to the neonatal intensive care unit in Wachemo University Comprehensive Specialized Hospital, Southern Ethiopia, 2022. *Front Pediatr* 2023; **11**: 1184205 [PMID: 37465417 DOI: 10.3389/fped.2023.1184205]
- 2 **Yadav P**, Yadav SK. Progress in Diagnosis and Treatment of Neonatal Sepsis: A Review Article. *JNMA J Nepal Med Assoc* 2022; **60**: 318-324 [PMID: 35633256 DOI: 10.31729/jnma.7324]
- 3 **Hogden L**, Munger K, Duffek S. Neonatal Respiratory Distress. *SD Med* 2021; **74**: 28-35 [PMID: 33691054]
- 4 **Duke T**. Neonatal pneumonia in developing countries. *Arch Dis Child Fetal Neonatal Ed* 2005; **90**: F211-F219 [PMID: 15846010 DOI: 10.1136/adc.2003.048108]
- 5 **Nissen MD**. Congenital and neonatal pneumonia. *Paediatr Respir Rev* 2007; **8**: 195-203 [PMID: 17868917 DOI: 10.1016/j.prrv.2007.07.001]
- 6 **Cohen AL**, Hyde TB, Verani J, Watkins M. Integrating pneumonia prevention and treatment interventions with immunization services in resource-poor countries. *Bull World Health Organ* 2012; **90**: 289-294 [PMID: 22511825 DOI: 10.2471/BLT.11.094029]
- 7 **Shakeeb N**, Varkey P, Ajit A. Human Saliva as a Diagnostic Specimen for Early Detection of Inflammatory Biomarkers by Real-Time RT-PCR. *Inflammation* 2021; **44**: 1713-1723 [PMID: 34031776 DOI: 10.1007/s10753-021-01484-1]
- 8 **Sproston NR**, Ashworth JJ. Role of C-Reactive Protein at Sites of Inflammation and Infection. *Front Immunol* 2018; **9**: 754 [PMID: 29706967 DOI: 10.3389/fimmu.2018.00754]
- 9 **Noh JY**. Megakaryopoiesis and Platelet Biology: Roles of Transcription Factors and Emerging Clinical Implications. *Int J Mol Sci* 2021; **22** [PMID: 34502524 DOI: 10.3390/ijms22179615]
- 10 **Kaushansky K**. The molecular mechanisms that control thrombopoiesis. *J Clin Invest* 2005; **115**: 3339-3347 [PMID: 16322778 DOI: 10.1172/JCI26674]
- 11 **Catal F**, Tayman C, Tonbul A, Akça H, Kara S, Tatlı MM, Oztekin O, Bilici M. Mean platelet volume (MPV) may simply predict the severity of sepsis in preterm infants. *Clin Lab* 2014; **60**: 1193-1200 [PMID: 25134389 DOI: 10.7754/clin.lab.2013.130501]
- 12 **Omran A**, Sobh H, Abdalla MO, El-Sharkawy S, Rezk AR, Khashana A. Salivary and Serum Interleukin-10, C-Reactive Protein, Mean Platelet Volume, and CRP/MPV Ratio in the Diagnosis of Late-Onset Neonatal Sepsis in Full-Term Neonates. *J Immunol Res* 2021; **2021**: 4884537 [PMID: 34676267 DOI: 10.1155/2021/4884537]
- 13 **Fitchett EJA**, Seale AC, Vergnano S, Sharland M, Heath PT, Saha SK, Agarwal R, Ayede AI, Bhutta ZA, Black R, Bojang K, Campbell H, Cousens S, Darmstadt GL, Madhi SA, Meulen AS, Modi N, Patterson J, Qazi S, Schrag SJ, Stoll BJ, Wall SN, Wammanda RD, Lawn JE; SPRING (Strengthening Publications Reporting Infection in Newborns Globally) Group. Strengthening the Reporting of Observational Studies in Epidemiology for Newborn Infection (STROBE-NI): an extension of the STROBE statement for neonatal infection research. *Lancet Infect Dis* 2016; **16**: e202-e213 [PMID: 27633910 DOI: 10.1016/S1473-3099(16)30082-2]
- 14 **Perkins BS**, Brandon DH, Kahn DJ. Development of a Novel Assessment Tool and Code Sepsis Checklist for Neonatal Late-Onset Sepsis. *Adv Neonatal Care* 2022; **22**: 6-14 [PMID: 34334674 DOI: 10.1097/ANC.0000000000000896]
- 15 **Hooven TA**, Polin RA. Pneumonia. *Semin Fetal Neonatal Med* 2017; **22**: 206-213 [PMID: 28343909 DOI: 10.1016/j.siny.2017.03.002]
- 16 **Haney PJ**, Bohlman M, Sun CC. Radiographic findings in neonatal pneumonia. *AJR Am J Roentgenol* 1984; **143**: 23-26 [PMID: 6610324 DOI: 10.2214/ajr.143.1.23]
- 17 **Toro-Huamanchumo CJ**, Cabanillas-Ramirez C, Quispe-Vicuña C, Caballero-Alvarado JA, León-Figueroa DA, Cruces-Tirado N, Barboza JJ. Mean Platelet Volume in Neonatal Sepsis: Meta-Analysis of Observational Studies. *Children (Basel)* 2022; **9** [PMID: 36553264 DOI: 10.3390/children9121821]
- 18 **Tanju C**, Ekrem G, Berksoy Emel A, Nur A. Mean platelet volume as a negative marker of inflammation in children with rotavirus gastroenteritis. *Iran J Pediatr* 2014; **24**: 617-622 [PMID: 25793071]
- 19 **Wang X**, Xu XL, Li XM, Zhao R, Yang X, Cong HL. Diagnostic Value of Mean Platelet Volume Combined With Troponin I for Acute Coronary Syndrome. *Am J Med Sci* 2016; **352**: 159-165 [PMID: 27524214 DOI: 10.1016/j.amjms.2016.04.014]
- 20 **Oktay MM**, Boğan M, Çolak ST, Sabak M, Gümüşboğa H, Eren SH. Evaluation of the diagnostic value of platelet indices in pediatric acute appendicitis. *J Int Med Res* 2020; **48**: 300060520946515 [PMID: 32962484 DOI: 10.1177/0300060520946515]
- 21 **Omran A**, Ali M, Saleh MH, Zekry O. Salivary C-reactive protein and mean platelet volume in diagnosis of late-onset neonatal pneumonia. *Clin Respir J* 2018; **12**: 1644-1650 [PMID: 29028152 DOI: 10.1111/crj.12723]
- 22 **Pamudji K**, Kardana I. Diagnostic value of mean platelet volume in neonatal sepsis. *Paediatrica Indonesiana* 2019; **59**: 289-293 [DOI: 10.14238/pi59.6.2019.289-93]
- 23 **Wang J**, Wang Z, Zhang M, Lou Z, Deng J, Li Q. Diagnostic value of mean platelet volume for neonatal sepsis: A systematic review and meta-analysis. *Medicine (Baltimore)* 2020; **99**: e21649 [PMID: 32769935 DOI: 10.1097/MD.00000000000021649]
- 24 **Panda SK**, Nayak MK, Thangaraj J, Das P, Pugalia R. Platelet parameters as a diagnostic marker in early diagnosis of neonatal sepsis- Seeking newer answers for older problems. *J Family Med Prim Care* 2022; **11**: 1748-1754 [PMID: 35800552 DOI: 10.4103/jfmpc.jfmpc_1271_21]
- 25 **Barekatin B**, HasanGhalyaei N, Mohammadzadeh M, Tavakolfard N. Investigation of salivary C-reactive protein and interleukin-18 for the diagnosis of neonatal sepsis. *J Res Med Sci* 2021; **26**: 131 [PMID: 35126594 DOI: 10.4103/jrms.JRMS_1256_20]
- 26 **Kumar R**, Musoke R, Macharia WM, Revathi G. Validation of c-reactive protein in the early diagnosis of neonatal sepsis in a tertiary care hospital in Kenya. *East Afr Med J* 2010; **87**: 255-261 [PMID: 23057268 DOI: 10.4314/eamj.v87i6.63084]
- 27 **Omran A**, Maarroof A, Mohammad MHS, Abdelwahab A. Salivary C-reactive protein, mean platelet volume and neutrophil lymphocyte ratio as diagnostic markers for neonatal sepsis. *J Pediatr (Rio J)* 2018; **94**: 82-87 [PMID: 28734690 DOI: 10.1016/j.jpmed.2017.03.006]
- 28 **Xiao X**, Xue L, Sheng HL, Xiao ZH. Correlation between serum levels of C-reactive protein and infant pneumonia: A meta-analysis. *Exp Ther Med* 2015; **9**: 2331-2338 [PMID: 26136982 DOI: 10.3892/etm.2015.2417]
- 29 **Li X**, Chen Z. Correlation between serum levels of C-reactive protein and neonatal pneumonia: A protocol for systematic review and meta-analysis. *Medicine (Baltimore)* 2021; **100**: e25977 [PMID: 34011087 DOI: 10.1097/MD.00000000000025977]
- 30 **Chirico G**, Loda C. Laboratory aid to the diagnosis and therapy of infection in the neonate. *Pediatr Rep* 2011; **3**: e1 [PMID: 21647274 DOI: 10.4081/pr.2011.e1]
- 31 **Chiesa C**, Natale F, Pascone R, Osborn JF, Pacifico L, Bonci E, De Curtis M. C reactive protein and procalcitonin: reference intervals for preterm and term newborns during the early neonatal period. *Clin Chim Acta* 2011; **412**: 1053-1059 [PMID: 21338596 DOI: 10.1016/j.cca.2011.02.020]

- 32 **Laborada G**, Rego M, Jain A, Guliano M, Stavola J, Ballabh P, Krauss AN, Auld PA, Nesin M. Diagnostic value of cytokines and C-reactive protein in the first 24 hours of neonatal sepsis. *Am J Perinatol* 2003; **20**: 491-501 [PMID: 14703598 DOI: 10.1055/s-2003-45382]
- 33 **Ishibashi M**, Takemura Y, Ishida H, Watanabe K, Kawai T. C-reactive protein kinetics in newborns: application of a high-sensitivity analytic method in its determination. *Clin Chem* 2002; **48**: 1103-1106 [PMID: 12089183]
- 34 **Li CX**, Zhang L, Yan YR, Ding YJ, Lin YN, Zhou JP, Li N, Li HP, Li SQ, Sun XW, Li QY. A narrative review of exploring potential salivary biomarkers in respiratory diseases: still on its way. *J Thorac Dis* 2021; **13**: 4541-4553 [PMID: 34422380 DOI: 10.21037/jtd-21-202]
- 35 **Tsai CM**, Tang KS, Cheng MC, Liu TY, Huang YH, Chen CC, Yu HR. Use of saliva sample to detect C-reactive protein in children with pneumonia. *Pediatr Pulmonol* 2020; **55**: 2457-2462 [PMID: 32633868 DOI: 10.1002/ppul.24947]
- 36 **Iyengar A**, Paulus JK, Gerlanc DJ, Maron JL. Detection and potential utility of C-reactive protein in saliva of neonates. *Front Pediatr* 2014; **2**: 131 [PMID: 25485262 DOI: 10.3389/fped.2014.00131]
- 37 **Babaei M**, Rezaei S, Saghafi Khadem S, Shirinbak I, Basir Shabestari S. The Role of Salivary C-Reactive Protein in Systemic and Oral Disorders: A Systematic Review. *Med J Islam Repub Iran* 2022; **36**: 138 [PMID: 36479533 DOI: 10.47176/mjiri.36.138]
- 38 **Tosson AMS**, Koptan D, Abdel Aal R, Abd Elhady M. Evaluation of serum and salivary C-reactive protein for diagnosis of late-onset neonatal sepsis: A single center cross-sectional study. *J Pediatr (Rio J)* 2021; **97**: 623-628 [PMID: 33582092 DOI: 10.1016/j.jpmed.2021.01.004]
- 39 **Pay JB**, Shaw AM. Towards salivary C-reactive protein as a viable biomarker of systemic inflammation. *Clin Biochem* 2019; **68**: 1-8 [PMID: 30995442 DOI: 10.1016/j.clinbiochem.2019.04.006]
- 40 **Ragab SM**, El-Sayed HM, El-Deeb SM, El-Arabi RH. Salivary C-reactive protein, mean platelet volume, and neutrophil-lymphocyte ratio as diagnostic markers of neonatal sepsis. *Menoufia Med J* 2022; **35**: 1810-1815
- 41 **Ball S**, Dennis JA, Bedanie G, Nugent K. Relation between mean platelet volume and C-reactive protein. *Proc (Bayl Univ Med Cent)* 2020; **33**: 163-168 [PMID: 32313453 DOI: 10.1080/08998280.2019.1710658]
- 42 **Omran A**, Abohadid H, Mohammad MHS, Shalaby S. Salivary C-Reactive Protein and Mean Platelet Volume in the Diagnosis and Follow-Up of Community-Acquired Pneumonia in Infants. *Pediatr Allergy Immunol Pulmonol* 2021; **34**: 141-146 [PMID: 34860599 DOI: 10.1089/ped.2021.0077]
- 43 **Eichberger J**, Resch E, Resch B. Diagnosis of Neonatal Sepsis: The Role of Inflammatory Markers. *Front Pediatr* 2022; **10**: 840288 [PMID: 35345614 DOI: 10.3389/fped.2022.840288]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: office@baishideng.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

