

Supplementary 1 Endoscopic methods and outcomes of patients with Calcifying fibrous tumors

CFTs (n = 36)	
Growth pattern, n (%)	
Intraluminal growth	32 (88.9)
Extraluminal growth	4 (11.1)
Smooth mucosa, n (%)	36 (100)
Irregular morphology, n (%)	2 (5.6)
Max diameter (cm), mean ± SD	10.6 ± 4.3
Location, n (%)	
Esophagus	3 (8.3)
Stomach	26 (72.2)
Gastric fundus	4 (15.4)
Gastric body	20 (76.9)
Gastric antrum	2 (7.7)
Small intestine	1 (2.8)
Colon	2 (5.6)
Rectum	4 (11.1)
Layer, n (%)	
Submucosa	7 (19.4)
Muscularis propria	29 (80.6)
Accompanied CFTs in other parts of the body, n (%)	0 (0)
Surgical methods, n (%)	
EMR	1 (2.8)
ESD	14 (38.9)
EFTR	16 (44.4)
STER	5 (13.9)
Full-thickness resection	16 (44.4)
Suture method	
Unstitched	2 (5.6)
Metal clip	25 (69.4)

Metal clip and nylon rope	8 (22.2)
Metal stent	1 (2.8)
Surgery time (min), mean \pm SD	39.8 \pm 29.8
Gastrointestinal decompression	
None	20 (55.6)
Gastric tube	14 (38.9)
Anal tube	2 (5.6)
En bloc resection rate, <i>n</i> (%)	34 (94.4)
Complete resection rate, <i>n</i> (%)	34 (94.4)
Hospital stay (d), mean \pm SD	2.9 \pm 1.2
Complications, <i>n</i> (%)	0 (0)
New growth of CFTs, <i>n</i> (%)	2 (5.6)
Recurrence, <i>n</i> (%)	0 (0)

CFTs: Calcifying fibrous tumors; EMR: Endoscopic mucosal resection; EFTR: Endoscopic full-thickness resection; ESD: Endoscopic submucosal dissection; STER: Submucosal tunneling endoscopic resection.