

## Endoscopic ultrasound-guided biliary intervention in patients with surgically altered anatomy

Aroon Siripun, Pimsiri Sripongpun, Bancha Ovartharnporn

Aroon Siripun, Bancha Ovartharnporn, NKC Institute of Gastroenterology and Hepatology, Faculty of Medicine, Prince of Songkla University, Hat Yai, Songkhla 90110, Thailand  
Pimsiri Sripongpun, Department of Internal Medicine, Faculty of Medicine, Prince of Songkla University, Hat Yai, Songkhla 90110, Thailand

**Author contributions:** Siripun A and Sripongpun P performed the literature search and wrote the first draft of the paper; Ovartharnporn B reviewed and edited the article.

**Conflict-of-interest:** Siripun A, Sripongpun P and Ovartharnporn B have no conflict of interest to declare.

**Data sharing:** No additional data are available.

**Open-Access:** This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Correspondence to:** Bancha Ovartharnporn, MD, NKC Institute of Gastroenterology and Hepatology, Faculty of Medicine, Prince of Songkla University, 15 Karnjanavanich Rd., Hat Yai, Songkhla 90110, Thailand. [mrovarth@ji-net.com](mailto:mrovarth@ji-net.com)

Telephone: +66-81-7664879

Fax: +66-74-429436

Received: August 27, 2014

Peer-review started: August 31, 2014

First decision: November 27, 2014

Revised: January 8, 2015

Accepted: February 4, 2015

Article in press: February 9, 2015

Published online: March 16, 2015

2014 reporting on EUS-BD in patients with surgically altered anatomy using the terms "EUS drainage" and "altered anatomy". All relevant articles were accessed in full text. A manual search of the reference lists of relevant retrieved articles was also performed. Only full-text English papers were included. Data regarding age, gender, diagnosis, method of EUS-BD and intervention, type of altered anatomy, technical success, clinical success, and complications were extracted and collected. Anatomic alterations were categorized as: group 1, Billroth I; group 2, Billroth II; group 4, Roux-en-Y with gastric bypass; and group 3, all other types.

**RESULTS:** Twenty three articles identified in the literature search, three reports were from the same group with different numbers of cases. In total, 101 cases of EUS-BD in patients with altered anatomy were identified. Twenty-seven cases had no information and were excluded. Seventy four cases were included for analysis. Data of EUS-BD in patients categorized as group 1, 2 and 4 were limited with 2, 3 and 6 cases with EUS-BD done respectively. Thirty four cases with EUS-BD were reported in group 3. The pooled technical success, clinical success, and complication rates of all reports with available data were 89.18%, 91.07% and 17.5%, respectively. The results are similar to the reported outcomes of EUS-BD in general, however, with limited data of EUS-BD in patients with altered anatomy rendered it difficult to draw a firm conclusion.

**CONCLUSION:** EUS-BD may be an option for patients with altered anatomy after a failed endoscopic-retrograde-cholangiography in centers with expertise in EUS-BD procedures in a research setting.

**Key words:** Endoscopic ultrasound-guided antegrade approach; Endoscopic ultrasound-guided biliary drainage; Endoscopic ultrasound-guided choledochoduodenostomy; Endoscopic ultrasound-guided hepaticogastrostomy; Endoscopic ultrasound-rendezvous technique; Surgically altered anatomy; Overtube-assisted enteroscopy-endoscopic retrograde cholangiopancreatography

### Abstract

**AIM:** To evaluate the efficacy of endoscopic ultrasound guided biliary drainage (EUS-BD) in patients with surgically altered anatomies.

**METHODS:** We performed a search of the MEDLINE database for studies published between 2001 to July

**Core tip:** Endoscopic retrograde cholangiopancreatography (ERCP) in patients with surgically altered anatomy is challenging, with a failure rate as high as 26%. Data of endoscopic ultrasound-guided biliary drainage (EUS-BD) in patients with altered anatomy from the literature show a similar efficacy to that of EUS-BD in general. EUS-BD may be selected as an alternative for patients with altered anatomy who failed overtube-assisted enteroscopy-ERCP in centers where the expertise in EUS-BD is available. However, the EUS-BD approach should be performed in a research setting based on the current stage of EUS-BD techniques.

Siripun A, Sripongpun P, Ovartharnporn B. Endoscopic ultrasound-guided biliary intervention in patients with surgically altered anatomy. *World J Gastrointest Endosc* 2015; 7(3): 283-289 Available from: URL: <http://www.wjgnet.com/1948-5190/full/v7/i3/283.htm> DOI: <http://dx.doi.org/10.4253/wjge.v7.i3.283>

## INTRODUCTION

Endoscopic retrograde cholangiopancreatography (ERCP) has been widely accepted as a standard procedure with a high success rate for the management of biliary disorders<sup>[1]</sup>. However, conventional ERCP in patients with surgically altered anatomy is technically difficult, and is accompanied by a relatively high rate of complications. In large case series, technical failures varied from 13% to 67%, and the rate of perforation was as high as 18%, with a mortality rate of 3%<sup>[2,3]</sup>. ERCP with overtube-assisted enteroscopy (OAE-ERCP), with a balloon or spiral overtube, achieved a success rate of approximately 74% and a 3.4% complication rate in patients with surgically altered anatomies<sup>[4]</sup>. Percutaneous transhepatic biliary drainage (PTBD) is a well-established technique that is usually selected as an alternative in patients with failed ERCP. However, despite the high clinical success rate, the PTBD approach is associated with 0.5%-15% morbidity and 0%-4.9% mortality rates<sup>[5]</sup>.

Endoscopic ultrasound-guided biliary drainage (EUS-BD) was first reported in 2001 by Giovannini *et al*<sup>[6]</sup>. Subsequently, many groups reported the utilization of EUS-BD with various approaches as an alternative biliary drainage for failed ERCPs, with an average success rate varying from 77% to 94% and complication rate of 19%-27%<sup>[7]</sup>. This method may be an option for patients with altered anatomy for whom OAE-ERCP is difficult, as reflected by a high failure rate (26%)<sup>[4]</sup>. However, the role of EUS-BD in patients with altered anatomy and failed ERCP is not well defined, and the suitability of this as an alternative drainage procedure is unclear. This review analyzes the clinical efficacy, complications, clinical implication, and

limitations of EUS-BD in patients with surgically altered anatomy from data available in the literature.

## MATERIALS AND METHODS

A PubMed search of the MEDLINE database was conducted for articles published between 2001 and July 2014 using the terms "EUS drainage" and "altered anatomy". A manual search of the reference lists of relevant retrieved articles was also performed. Only full-text English papers were included. The computerized endoscopic data at our center were also searched for additional cases of EUS-BD in altered anatomy conducted after our published data<sup>[8]</sup>. Data regarding age, gender, diagnosis, method of EUS-BD and intervention, type of altered anatomy, technical success, clinical success, and complications were collected.

### Classification of EUS-BD techniques

The three main techniques of EUS-guided procedures for biliary drainage that were included were: antegrade EUS-BD, transluminal drainage, and the rendezvous method<sup>[7,9,10]</sup>. Antegrade EUS-BD involves intervention *via* an antegrade route across the ampulla or anastomosis. Transluminal drainage encompasses transesophageal, transgastric (hepaticogastrostomy), transduodenal (choledochoduodenostomy, EUS-CDS), or transjejunal approaches. The rendezvous method involves EUS-guided placement of a guide-wire across the ampulla or anastomosis that is exchanged with a standard duodenoscope or enteroscope to perform the intervention.

### Classification of surgically altered anatomy

Anatomic alterations were categorized as: group 1, Billroth I; group 2, Billroth II; group 3, Roux-en-Y with pancreaticoduodenectomy (with or without a modified Child procedure), pylorus-preserving pancreaticoduodenectomy, hepaticojejunostomy, choledochojejunostomy, and total or partial gastrectomy, distal gastrectomy without specific anastomosis mentioned, and hepatic and bile duct resection; and group 4, Roux-en-Y with gastric bypass (RY-GB).

### Statistical analysis

The descriptive data of age was analyzed by using Minitab 15<sup>®</sup> and no other statistical analysis was done since this study was a descriptive review.

## RESULTS

### EUS-BD IN patients with surgically altered anatomy

Of the 23 articles identified in the literature search, three reports were from the same group with different numbers of cases<sup>[11-13]</sup>. As no details regarding individual cases were available from these three reports, the cases with the same type of surgically

**Table 1** Demographics of endoscopic ultrasound-guided biliary drainage patients with altered anatomy[8,11,14,18-32] (*n* = 74)

Characteristic	Value
Age, yr ( <i>n</i> = 36)	64.14 ± 15.08
Sex, female/male ( <i>n</i> = 36)	22/14
Diagnosis ( <i>n</i> = 38)	
Bile duct stone	14
Benign stricture	10
Malignant stricture	11
Gastric cancer	2
Occluded stent	1
Outcome of EUS-BD, <i>n</i> (%)	
Clinical success	51 (91.1)
Complications	7 (17.5)
Technical success	66 (89.2)
EUS-BD technique with technical success, <i>n</i>	
Anterograde	28
Hepaticogastrostomy	18
Hepaticojejunostomy	2
Hepaticoesophagostomy	1
Rendezvous	3
Unavailable	14

EUS-BD: Endoscopic ultrasound-guided biliary drainage.

altered anatomy and the same diagnosis were treated as one case. In total, 101 cases of EUS-BD in patients with altered anatomy were identified. Only one report was a case series<sup>[14]</sup>, all other reports were case reports or reports of EUS-BD that included normal and altered anatomy patients. Twenty-seven cases had no information and were excluded<sup>[12,15-17]</sup>, leaving 74 patients with altered anatomy who underwent EUS-BD<sup>[8,11,14,18-32]</sup>. Available demographic and procedural information of these cases is presented in Table 1. Of the 40 cases reporting complications, there were incidences of mild pancreatitis (*n* = 2), mild abdominal pain (*n* = 1), hematoma (*n* = 1), cholangitis (*n* = 1), minor bleeding (*n* = 1), and surgical repositioning of a stent in the peritoneum (*n* = 1), with no mortalities.

### EUS-BD in altered anatomy subtypes

The classification of altered anatomy types is listed in Table 2. In group 1, one case of EUS-BD with common bile duct (CBD) stones was reported<sup>[25]</sup>, and EUS-BD was performed in one patient with distal CBD stricture at our center. Both had successful clinical outcomes. There were three cases with EUS-BD in group 2: two cases with malignant stricture and one case with a CBD stone, in whom the EUS-BD failed<sup>[11,32]</sup>.

Within group 3, Roux-en-Y was performed with hepaticojejunostomy (*n* = 10), choledochojejunostomy (*n* = 1), total gastrectomy (*n* = 5), subtotal gastrectomy (*n* = 2), Whipple's operation (*n* = 6), distal gastrectomy (*n* = 2), pylorus-preserving pancreaticoduodenectomy (*n* = 5), pancreaticoduodenectomy with a modified Child procedure (*n* = 2), and hepatic and bile duct resection (*n* = 1)<sup>[8,11,14,18-32]</sup>. The diagnoses in group 3 included bile duct stones (*n* = 9), benign stricture (*n* = 9), malignant stricture (*n* = 9), occluded metallic

**Table 2** Results of endoscopic ultrasound-guided biliary drainage in altered anatomy subgroups

Group	Subtype	No. of cases	Technical success	Clinical success	Complications
1	Billroth I	2	2	2	0
2	Billroth II	3	2	2	0
3	Roux-en-Y	34	33 (97.0%)	23 (92.0%)	5
4	Roux-en-Y gastric bypass	6	6	5	1

Twenty-seven cases with unspecified type and no details available were excluded.

stents placed by percutaneous route (*n* = 1), or were unspecified (*n* = 6). The success rates were in the range reported for EUS-BD in general. However, the missing data in a large proportion of patients in this group rendered it difficult to draw a firm conclusion.

Six cases within group 4 received EUS-BD with RY-GB<sup>[33]</sup>. These patients all had CBD stones, and stone clearance was achieved with EUS-BD in five of these, with a failure in one with a hematoma.

In 27 cases that were excluded from analysis due to insufficient information<sup>[12,15-17]</sup>, the overall technical success rate (including patients with altered anatomy) varied from 67.2% to 94.0%, the clinical success rate varied from 63.2% to 97.0%, and the complication rate varied from 12.0% to 23.2%.

## DISCUSSION

### EUS-BD vs PTBD

PTBD is a traditional alternative for patients with a failed ERCP, though it is associated with a risk of complication and significant morbidity<sup>[5]</sup>. One of the major drawbacks of PTBD is external bile loss, which leads to a decreased total bile pool. Theoretically, maintenance of enterohepatic bile circulation is important for host defense function. Kamiya *et al.*<sup>[33]</sup> reported that bile replacement by oral intake of the externally diverted bile helped restore gut barrier function in patients with bile duct obstruction, but internal drainage is still more physiologic than external drainage. Moreover, the burden to the patients or family members caring for the catheter is considerable, and individuals who bathe twice daily may be disturbed by the inability to do so, thus decreasing their quality of life. In one retrospective study that compared 22 EUS-BD patients with 51 PTBD patients, the procedures showed a similar clinical success, but EUS-BD was associated with fewer adverse events and was less costly in the long term<sup>[34]</sup>. However, PTBD in their study had a 100% success rate, which was significantly higher than the 86.4% with EUS-BD. Another retrospective study compared 25 cases with EUS-BD with 26 cases with PTBD, and showed that EUS-BD was superior to PTBD in terms of success and complication rates<sup>[35]</sup>. In contrast, a prospective study showed a similar efficacy between EUS-BD performed

in 13 patients and PTBD in 12 patients<sup>[36]</sup>. Taken together, these data suggest that EUS-BD is a suitable alternative in patients with failed ERCP, and it may be an option in the centers where EUS-BD is available.

### **Role of EUS-BD in altered anatomy patients**

The available data suggest that EUS-BD is as effective in patients with altered anatomy as in general patients. EUS-BD is still in a state of development, with proper procedural techniques under refinement. Furthermore, EUS-BD for patients with altered anatomy and failed ERCP should be assessed in a research setting to properly define its role. A standardized treatment algorithm for selection of EUS-BD techniques based on the clinical context may improve the outcome<sup>[26]</sup>.

In patients with a Billroth I operation, the straight anatomy of the stomach and duodenum cause the tip of a standard duodenoscope to come too close to the papilla, making it difficult to position the ERCP catheter along the axis of the bile duct, leading to a failed procedure in some patients<sup>[37]</sup>. In patients with Billroth II anatomy, OAE-ERCP has an endoscopic success rate of 96% and a successful ERCP rate of 90%<sup>[4]</sup>. ERCP is the most difficult in patients with RY-GB, and OAE-ERCP has an endoscopic success rate of only 80% and successful ERCP rate of only 70%<sup>[4]</sup>. In post RY-GB patients with failed OAE-ERCP, laparoscopy-assisted ERCP may be an alternative, as the results in four publications<sup>[38-41]</sup> demonstrated a high success rate of 90%-100%. However, these studies were limited by the number of patients, longer procedure time, the need for a laparoscopic doctor, and a much higher cost of treatment. The data supporting the role of EUS-BD in groups 1, 2 and 4 were very limited, and need further evaluation.

EUS-BD may be a suitable alternative in patients with failed OAE-ERCP with altered anatomy classified as group 3. In patients with benign stricture, the accepted treatment includes extended multiple plastic stents or metallic stent placement<sup>[42-44]</sup>. Short-term outcome of EUS-BD for a small number of these patients was promising, though no long-term data is available<sup>[8,11,14,18,21,27,31]</sup>. Anterograde balloon dilation has been reported in very few cases, with a successful short-term outcome<sup>[8,14,21]</sup>, and transgastric placement of multiple plastic stents across the anastomotic stricture was feasible in select patients. One patient in our report had a good long-term result after three years<sup>[8]</sup>. Because of the repeated nature of the procedures in this group for the assessment of stricture patency or insertion of an additional stent, EUS-BD with anterograde or rendezvous techniques may be initially selected as a bridging procedure in the patients with endoscopic access of the papilla or biliary anastomotic site, but failed ERCP cannulation. At present, EUS-BD in patients with benign stricture and failed OAE-ERCP access to the papilla or biliary anastomosis is challenging. In patients with malignant strictures, the same approach is applicable, but

transluminal drainage is preferred, as repeated procedures may be easier using a standard endoscope. In patients with bile duct stones associated with altered anatomy and OAE-ERCP access to the papilla or biliary anastomosis with failed ERCP cannulation, EUS-BD with anterograde or rendezvous procedures may be preferred for stone removal<sup>[4,45]</sup>. EUS-BD with anterograde stone removal using balloon dilation with the stones pushed across the ampulla or anastomosis was reported in 11 patients with one failure<sup>[8,14,22]</sup>, and may be an option in select patients with failed OAE-ERCP access to the papilla or biliary anastomosis. Placement of a transgastric nasobiliary drainage tube or a plastic stent to maintain access for subsequent repeated procedures was also an option<sup>[8,14]</sup>. The details of the procedure should be customized based on clinical setting.

### **EUS-BD as an initial modality**

OAE-ERCP is increasingly used in patients with altered anatomy with more supporting data compared with EUS-BD<sup>[4,8,11,14,18-32,45]</sup>, though no comparative studies are available. In patients with benign strictures, standard ERCP (for patients with Billroth I or II anatomy, Whipple's operation,) or OAE-ERCP is more suitable because of the likelihood for repeated procedures for additional stent placement or stent exchange. In patients with malignant strictures, EUS-BD may be an alternative to PTBD in centers with appropriate expertise when OAE-ERCP is not available, however, this should be done in a research setting. For patients with bile duct stones, OAE-ERCP may be suitable as the options for treatment of the stones are more readily available, and EUS-BD should be reserved for patients in whom this procedure fails.

### **Limitations of EUS-BD**

As it is difficult to pass the linear EUS endoscope into the afferent limb<sup>[46]</sup>, EUS-CDS is not the appropriate option for patients with altered anatomies. The EUS-BD drainage access is limited to the left biliary system, and requires the presence of a dilated ductal system. Manipulation of the guide-wire to cross a stricture or papilla may be difficult, and the guide-wire can be sheared<sup>[47,48]</sup>. As only limited data for anterograde EUS-BD were available<sup>[48]</sup>, the success rate may be lower with a lower complication rate compared with other techniques<sup>[7]</sup>. EUS-hepaticogastrostomy is limited by the lack of adherence between the stomach and the liver, which may increase the risk of stent dislocation and lead to bile leak. The risk of bleeding from the liver may also increase<sup>[48]</sup>. The main limitation of the rendezvous method is the requirement of an endoscopically accessible papilla or anastomosis, which is always troublesome in cases of surgically altered anatomy. In addition, the rendezvous procedure requires exchanging the echoendoscope for a duodenoscope, during which guide-wire access can be lost<sup>[48]</sup>.

The majority of the data concerning EUS-BD is reported by experts, and may not translate to clinical practice. For example, in a national study in Spain involving community endoscopists, EUS-BD had a lower success rate (67.2%) and a complication rate of 23.2%<sup>[15]</sup>. Moreover, there is no well-designed EUS-training system and training using swine models, or computer-based simulators are expensive and not accessible by all trainees<sup>[47]</sup>. This may hinder the establishment of skills in therapeutic EUS.

## FUTURE DEVELOPMENTS

Most of EUS-BDs were performed with conventional fine-needle aspiration needles. The new 19-gauge blunt tip (Echo-HD; Cook Medical, Bloomington, IN, United States) may reduce catching at the needle tip during to-and-fro manipulation of a guide-wire that may reduce shearing. Needle-knife dilation was reported to increase the risk of complications in EUS-BD<sup>[49]</sup>. The tip of the needle knife may not align with the axis of the guide-wire, thus a 6 Fr catheter with diathermic ring (Endoflex, Voerde, Germany) was used in some centers<sup>[47]</sup>. A prototype compression coil and twin-headed needle may simplify the EUS-BD procedure, and shows promise for use in EUS-CDS in a study in canines<sup>[50]</sup>. The development of a forward-viewing echoendoscope allows simultaneous visualization of the endoscopic and EUS operating fields, while the perpendicular access and lack of angulation at the exit of the working channel allow for easy introduction of a 19-gauge needle and passing of the stent without indentation<sup>[51]</sup>. Although the forward-viewing echoendoscope showed a high success rate for EUS-CDS in a prospective case series<sup>[52]</sup>, further studies are needed to confirm its advantage in EUS-BD. Multiple exchanges over the wire during EUS-BD may increase the risk of leakage, increase the procedure time, and increase the chance to lose guide-wire access<sup>[47]</sup>. Non-exchange systems have been evaluated in experimental animal studies<sup>[53,54]</sup>, and may minimize the aforementioned drawbacks of the current EUS-BD technique when the devices are available in the future.

## CONCLUSION

EUS-guided biliary intervention is technically feasible and the available data indicate a high success rate in patients with surgically altered anatomies. Although the complication rate may be higher than for OAE-ERCP in patients with altered anatomy (17.5% vs 3.4%<sup>[4]</sup>), EUS-BD may be a rescue option in patients for whom OAE-ERCP has failed when conducted within centers with appropriate expertise and in a research setting. A standardized algorithm for using different EUS-BD techniques, refinement of these methods, and the development of new devices may improve the efficacy of EUS-BD and minimize the complication

rate. The role of forward-viewing echoendoscope and comparison with the current standard EUS endoscope remain to be assessed.

## COMMENTS

### Background

Surgically altered anatomy is a consequence of an operation for treatment of a specific disease. This precludes a normal access to the bile duct opening by a standard duodenoscope in many cases and an over-tube assisted endoscopy (OAE) is usually needed to access the bile duct. However, OAE has a failure rate as high as 26%. Endoscopic ultrasound guided biliary drainage (EUS-BD), a recently developed technique, showed a high success rate. The efficacy of EUS-BD in altered anatomy is not well defined.

### Research frontiers

To the best of our knowledge, no review of EUS-BD in surgically altered anatomy has been previously published. The objective of this study was to review systematically the efficacy of EUS-BD in altered anatomy.

### Innovations and breakthroughs

EUS-BD in the setting of surgically altered anatomy has an efficacy similar to EUS-BD in general, nonetheless, the available data were limited and further studies to evaluate the role of EUS-BD in altered anatomy are needed.

### Applications

EUS-BD may be an option for patient with surgically altered anatomy with a failed OAE therapeutic intervention but this should be done in centers with the expertise in EUS-BD in a research setting.

### Terminology

EUS-BD is a technique using an endoscope with ultrasound technology to visualize a bile duct. A needle puncture of the bile duct was done under ultrasound guidance then intervention was done using various kinds of endoscopic accessories. OAE is a technique of endoscopy that utilizing an over-tube with a balloon at the tip or an over-tube with a spiral configuration to facilitate the insertion of an enteroscope.

### Peer-review

This study was well investigated and will give us important information especially in clinical gastroenterology.

## REFERENCES

- Freeman ML, Guda NM. ERCP cannulation: a review of reported techniques. *Gastrointest Endosc* 2005; **61**: 112-125 [PMID: 15672074 DOI: 10.1016/S0016-5107(04)02463-0]
- Hintze RE, Veltke W, Adler A, Abou-Rebyeh H. Endoscopic sphincterotomy using an S-shaped sphincterotome in patients with a Billroth II or Roux-en-Y gastrojejunostomy. *Endoscopy* 1997; **29**: 74-78 [PMID: 9101142 DOI: 10.1055/s-2007-1004078]
- Wright BE, Cass OW, Freeman ML. ERCP in patients with long-limb Roux-en-Y gastrojejunostomy and intact papilla. *Gastrointest Endosc* 2002; **56**: 225-232 [PMID: 12145601]
- Skinner M, Popa D, Neumann H, Wilcox CM, Mönkemüller K. ERCP with the overtube-assisted enteroscopy technique: a systematic review. *Endoscopy* 2014; **46**: 560-572 [PMID: 24839188 DOI: 10.1055/s-0034-1365698]
- Kühn JP, Busemann A, Lerch MM, Heidecke CD, Hosten N, Puls R. Percutaneous biliary drainage in patients with nondilated intrahepatic bile ducts compared with patients with dilated intrahepatic bile ducts. *AJR Am J Roentgenol* 2010; **195**: 851-857 [PMID: 20858809 DOI: 10.2214/AJR.09.3461]
- Giovannini M, Moutardier V, Pesenti C, Bories E, Lelong B, Delperio JR. Endoscopic ultrasound-guided bilioduodenal anastomosis: a new technique for biliary drainage. *Endoscopy* 2001; **33**: 898-900 [PMID: 11571690 DOI: 10.1055/s-2001-17324]
- Iwashita T, Doi S, Yasuda I. Endoscopic ultrasound-guided biliary drainage: a review. *Clin J Gastroenterol* 2014; **7**: 94-102 [PMID: 24765215 DOI: 10.1007/s12328-014-0467-5]
- Attasaryanya S, Netinasunton N, Jongboonyanuparp T, Sottisuporn J, Witeerungrot T, Pirathvisuth T, Ovartlarnporn B. The Spectrum of

- Endoscopic Ultrasound Intervention in Biliary Diseases: A Single Center's Experience in 31 Cases. *Gastroenterol Res Pract* 2012; **2012**: 680753 [PMID: 22654900 DOI: 10.1155/2012/680753]
- 9 **Sarkaria S**, Lee HS, Gaidhane M, Kahaleh M. Advances in endoscopic ultrasound-guided biliary drainage: a comprehensive review. *Gut Liver* 2013; **7**: 129-136 [PMID: 23560147 DOI: 10.5009/gnl.2013.7.2.129]
- 10 **Kedia P**, Gaidhane M, Kahaleh M. Endoscopic guided biliary drainage: how can we achieve efficient biliary drainage? *Clin Endosc* 2013; **46**: 543-551 [PMID: 24143319 DOI: 10.5946/ce.2013.46.5.543]
- 11 **Kahaleh M**, Hernandez AJ, Tokar J, Adams RB, Shami VM, Yeaton P. Interventional EUS-guided cholangiography: evaluation of a technique in evolution. *Gastrointest Endosc* 2006; **64**: 52-59 [PMID: 16813803 DOI: 10.1016/j.gie.2006.01.063]
- 12 **Maranki J**, Hernandez AJ, Arslan B, Jaffan AA, Angle JF, Shami VM, Kahaleh M. Interventional endoscopic ultrasound-guided cholangiography: long-term experience of an emerging alternative to percutaneous transhepatic cholangiography. *Endoscopy* 2009; **41**: 532-538 [PMID: 19533558 DOI: 10.1055/s-0029-1214712]
- 13 **Kahaleh M**, Wang P, Shami VM, Tokar J, Yeaton P. EUS-guided transhepatic cholangiography: report of 6 cases. *Gastrointest Endosc* 2005; **61**: 307-313 [PMID: 15729253 DOI: 10.1016/S0016-5107(04)02585-4]
- 14 **Iwashita T**, Yasuda I, Doi S, Uemura S, Mabuchi M, Okuno M, Mukai T, Itoi T, Moriwaki H. Endoscopic ultrasound-guided antegrade treatments for biliary disorders in patients with surgically altered anatomy. *Dig Dis Sci* 2013; **58**: 2417-2422 [PMID: 23535877 DOI: 10.1007/s10620-013-2645-6]
- 15 **Vila JJ**, Pérez-Miranda M, Vazquez-Sequeiros E, Abadia MA, Pérez-Millán A, González-Huix F, Gornals J, Iglesias-García J, De la Serna C, Aparicio JR, Subtil JC, Alvarez A, de la Morena F, García-Cano J, Casi MA, Lancho A, Barturen A, Rodríguez-Gómez SJ, Repiso A, Juzgado D, Igea F, Fernandez-Urien I, González-Martin JA, Armengol-Miró JR. Initial experience with EUS-guided cholangiopancreatography for biliary and pancreatic duct drainage: a Spanish national survey. *Gastrointest Endosc* 2012; **76**: 1133-1141 [PMID: 23021167 DOI: 10.1016/j.gie.2012.08.001]
- 16 **Iwashita T**, Lee JG, Shinoura S, Nakai Y, Park DH, Muthusamy VR, Chang KJ. Endoscopic ultrasound-guided rendezvous for biliary access after failed cannulation. *Endoscopy* 2012; **44**: 60-65 [PMID: 22127960 DOI: 10.1055/s-0030-1256871]
- 17 **Khashab MA**, Valeshabad AK, Modayil R, Widmer J, Saxena P, Idrees M, Iqbal S, Kalloo AN, Stavropoulos SN. EUS-guided biliary drainage by using a standardized approach for malignant biliary obstruction: rendezvous versus direct transluminal techniques (with videos). *Gastrointest Endosc* 2013; **78**: 734-741 [PMID: 23886353 DOI: 10.1016/j.gie.2013.05.013]
- 18 **Artifon EL**, Safate-Ribeiro AV, Ferreira FC, Poli-de-Figueiredo L, Rasslan S, Carnevale F, Otoch JP, Sakai P, Kahaleh M. EUS-guided antegrade transhepatic placement of a self-expandable metal stent in hepatico-jejunal anastomosis. *JOP* 2011; **12**: 610-613 [PMID: 22072253]
- 19 **Ogura T**, Masuda D, Imoto A, Takeushi T, Kamiyama R, Mohamed M, Umegaki E, Higuchi K. EUS-guided hepaticogastrostomy combined with fine-gauge antegrade stenting: a pilot study. *Endoscopy* 2014; **46**: 416-421 [PMID: 24573771 DOI: 10.1055/s-0034-1365020]
- 20 **Nguyen-Tang T**, Binmoeller KF, Sanchez-Yague A, Shah JN. Endoscopic ultrasound (EUS)-guided transhepatic antegrade self-expandable metal stent (SEMS) placement across malignant biliary obstruction. *Endoscopy* 2010; **42**: 232-236 [PMID: 20119894 DOI: 10.1055/s-0029-1243858]
- 21 **Park do H**, Jang JW, Lee SS, Seo DW, Lee SK, Kim MH. EUS-guided transhepatic antegrade balloon dilation for benign bilioenteric anastomotic strictures in a patient with hepaticojejunostomy. *Gastrointest Endosc* 2012; **75**: 692-693 [PMID: 21679943 DOI: 10.1016/j.gie.2011.04.013]
- 22 **Weilert F**, Binmoeller KF, Marson F, Bhat Y, Shah JN. Endoscopic ultrasound-guided antegrade treatment of biliary stones following gastric bypass. *Endoscopy* 2011; **43**: 1105-1108 [PMID: 22057823 DOI: 10.1055/s-0030-1256961]
- 23 **Prachayakul V**, Aswakul P. A novel technique for endoscopic ultrasound-guided biliary drainage. *World J Gastroenterol* 2013; **19**: 4758-4763 [PMID: 23922474 DOI: 10.3748/wjg.v19.i29.4758]
- 24 **Shah JN**, Marson F, Weilert F, Bhat YM, Nguyen-Tang T, Shaw RE, Binmoeller KF. Single-operator, single-session EUS-guided antegrade cholangiopancreatography in failed ERCP or inaccessible papilla. *Gastrointest Endosc* 2012; **75**: 56-64 [PMID: 22018554 DOI: 10.1016/j.gie.2011.08.032]
- 25 **Püspök A**, Lomoschitz F, Dejaco C, Hejna M, Sautner T, Gangl A. Endoscopic ultrasound guided therapy of benign and malignant biliary obstruction: a case series. *Am J Gastroenterol* 2005; **100**: 1743-1747 [PMID: 16086710 DOI: 10.1111/j.1572-0241.2005.41806.x]
- 26 **Park do H**, Jeong SU, Lee BU, Lee SS, Seo DW, Lee SK, Kim MH. Prospective evaluation of a treatment algorithm with enhanced guidewire manipulation protocol for EUS-guided biliary drainage after failed ERCP (with video). *Gastrointest Endosc* 2013; **78**: 91-101 [PMID: 23523301 DOI: 10.1016/j.gie.2013.01.042]
- 27 **Bories E**, Pesenti C, Caillol F, Lopes C, Giovannini M. Transgastric endoscopic ultrasonography-guided biliary drainage: results of a pilot study. *Endoscopy* 2007; **39**: 287-291 [PMID: 17357952 DOI: 10.1055/s-2007-966212]
- 28 **Ma J**, Liu Y, Li Z, Jin Z. Endoscopic ultrasound-guided transgastric biliary drainage after partial gastrectomy. *Endoscopy* 2011; **43** Suppl 2 UCTN: E102 [PMID: 21424995 DOI: 10.1055/s-0030-1256104]
- 29 **Kawakubo K**, Isayama H, Kato H, Itoi T, Kawakami H, Hanada K, Ishiwatari H, Yasuda I, Kawamoto H, Itokawa F, Kuwatani M, Iiboshi T, Hayashi T, Doi S, Nakai Y. Multicenter retrospective study of endoscopic ultrasound-guided biliary drainage for malignant biliary obstruction in Japan. *J Hepatobiliary Pancreat Sci* 2014; **21**: 328-334 [PMID: 24026963 DOI: 10.1002/jhbp.27]
- 30 **Horaguchi J**, Fujita N, Noda Y, Kobayashi G, Ito K, Obana T, Takasawa O, Koshita S, Kanno Y. Endosonography-guided biliary drainage in cases with difficult transpapillary endoscopic biliary drainage. *Dig Endosc* 2009; **21**: 239-244 [PMID: 19961522 DOI: 10.1111/j.1443-1661.2009.00899.x]
- 31 **Will U**, Thieme A, Fuedner F, Gerlach R, Wanzar I, Meyer F. Treatment of biliary obstruction in selected patients by endoscopic ultrasonography (EUS)-guided transluminal biliary drainage. *Endoscopy* 2007; **39**: 292-295 [PMID: 17357950 DOI: 10.1055/s-2007-966215]
- 32 **Burmester E**, Niehaus J, Leineweber T, Huetteroth T. EUS-cholangio-drainage of the bile duct: report of 4 cases. *Gastrointest Endosc* 2003; **57**: 246-251 [PMID: 12556796 DOI: 10.1067/mge.2003.85]
- 33 **Kamiya S**, Nagino M, Kanazawa H, Komatsu S, Mayumi T, Takagi K, Asahara T, Nomoto K, Tanaka R, Nimura Y. The value of bile replacement during external biliary drainage: an analysis of intestinal permeability, integrity, and microflora. *Ann Surg* 2004; **239**: 510-517 [PMID: 15024312 DOI: 10.1097/01.sla.0000118594.23874.89]
- 34 **Khashab MA**, Valeshabad AK, Afghani E, Singh VK, Kumbhari V, Messallam A, Saxena P, El Zein M, Lennon AM, Canto MI, Kalloo AN. A comparative evaluation of EUS-guided biliary drainage and percutaneous drainage in patients with distal malignant biliary obstruction and failed ERCP. *Dig Dis Sci* 2015; **60**: 557-565 [PMID: 25081224 DOI: 10.1007/s10620-014-3300-6]
- 35 **Bapaye A**, Dubale N, Aher A. Comparison of endosonography-guided vs. percutaneous biliary stenting when papilla is inaccessible for ERCP. *United European Gastroenterol J* 2013; **1**: 285-293 [PMID: 24917973 DOI: 10.1177/2050640613490928]
- 36 **Artifon EL**, Aparicio D, Paione JB, Lo SK, Bordini A, Rabello C, Otoch JP, Gupta K. Biliary drainage in patients with unresectable, malignant obstruction where ERCP fails: endoscopic ultrasonography-guided choledochoduodenostomy versus percutaneous drainage. *J Clin Gastroenterol* 2012; **46**: 768-774 [PMID: 22810111 DOI: 10.1097/MCG.0b013e31825f264c]

- 37 **Imazu H**, Kanazawa K, Ikeda K, Kakutani H, Sumiyama K, Ang TL, Omar S, Tajiri H. Initial evaluation of a novel multibending backward-oblique viewing duodenoscope in endoscopic retrograde cholangiopancreatography. *Endoscopy* 2012; **44**: 99-102 [PMID: 22068702 DOI: 10.1055/s-0031-1291445]
- 38 **Schreiner MA**, Chang L, Gluck M, Irani S, Gan SI, Brandabur JJ, Thirlby R, Moonka R, Kozarek RA, Ross AS. Laparoscopy-assisted versus balloon enteroscopy-assisted ERCP in bariatric post-Roux-en-Y gastric bypass patients. *Gastrointest Endosc* 2012; **75**: 748-756 [PMID: 22301340 DOI: 10.1016/j.gie.2011.11.019]
- 39 **Lopes TL**, Clements RH, Wilcox CM. Laparoscopy-assisted ERCP: experience of a high-volume bariatric surgery center (with video). *Gastrointest Endosc* 2009; **70**: 1254-1259 [PMID: 19846085 DOI: 10.1016/j.gie.2009.07.035]
- 40 **Desai SV**, Naveed M, Jazwinski A, Jowell PS, Branch MS. Spiral enteroscopy versus laparoscopic-assisted endoscopy for completion of ERCP in patients with roux-en-y gastric bypass surgery [abstract]. *Gastrointest Endosc* 2011; **73**: AB122 [DOI: 10.1016/j.gie.2011.03.039]
- 41 **Bertin PM**, Singh K, Arregui ME. Laparoscopic transgastric endoscopic retrograde cholangiopancreatography (ERCP) after gastric bypass: case series and a description of technique. *Surg Endosc* 2011; **25**: 2592-2596 [PMID: 21416184 DOI: 10.1007/s00464-011-1593-5]
- 42 **Devière J**, Nageshwar Reddy D, Püspök A, Ponchon T, Bruno MJ, Bourke MJ, Neuhaus H, Roy A, González-Huix Lladó F, Barkun AN, Kortan PP, Navarrete C, Peetermans J, Blero D, Lakhtakia S, Dolak W, Lepilliez V, Poley JW, Tringali A, Costamagna G. Successful management of benign biliary strictures with fully covered self-expanding metal stents. *Gastroenterology* 2014; **147**: 385-395; quiz e15 [PMID: 24801350 DOI: 10.1053/j.gastro.2014.04.043]
- 43 **Baron TH**, Davee T. Endoscopic management of benign bile duct strictures. *Gastrointest Endosc Clin N Am* 2013; **23**: 295-311 [PMID: 23540962 DOI: 10.1016/j.giec.2013.01.001]
- 44 **Costamagna G**, Boškoski I. Current treatment of benign biliary strictures. *Ann Gastroenterol* 2013; **26**: 37-40 [PMID: 24714594]
- 45 **Shah RJ**, Smolkin M, Yen R, Ross A, Kozarek RA, Howell DA, Bakis G, Jonnalagadda SS, Al-Lehibi AA, Hardy A, Morgan DR, Sethi A, Stevens PD, Akerman PA, Thakkar SJ, Brauer BC. A multicenter, U.S. experience of single-balloon, double-balloon, and rotational overtube-assisted enteroscopy ERCP in patients with surgically altered pancreaticobiliary anatomy (with video). *Gastrointest Endosc* 2013; **77**: 593-600 [PMID: 23290720 DOI: 10.1016/j.gie.2012.10.015]
- 46 **Wilson JA**, Hoffman B, Hawes RH, Romagnuolo J. EUS in patients with surgically altered upper GI anatomy. *Gastrointest Endosc* 2010; **72**: 947-953 [PMID: 21034896 DOI: 10.1016/j.gie.2010.07.016]
- 47 **Kahaleh M**, Artifon EL, Perez-Miranda M, Gupta K, Itoi T, Binmoeller KF, Giovannini M. Endoscopic ultrasonography guided biliary drainage: summary of consortium meeting, May 7th, 2011, Chicago. *World J Gastroenterol* 2013; **19**: 1372-1379 [PMID: 23538784 DOI: 10.3748/wjg.v19.i9.1372]
- 48 **Binmoeller KF**, Nguyen-Tang T. Endoscopic ultrasound-guided antegrade cholangiopancreatography. *J Hepatobiliary Pancreat Sci* 2011; **18**: 319-331 [PMID: 21190119 DOI: 10.1007/s00534-010-0358-1]
- 49 **Park do H**, Jang JW, Lee SS, Seo DW, Lee SK, Kim MH. EUS-guided biliary drainage with transluminal stenting after failed ERCP: predictors of adverse events and long-term results. *Gastrointest Endosc* 2011; **74**: 1276-1284 [PMID: 21963067 DOI: 10.1016/j.gie.2011.07.054]
- 50 **Chang KJ**, Chao HH. EUS-Guided choledochoduodenostomy (ECD) for immediate and long-term treatment of biliary obstruction using prototype compression coil and twin-head needle. *Gastrointest Endosc* 2011; **73**: AB326
- 51 **Fusaroli P**, Ceroni L, Caletti G. Forward-view Endoscopic Ultrasound: A Systematic Review of Diagnostic and Therapeutic Applications. *Endosc Ultrasound* 2013; **2**: 64-70 [PMID: 24949367 DOI: 10.4103/2303-9027.117689]
- 52 **Hara K**, Yamao K, Hijioka S, Mizuno N, Imaoka H, Tajika M, Kondo S, Tanaka T, Haba S, Takeshi O, Nagashio Y, Obayashi T, Shinagawa A, Bhatia V, Shimizu Y, Goto H, Niwa Y. Prospective clinical study of endoscopic ultrasound-guided choledochoduodenostomy with direct metallic stent placement using a forward-viewing echoendoscope. *Endoscopy* 2013; **45**: 392-396 [PMID: 23338620 DOI: 10.1055/s-0032-1326076]
- 53 **Binmoeller KF**, De La Mora-Levy JG. An Exchange-Free Device for Advanced Translumenal Therapy. *Gastrointest Endosc* 2010; **71**: AB349
- 54 **Lee TH**, Choi JH, Lee SS, Cho HD, Seo DW, Park SH, Lee SK, Kim MH, Park do H. A pilot proof-of-concept study of a modified device for one-step endoscopic ultrasound-guided biliary drainage in a new experimental biliary dilatation animal model. *World J Gastroenterol* 2014; **20**: 5859-5866 [PMID: 24914346 DOI: 10.3748/wjg.v20.i19.5859]

**P- Reviewer:** Fabbri C, Hak NG, Naito Y, Rerknimitr R  
**S- Editor:** Tian YL **L- Editor:** A **E- Editor:** Zhang DN





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>

