

Advanced non-small cell lung cancer in elderly patients: A review

Lucio Buffoni, Lorena Consito, Andrea Riccardo Filippi, Enrico Ruffini, Paolo Solidoro, Paolo Bironzo, Maria Antonietta Satolli, Marina Schena, Libero Ciuffreda

Lucio Buffoni, Lorena Consito, Paolo Bironzo, Marina Schena, Libero Ciuffreda, Department of Medical Oncology, San Giovanni Battista Molinette Hospital, 10126 Turin, Italy

Andrea Riccardo Filippi, Department of Oncology, Radiation Oncology, University of Turin, 10124 Turin, Italy

Enrico Ruffini, Department of Thoracic Surgery, School of Medicine, University of Turin, 10124 Turin, Italy

Paolo Solidoro, Division of Pulmonology, University of Turin, 10124 Turin, Italy

Maria Antonietta Satolli, Department of Oncology, University of Turin, 10124 Turin, Italy

Author contributions: Being a review article authors contributed to conception and design of the review, acquisition of data, and analysis and interpretation of data; drafting the article and final approval of the version of the article to be published.

Conflict-of-interest statement: The authors disclosed any conflict of interest with the material included in the study.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Correspondence to: Lucio Buffoni, MD, Department of Medical Oncology, San Giovanni Battista Molinette Hospital, 15, Via Cherasco, 10126 Turin, Italy. buffonil75@gmail.com
Telephone: +39-11-6335189
Fax: +39-11-6335189

Received: September 24, 2014
Peer-review started: September 26, 2014
First decision: December 17, 2014
Revised: March 3, 2015

Accepted: April 1, 2015

Article in press: April 7, 2015

Published online: July 28, 2015

Abstract

Over 50% of patients diagnosed with non-small-cell lung cancer (NSCLC) are 65 years old while 30% exceed 70 years old. Comparing elderly patients to their younger counterpart they poorly tolerate chemotherapy due to progressive reduction of organ function and age-related co-existing pathologies. Due to this reason elderly are usually excluded from platinum-based chemotherapy, which still represent the standard of care for advanced NSCLC. In every-day practice, single-agent schedule with a third-generation drug is the recommended option for elderly patients with advanced NSCLC. A modest increase in toxicity for elderly patients has been demonstrated by subgroup analyses concluding for platinum-based combination chemotherapy being similar in young patients and fit elderly. Even though the cited evidence, feasibility of chemotherapy based on platinum remains an open question. Prospective randomised trials are warranted in order to change guide lines and give the clinicians a new therapeutic option. Recent emerging role of molecular target in selecting patients for new targeted therapies suggest dedicated trials for elderly patients. The same is for more accurate evaluation of elderly patients with increasing evidence for a comprehensive geriatric assessment as a valid tool for customized treatment in NSCLC elderly patients. Suitable evidences for the treatment of elderly patients affected by advanced NSCLC together with more appropriate and validated tools for patients selection are reviewed along the manuscript.

Key words: Lung cancer; Elderly; Chemotherapy; Target therapy

© The Author(s) 2015. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Due to progressive ageing of population in the next few years a consistent proportion of non-small-cell lung cancer (NSCLC) patients will be diagnosed over the age of 70 years old. Guide lines indications together with results from most recent phase III trials dedicated to elderly patients are discussed along the review. Special attention has been deserved to toxicity profile. Recent emerging role of molecular target in selecting patients for new targeted therapies suggest dedicated trials as for more accurate evaluation with increasing evidence for a comprehensive geriatric assessment as a valid tool for treatment selection in NSCLC elderly patients.

Buffoni L, Consito L, Filippi AR, Ruffini E, Solidoro P, Bironzo P, Satolli MA, Schena M, Ciuffreda L. Advanced non-small cell lung cancer in elderly patients: A review. *World J Respirol* 2015; 5(2): 102-111 Available from: URL: <http://www.wjgnet.com/2218-6255/full/v5/i2/102.htm> DOI: <http://dx.doi.org/10.5320/wjr.v5.i2.102>

INTRODUCTION

Non-small-cell lung cancer (NSCLC) still represents the most frequent cause of cancer-related deaths in Europe and many western countries. Median age at diagnosis of NSCLC is around 70 years old with 50% and more diagnosed over 65 and 30% over 70^[1,2]. Continuous shifting toward an older population will cause oncologists seeing more elderly patients with lung cancer in years to come.

The highest percentage of patients presents with metastatic disease at diagnosis with chemotherapy standing for the gold standard of management. It is demonstrated that many older patients with metastatic NSCLC are not treated due to many reasons^[3,4]. Aging is cause of physiological changes in organ functions, as for example renal and liver function. Co-morbidities and consequent drugs intake which may condition chemotherapy administration and toxicity profile^[5]. An immediate consequence is a sort of under-treatment for this setting of patients^[6].

It is known that these patients are not candidated to clinical trials causing difficulty in reaching evidence-based clinical recommendation. In current clinical practice the choice of treatment is still very often based on a old-believe that cancer in older people behave in a less aggressive manner^[7].

A significant difference has been documented on a survey which considered trials for cancer drug registration from 1995 to 2002 with only 35% of elderly patients enrolled on clinical trials^[8]. Such a discordance can greatly affect the reproducibility of trial results.

Chronological age should be no more considered the correct parameter on which candidate patients to receive specific treatment. Biological age should be defined through laboratory tests and geriatric assessment evaluation. However chronological age still represent a reference for clinical trials: 70 years old is considered a reasonable cut-off considering the incidence of age-related variations starts to increase after the age of 70^[9]. Many evidences suggest an advanced median age of elderly patients in dedicated trials with milder reported toxicity respect to that reported in age-unspecified studies^[10].

SINGLE-AGENT CHEMOTHERAPY

The ELVIS trial was the first phase III trial conducted in advanced NSCLC patients aged ≥ 70 years old. Patients were randomized to receive vinorelbine at a dose of 30 mg/m² on days 1 and 8 or best supportive care. Advantage in term of survival and increased quality of life (QoL) were demonstrated for vinorelbine arm^[11].

Regarding the role of gemcitabine in the treatment of elderly patients affected by advanced NSCLC many specifically addressed phase II trials confirmed its interesting role with an overall response rates (ORR) of 18% to 38% and MST of 6.8 to 9 mo. Toxicity was mild with only sporadic grade 3 to 4 hematologic toxicities in two of them^[12-15].

Other specific trials demonstrated satisfying data regarding both activity and feasibility for paclitaxel and docetaxel: ORR 3%-23% and MST 6.8-10.3 mo^[16]. Overall response rate (ORR) and median survival time (MST) were as follow: 3%-23% and 6.8-10.3 mo, respectively, with acceptable toxicity^[16-21].

When docetaxel was compared to vinorelbine in a randomized phase III trial no significant difference in MST was encountered, while longer progression-free survival (PFS) and higher ORR were demonstrated for docetaxel^[22].

NON-PLATINUM-BASED POLICHEMOTHERAPY

Among non-platinum-based regimen the association of gemcitabine plus vinorelbine has been widely studied. At least two phase III trials compared the combination with single agent schedule. Frasci *et al*^[23] obtained survival advantage for combination respect to single agent in term of MST and ORR.

The Multicenter Italian Lung cancer in the Elderly Study (MILES) trial with its 700 elderly patients affected by advanced NSCLC, represents a milestone in this setting. Patients were randomised to receive vinorelbine or gemcitabine or combination. Considering ORR, time to progression (TTP), MST or QoL no advantage was demonstrated over single-agent therapy in favour of combination therapy. Single-agent therapy confirmed

Table 1 Results from phase III trials of advanced non-small-cell lung cancer in elderly patients: Non-platinum based chemotherapy

Ref.	Regimen	Age (yr)	No. of patients	RR (%)	MST (mo)
Gridelli ^[11]	Vinorelbine	70	76	20	6.5
	<i>vs</i>				
Frasci <i>et al</i> ^[23]	Best Supportive Care		78	NA	4.8
	Vinorelbine	70	60	15	4.2
Gridelli <i>et al</i> ^[24]	<i>vs</i>				
	Vinorelbine + Gemcitabine		60	22	6.7
Kudoh <i>et al</i> ^[22]	Vinorelbine or	70	233	18	8.3
	Gemcitabine		233	16	6.5
Kudoh <i>et al</i> ^[22]	<i>vs</i>				
	Vinorelbine + Gemcitabine		232	21	6.9
Quoix <i>et al</i> ^[54]	Vinorelbine	≥ 70	91	9.9	9.9
	<i>vs</i>				
Quoix <i>et al</i> ^[54]	Docetaxel		88	22.7	14.3
	Vinorelbine or Gemcitabine	≥ 70	226		10.3
Hainsworth <i>et al</i> ^[25]	<i>vs</i>				
	Carboplatin/paclitaxel		225		6.2
Hainsworth <i>et al</i> ^[25]	Docetaxel	≥ 65	171	17	5.1
	<i>vs</i>				
	Docetaxel/gemcitabine		174	25	5.5

RR: Response rate; MST: Median survival time; NA: Not applicable.

its role in this setting^[24].

Hainsworth *et al*^[25] demonstrated that docetaxel plus gemcitabine *vs* weekly docetaxel alone did not improve survival in a population of 350 accrued and randomized elderly patients affected by advanced NSCLC. Moreover, in these two trials the doublet was slightly more toxic than single agent treatment^[25] (Table 1).

PLATINUM-BASED CHEMOTHERAPY

Cisplatin represents standard treatment for advanced NSCLC. Nephrotoxicity, ototoxicity and neurotoxicity are the most common non-haematological toxicities attributed to cisplatin in addition to haematological ones.

Carboplatin is responsible for lower incidence of nausea, nephrotoxicity and neurotoxicity, although safety remains an issue also due to its administration in combination with other myelotoxic agents.

Retrospective analyses of platinum-based chemotherapy

In the last decade many data were collected from large randomised trials not selected for elderly patients.

No significant differences in terms of efficacy were shown in retrospective analysis from ECOG 5592 comparing effects of two different platinum-based schedules in patients older than 70 years respect to younger counterpart. In terms of toxicity elderly patients had worse leukopenia and neuropsychiatric disorders^[26]. ECOG 1594 trial compared four treatment combinations in first-line with no significant differences for RR and MST in retrospective subset analysis for the 227 patients (20%) aged ≥ 70. Significant ($P = 0.04$) major grade 4 toxicities were reported in the elderly subgroup^[27]. Always referring to retrospective analysis the Southwest Oncology Group (SWOG) 9509 trial,

and the SWOG 9308 trial, documented no significant age influence on MST, TTP and toxicity^[28].

In more recent years, the CALGB compared carboplatin plus paclitaxel with paclitaxel. MST was similar between patients aged > 70 and their younger counterpart. A secondary analysis evidenced a survival advantage for the doublet in the elderly patients^[29].

TAX 326 compared first-line cisplatin plus vinorelbine or docetaxel. In the subset analysis considering patients aged ≥ 65 increase in survival was obtained with docetaxel plus cisplatin respect to vinorelbine/cisplatin. Lowest incidence in toxicity was registered for docetaxel plus carboplatin arm^[30].

Comparing weekly paclitaxel or standard dose paclitaxel in combination with carboplatin in the elderly, the fractionated regimen produced higher RR, TTP and MST. Significant less neuropathy was encountered in the experimental arm^[31-34] (Table 2).

Platinum-based chemotherapy: Prospectives studies

Many prospective phase II trials evaluating third-generation cytotoxic agents with modified platinum-based schedules were performed in the last 20 years. Cisplatin and gemcitabine combination was tested in four phase II trials reaching an ORR around 40% and a MST of 10 mo^[35-38]. At least three phase II trials tested cisplatin/vinorelbine schedule with similar ORR and MST^[39-41]. Both combinations demonstrated to be safe. Better results in terms of ORR and MST were demonstrated by Ohe *et al*^[42] in patients aged ≥ 75 years by adding docetaxel to weekly cisplatin^[42,43].

In 2007 Gridelli *et al*^[44] tested in the MILES 2P the feasibility of cisplatin with gemcitabine or vinorelbine in elderly patients. Both combinations were feasible and active with the former combination being the preferred one for a direct comparison with standard single-agent

Table 2 Retrospective data analyses of elderly patients enrolled in phase III trials with cisplatin- or carboplatin-based chemotherapy

Ref.	Treatment	Age (yr)	No. of patients	RR	MST (mo)	P value
Nguyen <i>et al</i> ^[27]	CDDP + GEM	≥ 70	53	15%	7.7	NS
		< 70	207	29%	9.4	
Kelly <i>et al</i> ^[28]	CBDCA + TAX	≥ 70	117	NR	6.9	0.06
		< 70	491	NR	8.6	
Langer <i>et al</i> ^[26]	CDDP + VP-16	≥ 70	86	23.3%	8.5	NS
		< 70	488	21.5%	9.1	
Rocha Lima <i>et al</i> ^[32]	CDDP + VBL	≥ 70	31	16%	5.7	NS
		< 70	222	31%	8.0	
Hensing <i>et al</i> ^[33]	CBDCA + TAX	≥ 70	67	27%	7.1	NS
		< 70	163	20%	7.8	
Belani <i>et al</i> ^[30]	CDDP + TXT	≥ 65	149	NR	12.6	NS
		All ages	408	32%	11.3	
	CDDP + VNR	≥ 65	134	NR	9.9	NS
		All ages	404	25%	10.1	
Belani <i>et al</i> ^[30]	CBDCA + TAXw	≥ 65	118	NR	9.0	NS
		All ages	406	24%	9.4	
	CBDCA + TAX	≥ 70	70	25.7%	9.2	NR
		< 70	147	28.6%	9.6	
Lilenbaum <i>et al</i> ^[29]	CBDCA + TAX	≥ 70	63	19%	7.7	NR
		< 70	151	19.2%	11.4	
		≥ 70	77	36%	8.0	
< 70	207	30%	8.5			

CDDP: Cisplatin; CBDCA: Carboplatin; TAX: Paclitaxel; TXT: Docetaxel; VNR: Vinorelbine; GEM: Gemcitabine; VBL: Vinblastine; VP-16: Etoposide; RR: Response rate; MST: Median survival time; S: Survival; NR: Not reported; NS: Not significant.

Table 3 Phase II trials of cisplatin-based chemotherapy with third-generation agents and modified schedules or attenuated doses of cisplatin

Ref.	Regimen	CDDP dose	Age (yr)	No. of patients	RR	MST (mo)
¹ Mattioli <i>et al</i> ^[39]	CDDP + VNR	25 mg/m ² , weekly	> 65	36	36%	11
Pereira <i>et al</i> ^[40]	CDDP + VNR	60-90 mg/m ²	> 70	44	50%	7.5
Buffoni <i>et al</i> ^[41]	CDDP + VNR	30 mg/m ² , day 1 and 8	≥ 70	30	33%	7.4
Lippe <i>et al</i> ^[35]	CDDP + GEM	35 mg/m ² , weekly	≥ 65	29	48%	10
Berardi <i>et al</i> ^[36]	CDDP + GEM	35 mg/m ² , weekly	≥ 70	48	31.8%	9
Feliu <i>et al</i> ^[37]	CDDP + GEM	50 mg/m ²	≥ 70	46	35%	10.2
Moscetti <i>et al</i> ^[38]	CDDP + GEM	75 mg/m ² , day 2	≥ 65	46	45.6%	15
Ohe <i>et al</i> ^[42]	CDDP + TXT	25 mg/m ² , weekly	≥ 75	33	52%	15.8

¹Including 3 unfit patients; CDDP: Cisplatin; VNR: Vinorelbine; GEM: Gemcitabine; TXT: Docetaxel; RR: Response rate; MST: Median survival time; S: Survival.

chemotherapy in this setting (Table 3).

Carboplatin plus vinorelbine combination was tested in two phase II studies without any clinical benefit compared to standard treatment^[45,46]. More favourable results were reported for the combination of low-dose carboplatin (AUC 4) and gemcitabine accompanied by acceptable toxicity^[47].

Modified carboplatin/paclitaxel schedules reached in many trials 70% RR and 14 mo MST^[10,48-53].

Recently IFCT-0501 confronted attenuated carboplatin-paclitaxel schedule vs standard single agent monotherapy. Combination and standard treatment showed respectively 10.3 and 6.2 mo in terms of OS. Regarding toxicity, grade 3-4 neutropenia and thrombocytopenia were 54.3% and 6.3% for doublet respect to 14.3% and 1% for single agent. Considering non-hematological toxicity, neuropathy resulted significantly more frequent in the doublet arm

vs single agent arm. Considering QoL, role functioning and fatigue were worse in the doublet group than in single agent. Authors concluded that despite increased toxic effects for doublet, this should be considered as standard treatment in first line setting^[54].

For the future better CDDP-based schedule in terms of activity and tolerability should be tested by direct comparison and considering emerging strong data about histology (Table 4).

TARGETED THERAPIES

Bevacizumab

The SAIL study assessed the addition of bevacizumab to standard chemotherapy in terms of safety and efficacy in the first-line. When evaluating incidence of anti-VEGFR related side-effects in a planned analysis including elderly patients no difference was

Table 4 Prospective trials of first line platinum-based chemotherapy in elderly patients affected by advanced non-small-cell lung cancer

Ref.	Regimen	Phase	Age (yr)	No. of patients	Efficacy
Gridelli <i>et al</i> ^[44]	CDDP + GEM	I / II	≥ 70	159	OS 43.6 wk PFS 25.3 wk RR 43.5%
	CDDP + VNR				OS 33.1 wk PFS 21.1 wk RR 36.1%
Abe <i>et al</i> ^[43]	CDDP + DOC <i>vs</i> DOC	III	> 70	221	OS 13.3 wk OS 17.3 wk
Biesma <i>et al</i> ^[53]	CARBO + GEM	III	≥ 70	181	OS 8.6 mo RR 27%
	CARBO + PAC				OS 6.9 mo RR 19%
Quoix <i>et al</i> ^[54]	GEM or VNR	III	≥ 70	451	OS 6.2 mo RR 10.9%
	CARBO + PAC				OS 10.3 mo RR 29.5%

CDDP: Cisplatin; VNR: Vinorelbine; GEM: Gemcitabine; CARBO: Carboplatin; RR: Response rate; OS: Overall survival.

encountered. OS (14.6 mo in both groups), TTP (8.2 mo *vs* 7.6 mo), RR (49.3% *vs* 52.4%) and disease control rate (89.3% *vs* 88.4%) were similar in both arms^[55].

The ARIES trial evaluated bevacizumab in clinical practice. Six hundred and fifty enrolled patients were older than 70 and experienced similar adverse events than total population except for arterial thromboembolic events (slightly increased in patients ≥ 70 years old). Median PFS and OS were similar in both subgroups^[56].

EGFR tyrosine kinase inhibitors

Gefitinib/erlotinib: Gefitinib and erlotinib are reversible inhibitor of EGFR that competitively inhibits the binding of ATP.

Antitumor activity of single-agent gefitinib in patients unselected for EGFR status has been tested in many trials. When gefitinib was dispensed in the second line setting in an unselected populations with NSCLC obtained 5.3 mo of MST^[57].

In the subgroup of elderly patients gefitinib maintained its activity and good tolerance with no grade 3 or 4 side effects experienced^[58]. Cavina *et al*^[59] reported encouraging efficacy data with only 10% of grade 3 skin toxicity and 3% of diarrhoea. Gridelli *et al*^[60] reported in the same setting of patients similar MST and favourable safety profile. Cappuzzo *et al*^[61] observed reported 5 mo MST and mild side effects. Hotta *et al*^[62] studied gefitinib on patients aged ≥ 75 years: RR 17%; SD 43%; MST 7.6 mo. Grade 3-4 toxicity was encountered in 9% of patients (Table 5).

Gefitinib addition to chemotherapy has been tested. Stinchcombe *et al*^[63] combined weekly docetaxel plus daily oral gefitinib: RR 31%, MST 6.5

mo, 1-year survival 27%. The schedule resulted in excessive toxicity for elderly patients^[63]. Bepler *et al*^[64] added daily gefitinib to three-weekly docetaxel: RR 38%, SD 24%, MST 12.4 mo, and 1-year survival of 60%. Better tolerance was reported even if adverse effects required hospitalization in 6 patients^[64]. A phase II trial evaluated gefitinib with vinorelbine or gemcitabine. Vinorelbine plus gefitinib produced 72% of grade 3-4 neutropenia and 3 treatment-related deaths while the association with gemcitabine reported a lower activity (RR 5.7%, SD 14%, MST 9.1 mo) but a better safety profile (grade 3-4 neutropenia 11.4%; thrombocytopenia 8.6%, asthenia and diarrhoea 5.7%)^[65,66] (Table 6).

At least three single-arm trials tested the role of gefitinib in patients with EGFR mutation positive tumors. A phase II study conducted with erlotinib in patients older than 70 years in I line setting, showing encouraging activity (RR of 10.9%, SD of 54.5%) and MST (10.5 mo). Adverse events were mild. EGFR mutations have been detected in 3 out of 5 responsive patients to erlotinib treatment^[67]. Erlotinib improved also QoL and many disease related symptoms^[68] (Table 7).

The EURTAC trial population with its median age of 65 years old represent an older population respect to common trial population. This trial showed that erlotinib yielded a longer progression-free survival than chemotherapy^[69].

An age-unspecified trial in patients not selected for mutation status, the BR21, showed that in a second or third line setting, erlotinib improves survival but at a cost to older patients. Although older and younger patients achieved comparable PFS, OS and RRs, older patients suffered worse toxicity due to rash, fatigue and dehydration^[70].

Erlotinib performed better than vinorelbine in a subgroup of EGFR mutation positive elderly patients as shown in a prospective phase II trial while failed to gain advantage when added to gemcitabine or compared to it in molecularly not selected elderly patients^[71,72].

SECOND-LINE CHEMOTHERAPY

Retrospective subgroup analysis on elderly patients from phase III trials testing pemetrexed *vs* docetaxel obtained: TTP 4.6 mo *vs* 2.9 mo, MST 9.5 mo *vs* 7.7 mo, 12-mo survival was 20.4% *vs* 23.1%, 24-mo survival 6.1% *vs* 10.6%, respectively. Neutropenia, febrile neutropenia and anemia were more consistent in docetaxel arm^[73]. Second-line cytotoxic therapy appeared feasible for good performance status elderly patients.

Pemetrexed produced a more favourable toxicity profile compared to docetaxel. In a phase II trial a modified schedule of docetaxel (37.5 mg/m² on days 1 and 8 every three weeks) reported encouraging activity and acceptable toxicity profile^[74].

Table 5 Retrospective analyses of gefitinib in the treatment of unselected elderly patients with advanced non-small-cell lung cancer

Ref.	Previous chemotherapy	Age (yr)	No. of patients	RR (%)	SD (%)	MST (mo)	Toxicity G \geq 3
Copin <i>et al</i> ^[58]	Yes (61%) No (39%)	\geq 70	61	2 (3)	16 (26)	NR	None
Cavina <i>et al</i> ^[59]	Yes (64.5%) No (35.5%)	\geq 70	31	0 (0)	18 (58)	3.0	G3 skin 10% G3 diarrhoea 3%
Gridelli <i>et al</i> ^[60]	Yes (94.5%) No (5.5%)	\geq 70	18	0 (0)	2 (11)	4.4	None
Cappuzzo <i>et al</i> ^[61]	Yes (100%)	\geq 70	40	2 (5)	18 (45)	5.0	G4 diarrhoea 2.5%
Hotta <i>et al</i> ^[62]	Yes (57%) No (43%)	\geq 75	92	16 (17)	40 (43)	7.6	G3-4 toxicity 9%

RR: Response rate; SD: Stable disease; MST: Median survival time; G: Grade; NR: Not reported.

Table 6 Studies with gefitinib (250 mg/d) plus chemotherapy in the treatment of elderly patients (age \geq 70 years) with advanced non-small-cell lung cancer

Ref.	Treatment	Study phase	No. of patients	RR (%)	SD (%)	MST (mo)	Toxicity G \geq 3
Stinchcombe <i>et al</i> ^[63]	TXT 30-36 mg/m ² , days 1, 8, 15, Q4W	I / II	26	8 (31)	-	6.5	G3-5 toxicity 42%
Beppler <i>et al</i> ^[64]	TXT 75 mg/m ² , day 1, Q3W	II	21	8 (38)	5 (24)	12.4	G3-4 toxicity 28.5%
Scagliotti <i>et al</i> ^[65]	GEM 1200 mg/m ² , days 1, 8, Q3W <i>vs</i> VNR 30 mg/m ² , days 1, 8, Q3W	II Random	35 25	2 (5.7)	14 (40)	9.1 12.2	G3-4 neutropenia 11.4% G3-4 neutropenia 72% 3 toxic deaths

RR: Response rate; SD: Stable disease; MST: Median survival time; G: Grade; TXT: Docetaxel; GEM: Gemcitabine; VNR: Vinorelbine; Q4W: Every 4 wk; Q3W: Every 3 wk; Random: Randomised.

Table 7 Phase II study of single-agent erlotinib in the treatment of advanced non-small-cell lung cancer elderly patients

Ref.	Previous chemotherapy	Age (yr)	No. of patients	RR (%)	SD (%)	MST (mo)	Toxicity G \geq 3
Jackman <i>et al</i> ^[67]	No (100%)	\geq 70	58	6 (10.9)	30 (54.5)	10.5	G \geq 3 toxicity 30%

RR: Response rate; SD: Stable disease; MST: Median survival time; G: Grade.

Second-line cytotoxic therapy appeared feasible for good performance status elderly patients.

A recent retrospective review of elderly patients receiving a second line therapy analyzing 293 young patients (age < 70) and 168 patients (> 70) treated with second-line treatment (both chemotherapy and EGFR TKIs) showed no differences in both efficacy and toxicity between the two age group^[75].

GERIATRIC ASSESSMENT

As elderly patients represent a very heterogeneous group, their functional status cannot be established only basing on chronological parameters. Comprehensive Geriatric Assessment (CGA) is a multidisciplinary and multidimensional approach including many items: co-existing diseases, socioeconomic status, nutritional status, medicine intake, and the presence of geriatric syndromes.

Many works demonstrated to add very useful informations regarding functional assessment of elderly

patients permitting better evaluation in terms of prognostic aspects^[76].

A meta-analysis based on 28 trials demonstrated that CGA, when applied together with geriatric interventions is capable to reduce early re-hospitalization and mortality. Nowadays no phase III randomized trials are available.

More recent studies attributed to many CGA issues the power to predict the risk of chemotherapy toxicity^[77].

Puts *et al*^[78] reported data on CGA in cancer patients collected from 73 trials. Six reported a significant association with chemotherapy toxicity, 8 demonstrated association with mortality and other 2 a change of treatment indications after CGA assessment. CGA is recommended by the International Society of Geriatric Oncology (SIOG) and the EORTC^[78].

CONCLUSION

NSCLC remains the major cause of cancer-related

deaths.

Altered organ functions and higher incidence of co-morbidities usually affect treatment decisions. Platinum-based treatment while is a mainstay for advanced NSCLC in younger patients is still considered questionable for elderly patients due to higher risk of toxicity. However many retrospective subset analyses demonstrated only minimal or none differences between elderly and younger counterpart patients.

Dedicated trials for elderly population are anyway needed. Research must be empowered looking for new tools and items capable of better define "biological" and "chronological" aspects of ageing. At the same time several treatment options should still be evaluated as for example: doublets with and without platinum, maintenance strategies and biologic agents^[73].

A third-generation agent given alone is at the time the recommended option for elderly advanced NSCLC patients^[79-81].

The choice for the best agent should consider together expected toxicities, pharmacokinetics aspects, liver-kidney function and co-existing illnesses.

Doublets employing platinum could be consider a valid indication for fit elderly patients with normal organ function. Each patient should receive a functional assessment at baseline in order to better define the therapeutic options.

Considering continuous elucidation of mechanisms that contribute to the malignant phenotype and subsequent molecular targets for anticancer therapy several biologic agents have been introduced in the treatment of NSCLC and many are still under investigation.

Trials performed with new targeted agents in molecularly selected younger population evidenced very good toxicity profile. So new biologic drugs are better candidates than chemotherapy to be tested in elderly patients. Gefitinib and erlotinib have already proven to be effective in chemotherapy-refractory NSCLC patients. Their mild toxicity profile, experienced in elderly patients with advanced disease, makes them some of the best candidates to test prospectively as single-agent first-line treatment in molecularly selected elderly population, as an alternative to chemotherapy.

REFERENCES

- 1 **Howlader N**, Noone AM, Krapcho M, Garshell J, Miller D, Altekruse SF, Kosary CL, Yu M, Ruhl J, Tatalovich Z, Mariotto A, Lewis DR, Chen HS, Feuer EJ, Cronin KA, editors. SEER Cancer Statistics Review, 1975-2011, National Cancer Institute. Bethesda, MD, [based on November 2013 SEER data submission, posted to the SEER web site, April 2014]. Available from: URL: <http://seer.cancer.gov/csr/1975-2011/>
- 2 **Jemal A**, Murray T, Ward E, Samuels A, Tiwari RC, Ghafoor A, Feuer EJ, Thun MJ. Cancer statistics, 2005. *CA Cancer J Clin* 2006; **55**: 10-30 [PMID: 15661684]
- 3 **Davidoff AJ**, Tang M, Seal B, Edelman MJ. Chemotherapy and survival benefit in elderly patients with advanced non-small-cell lung cancer. *J Clin Oncol* 2010; **28**: 2191-2197 [PMID: 20351329 DOI: 10.1200/JCO.2009.25.4052]

- 4 **Lang K**, Marciniak MD, Faries D, Stokes M, Buesching D, Earle C, Treat J, Morissette N, Thompson D. Trends and predictors of first-line chemotherapy use among elderly patients with advanced non-small cell lung cancer in the United States. *Lung Cancer* 2009; **63**: 264-270 [PMID: 18649969 DOI: 10.1016/j.lungcan.2008.05.003]
- 5 **Yancik R**, Ganz PA, Varricchio CG, Conley B. Perspectives on comorbidity and cancer in older patients: approaches to expand the knowledge base. *J Clin Oncol* 2001; **19**: 1147-1151 [PMID: 11181680]
- 6 **Fentiman IS**, Tirelli U, Monfardini S, Schneider M, Festen J, Cognetti F, Aapro MS. Cancer in the elderly: why so badly treated? *Lancet* 1990; **335**: 1020-1022 [PMID: 1970072]
- 7 **Rossi A**, Maione P, Gridelli C. Cetuximab in advanced non-small cell lung cancer. *Crit Rev Oncol Hematol* 2006; **59**: 139-149 [PMID: 16806963]
- 8 **Talarico L**, Chen G, Pazdur R. Enrollment of elderly patients in clinical trials for cancer drug registration: a 7-year experience by the US Food and Drug Administration. *J Clin Oncol* 2004; **22**: 4626-4631 [PMID: 15542812]
- 9 **Balducci L**. Geriatric oncology: challenges for the new century. *Eur J Cancer* 2000; **36**: 1741-1754 [PMID: 10974621]
- 10 **Jatoi A**, Stella PJ, Hillman S, Mailliard JA, Vanone S, Perez EA, Cannon MW, Geyer S, Wiesenfeld M, Jett JR. Weekly carboplatin and paclitaxel in elderly non-small-cell lung cancer patients (& gt; or=65 years of age): a phase II North Central Cancer Treatment Group study. *Am J Clin Oncol* 2003; **26**: 441-447 [PMID: 14528068]
- 11 **Gridelli C**. The ELVIS trial: a phase III study of single-agent vinorelbine as first-line treatment in elderly patients with advanced non-small cell lung cancer. *Elderly Lung Cancer Vinorelbine Italian Study. Oncologist* 2001; **6** Suppl 1: 4-7 [PMID: 11181997]
- 12 **Ricci S**, Antonuzzo A, Galli L, Tibaldi C, Bertuccelli M, Lopes Pegna A, Petruzzelli S, Algeri R, Bonifazi V, Fioretto ML, Orlandini C, Conte PF. Gemcitabine monotherapy in elderly patients with advanced non-small cell lung cancer: a multicenter phase II study. *Lung Cancer* 2000; **27**: 75-80 [PMID: 10688490]
- 13 **Altavilla G**, Adamo V, Buemi B, Marabello G, Maisano R, Lupo G, Bene A, Bellocco G. Gemcitabine as single agent in the treatment of elderly patients with advanced non small cell lung cancer. *Anticancer Res* 2000; **20**: 3675-3678 [PMID: 11268438]
- 14 **Martoni A**, Di Fabio F, Guaraldi M, Piana E, Ramini R, Lelli G, Palomba G, Artioli F, Bandieri E, Robustelli Della Cuna G, Preti P. Prospective phase II study of single-agent gemcitabine in untreated elderly patients with stage IIIB/IV non-small-cell lung cancer. *Am J Clin Oncol* 2001; **24**: 614-617 [PMID: 11801766]
- 15 **Gridelli C**, Cigolari S, Gallo C, Manzione L, Ianniello GP, Frontini L, Ferrau F, Robbiati SF, Adamo V, Gasparini G, Novello S, Perrone F. Activity and toxicity of gemcitabine and gemcitabine + vinorelbine in advanced non-small-cell lung cancer elderly patients: Phase II data from the Multicenter Italian Lung Cancer in the Elderly Study (MILES) randomized trial. *Lung Cancer* 2001; **31**: 277-284 [PMID: 11165408]
- 16 **Nakamura Y**, Sekine I, Furuse K, Saijo N. Retrospective comparison of toxicity and efficacy in phase II trials of 3-h infusions of paclitaxel for patients 70 years of age or older and patients under 70 years of age. *Cancer Chemother Pharmacol* 2000; **46**: 114-118 [PMID: 10972480]
- 17 **Fidias P**, Supko JG, Martins R, Boral A, Carey R, Grossbard M, Shapiro G, Ostler P, Lucca J, Johnson BE, Skarin A, Lynch TJ. A phase II study of weekly paclitaxel in elderly patients with advanced non-small cell lung cancer. *Clin Cancer Res* 2001; **7**: 3942-3949 [PMID: 11751486]
- 18 **Garbo L**, Marsland T, Garfield D, Khan M, Asmar L. A phase II study of weekly paclitaxel (Taxol) in stage IIIB, IV, or relapsed after local therapy, non-small cell lung cancer (NSCLC) patients with a performance status of 2 and/or ≥ 70 years (yrs) of age, with (Paraplatin) administered at disease progression. *Proc Am Soc Clin Oncol* 2001; **20**: 267b (abstr 2821)
- 19 **West WH**, Birch R, Sysel IA, Schell FM, Liebmann J, Tongol J, Begas A. A phase II trial of weekly paclitaxel in elderly patients or those with decreased performance status with advanced non-small

- cell lung cancer. *Proc Am Soc Clin Oncol* 2001; **20**: 258b (abstr 2782)
- 20 **Lilenbaum R**, Rubin M, Samuel J, Boros L, Chidiac T, Seigel L, Graham P. A phase II randomized trial of docetaxel weekly or every 3 weeks in elderly and/or poor performance status patients with advanced non-small cell lung cancer. *J Clin Oncol* 2004; **22** (14S): 630s (abstr 7057)
- 21 **Hainsworth JD**, Burris HA, Billings FT, Bradof JE, Baker M, Greco FA. Weekly docetaxel with either gemcitabine or vinorelbine as second-line treatment in patients with advanced nonsmall cell lung carcinoma: Phase II trials of the Minnie Pearl Cancer Research Network. *Cancer* 2001; **92**: 2391-2398 [PMID: 11745295]
- 22 **Kudoh S**, Takeda K, Nakagawa K, Takada M, Katakami N, Matsui K, Shinkai T, Sawa T, Goto I, Semba H, Seto T, Ando M, Satoh T, Yoshimura N, Negoro S, Fukuoka M. Phase III study of docetaxel compared with vinorelbine in elderly patients with advanced non-small-cell lung cancer: results of the West Japan Thoracic Oncology Group Trial (WJTOG 9904). *J Clin Oncol* 2006; **24**: 3657-3663 [PMID: 16877734]
- 23 **Frasci G**, Lorusso V, Panza N, Comella P, Nicoletta G, Bianco A, De Cataldis G, Iannelli A, Bilancia D, Belli M, Massidda B, Piantedosi F, Comella G, De Lena M. Gemcitabine plus vinorelbine versus vinorelbine alone in elderly patients with advanced non-small-cell lung cancer. *J Clin Oncol* 2000; **18**: 2529-2536 [PMID: 10893283]
- 24 **Gridelli C**, Perrone F, Gallo C, Cigolari S, Rossi A, Piantedosi F, Barbera S, Ferrau F, Piazza E, Rosetti F, Clerici M, Bertetto O, Robbiati SF, Frontini L, Sacco C, Castiglione F, Favaretto A, Novello S, Migliorino MR, Gasparini G, Galetta D, Iaffaioli RV, Gebbia V. Chemotherapy for elderly patients with advanced non-small-cell lung cancer: the Multicenter Italian Lung Cancer in the Elderly Study (MILES) phase III randomized trial. *J Natl Cancer Inst* 2003; **95**: 362-372 [PMID: 12618501]
- 25 **Hainsworth JD**, Spigel DR, Farley C, Shipley DL, Bearden JD, Gandhi J, Ann Houston G, Anthony Greco F. Weekly docetaxel versus docetaxel/gemcitabine in the treatment of elderly or poor performance status patients with advanced nonsmall cell lung cancer: a randomized phase 3 trial of the Minnie Pearl Cancer Research Network. *Cancer* 2007; **110**: 2027-2034 [PMID: 17823908]
- 26 **Langer CJ**, Manola J, Bernardo P, Kugler JW, Bonomi P, Cella D, Johnson DH. Cisplatin-based therapy for elderly patients with advanced non-small-cell lung cancer: implications of Eastern Cooperative Oncology Group 5592, a randomized trial. *J Natl Cancer Inst* 2002; **94**: 173-181 [PMID: 11830607]
- 27 **Nguyen B**, Sandler A, Denham C. The safety and efficacy of gemcitabine plus cisplatin in the elderly chemo-naïve NSCLC patients (age \geq 70 years) as compared to those with age < 70 years. *Proc Am Soc Clin Oncol* 1999; **18**: 471a (abstr 1818)
- 28 **Kelly K**, Giarratta S, Hayes S, Akerley W, Hesketh P, Wozniak A, Albain K, Crowley J. Should older patient (pts) receive combination chemotherapy for advanced stage non-small cell lung cancer (NSCLC)? An analysis of Southwest Oncology Trials 9509 and 9308. *Proc Am Soc Clin Oncol* 2001; **20**: 329a (abstr 1313)
- 29 **Lilenbaum RC**, Herndon JE, List MA, Desch C, Watson DM, Miller AA, Graziano SL, Perry MC, Saville W, Chahinian P, Weeks JC, Holland JC, Green MR. Single-agent versus combination chemotherapy in advanced non-small-cell lung cancer: the cancer and leukemia group B (study 9730). *J Clin Oncol* 2005; **23**: 190-196 [PMID: 15625373]
- 30 **Belani CP**, Fossella F. Elderly subgroup analysis of a randomized phase III study of docetaxel plus platinum combinations versus vinorelbine plus cisplatin for first-line treatment of advanced nonsmall cell lung carcinoma (TAX 326). *Cancer* 2005; **104**: 2766-2774 [PMID: 16288485]
- 31 **Belani CP**, Wang W, Johnson DH, Wagner H, Schiller J, Veeder M, Mehta M. Phase III study of the Eastern Cooperative Oncology Group (ECOG 2597): induction chemotherapy followed by either standard thoracic radiotherapy or hyperfractionated accelerated radiotherapy for patients with unresectable stage IIIA and B non-small-cell lung cancer. *J Clin Oncol* 2005; **23**: 3760-3767 [PMID: 15837967]
- 32 **Rocha Lima CM**, Herndon JE, Kosty M, Clamon G, Green MR. Therapy choices among older patients with lung carcinoma: an evaluation of two trials of the Cancer and Leukemia Group B. *Cancer* 2002; **94**: 181-187 [PMID: 11815975]
- 33 **Hensing TA**, Peterman AH, Schell MJ, Lee JH, Socinski MA. The impact of age on toxicity, response rate, quality of life, and survival in patients with advanced, Stage IIIB or IV nonsmall cell lung carcinoma treated with carboplatin and paclitaxel. *Cancer* 2003; **98**: 779-788 [PMID: 12910523]
- 34 **Perrone F**, Gallo C, Gridelli C. Re: Cisplatin-based therapy for elderly patients with advanced non-small-cell lung cancer: implications of Eastern Cooperative Oncology Group 5592, a randomized trial. *J Natl Cancer Inst* 2002; **94**: 1029-1030; author reply 1030-1031 [PMID: 12096088]
- 35 **Lippe P**, Silva RR, Giuliodori L, Monterubbianesi MC, Mattioli R, Massacesi C, Cascinu S, Tummarello D, Cellerino R. Clinical benefit of gemcitabine-cisplatin in advanced non-small cell lung cancer elderly patients. *Anticancer Res* 2002; **22**: 1053-1059 [PMID: 12168900]
- 36 **Berardi R**, Porfiri E, Scartozzi M, Lippe P, Silva RR, Nacciarriti D, Menichetti ET, Tummarello D, Carle F, Piga A, Cellerino R. Elderly patients with advanced non-small cell lung cancer. A phase II study with weekly cisplatin and gemcitabine. *Oncology* 2003; **65**: 198-203 [PMID: 14657592]
- 37 **Feliu J**, Martín G, Madroñal C, Rodríguez-Jarzáiz A, Castro J, Rodríguez A, Checa T, Bolaño M, Casado E, González-Barón M. Combination of low-dose cisplatin and gemcitabine for treatment of elderly patients with advanced non-small-cell lung cancer. *Cancer Chemother Pharmacol* 2003; **52**: 247-252 [PMID: 12783203]
- 38 **Moscetti L**, Nelli F, Padalino D, Sperduti I, Giannarelli D, Pollera CF. Gemcitabine and cisplatin in the treatment of elderly patients with advanced non-small cell lung cancer: impact of comorbidities on safety and efficacy outcome. *J Chemother* 2005; **17**: 685-692 [PMID: 16433202]
- 39 **Mattioli R**, Morese R, Imperatori L, Laici G, Gattafoni P, Lippe P. Weekly cisplatin (P) and vinorelbine (V) is active, manageable and provides clinical benefit in advanced non-small cell lung cancer (ANSCLC) elderly or poor performance status (PS) patients. *Ann Oncol* 2002; **13** (suppl 5): 104 (abstr 377P)
- 40 **Pereira JR**, Martins SJ, Nikaedo SM, Ikari FK. Chemotherapy with cisplatin and vinorelbine for elderly patients with locally advanced or metastatic non-small-cell lung cancer (NSCLC). *BMC Cancer* 2004; **4**: 69 [PMID: 15456520]
- 41 **Buffoni L**, Dongiovanni D, Barone C, Fissore C, Ottaviani D, Dongiovanni V, Grillo R, Salvadori A, Birocco N, Schena M, Bertetto O. Fractionated dose of cisplatin (CDDP) and vinorelbine (VNB) chemotherapy for elderly patients with advanced non-small cell lung cancer: phase II trial. *Lung Cancer* 2006; **54**: 353-357 [PMID: 17049672]
- 42 **Ohe Y**, Niho S, Kakinuma R, Kubota K, Ohmatsu H, Goto K, Nokihara H, Kunitoh H, Saijo N, Aono H, Watanabe K, Tango M, Yokoyama A, Nishiwaki Y. A phase II study of cisplatin and docetaxel administered as three consecutive weekly infusions for advanced non-small-cell lung cancer in elderly patients. *Ann Oncol* 2004; **15**: 45-50 [PMID: 14679118]
- 43 **Abe T**, Takeda K, Ohe Y, Kudoh S, Ichinose Y, Okamoto H, Yamamoto N, Yoshioka H, Minato K, Sawa T, Iwamoto Y, Saka H, Mizusawa J, Shibata T, Nakamura S, Ando M, Yokoyama A, Nakagawa K, Saijo N, Tamura T. Randomized phase III trial comparing weekly docetaxel plus cisplatin versus docetaxel monotherapy every 3 weeks in elderly patients with advanced non-small-cell lung cancer: the intergroup trial JCOG0803/WJOG4307L. *J Clin Oncol* 2015; **33**: 575-581 [PMID: 25584004]
- 44 **Gridelli C**, Maione P, Illiano A, Piantedosi FV, Favaretto A, Bearz A, Robbiati SF, Filipazzi V, Lorusso V, Carrozza F, Iaffaioli RV, Manzione L, Gallo C, Morabito A, Perrone F. Cisplatin plus gemcitabine or vinorelbine for elderly patients with advanced non-small-cell lung cancer: the MILES-2P studies. *J Clin Oncol* 2007; **25**: 4663-4669 [PMID: 17925563]
- 45 **Colleoni M**, Vicario G, Pancheri F, Sgarbossa G, Nelli P, Manente

- P. Weekly carboplatin and vinorelbine in elderly patients with non-small-cell lung cancer (NSCLC). Proc 6th International Congress on Anti-Cancer Treatment. Paris, France: NSCLC, 2004: 194 (abstract 562)
- 46 **Santomaggio CR**, Tucci E. Chemoterapia con carboplatino (CBDCA) e vinorelbina (V) nell'anziano affetto da carcinoma polmonare non microcitoma, stadi avanzati (A-NSCLC). *J Chemother* 1996; **8** (suppl 3): 104
 - 47 **Maestu I**, Gómez-Aldaraví L, Torregrosa MD, Camps C, Llorca C, Bosch C, Gómez J, Giner V, Oltra A, Albert A. Gemcitabine and low dose carboplatin in the treatment of elderly patients with advanced non-small cell lung cancer. *Lung Cancer* 2003; **42**: 345-354 [PMID: 14644523]
 - 48 **Choi IS**, Kim BS, Park SR, Lee SY, Kim DY, Kim JH, Lee SH, Kim TY, Heo DS, Bang YJ, Kim NK. Efficacy of modified regimen with attenuated doses of paclitaxel plus carboplatin combination chemotherapy in elderly and/or weak patients with advanced non-small cell lung cancer. *Lung Cancer* 2003; **39**: 99-101 [PMID: 12499101]
 - 49 **Okamoto I**, Moriyama E, Fujii S, Kishi H, Nomura M, Goto E, Kiyofuji C, Imamura F, Mori T, Matsumoto M. Phase II study of carboplatin-paclitaxel combination chemotherapy in elderly patients with advanced non-small cell lung cancer. *Jpn J Clin Oncol* 2005; **35**: 188-194 [PMID: 15845567]
 - 50 **Marsland TA**, Garfield DH, Khan MM, Look RM, Boehm KA, Asmar L. Sequential versus concurrent paclitaxel and carboplatin for the treatment of advanced non-small cell lung cancer in elderly patients and patients with poor performance status: results of two Phase II, multicenter trials. *Lung Cancer* 2005; **47**: 111-120 [PMID: 15603861]
 - 51 **Inoue A**, Usui K, Ishimoto O, Matsubara N, Tanaka M, Kanbe M, Gomi K, Koinumaru S, Saijo Y, Nukiwa T. A phase II study of weekly paclitaxel combined with carboplatin for elderly patients with advanced non-small cell lung cancer. *Lung Cancer* 2006; **52**: 83-87 [PMID: 16483687]
 - 52 **Pujol JL**, Milleron B, Molinier O, Quoix E, Depierre A, Breton JL, Gervais R, Debieuvre D, Hominal S, Namouni F, Tonelli D. Weekly paclitaxel combined with monthly carboplatin in elderly patients with advanced non-small cell lung cancer: a multicenter phase II study. *J Thorac Oncol* 2006; **1**: 328-334 [PMID: 17409879]
 - 53 **Biesma B**, Wymenga AN, Vincent A, Dalesio O, Smit HJ, Stigt JA, Smit EF, van Felius CL, van Putten JW, Slaets JP, Groen HJ. Quality of life, geriatric assessment and survival in elderly patients with non-small-cell lung cancer treated with carboplatin-gemcitabine or carboplatin-paclitaxel: NVALT-3 a phase III study. *Ann Oncol* 2011; **22**: 1520-1527 [PMID: 21252061]
 - 54 **Quoix E**, Zalcmán G, Oster JP, Westeel V, Pichon E, Lavolé A, Dauba J, Debieuvre D, Souquet PJ, Bigay-Game L, Dansin E, Poudoux M, Molinier O, Vaylet F, Moro-Sibilot D, Herman D, Bennouna J, Tredaniel J, Ducoloné A, Lebitasy MP, Baudrin L, Laporte S, Milleron B. Carboplatin and weekly paclitaxel doublet chemotherapy compared with monotherapy in elderly patients with advanced non-small-cell lung cancer: IFCT-0501 randomised, phase 3 trial. *Lancet* 2011; **378**: 1079-1088 [PMID: 21831418]
 - 55 **Laskin J**, Crinò L, Felip E, Franke F, Gorbunova V, Groen H, Jiang GL, Reck M, Schneider CP. Safety and efficacy of first-line bevacizumab plus chemotherapy in elderly patients with advanced or recurrent nonsquamous non-small cell lung cancer: safety of avastin in lung trial (MO19390). *J Thorac Oncol* 2012; **7**: 203-211 [PMID: 22173662]
 - 56 **Wozniak AJ**, Garst J, Jahanzeb M, Kosty MP, Vidaver R, Beatty S, Teng S, Flick ED, Sing A, Lynch TJ. Clinical outcomes (CO) for special populations of patients (pts) with advanced non-small cell lung cancer (NSCLC): results from ARIES, a bevacizumab (BV) observational cohort study (OCS). *J Clin Oncol* 2013; **28**: abstr 7618
 - 57 **Ochs J**, Grous JJ, Warner KL. Final survival and safety results for 21064 non-small-cell lung cancer (NSCLC) patients who received compassionate use gefitinib in US expanded access program (EAP). *J Clin Oncol* 2004; **22** (14S): 631s (abstr 7060)
 - 58 **Copin M**, Kommareddy A, Behnken D, McLeod H, Trinkaus K, Ali A, Read W, Govindan R. Gefitinib in elderly patients with non-small cell lung cancer (NSCLC). *Proc Am Soc Clin Oncol* 2003; **22**: 758 (abstr 3048)
 - 59 **Cavina R**, Soto Parra HJ, Zucali PA, Campagnoli E, Latteri F, Biancofiore G, Abbadesse G, Morengi E, Santoro A. ZD 1839 (Iressa) in elderly patients with progressive pretreated non-small cell lung cancer (NSCLC): results at the Istituto Clinico Humanitas, Rozzano, Milano. *Lung Cancer* 2003; **41** (suppl 2): S248 (abstr P-618)
 - 60 **Gridelli C**, Maione P, Castaldo V, Rossi A. Gefitinib in elderly and unfit patients affected by advanced non-small-cell lung cancer. *Br J Cancer* 2003; **89**: 1827-1829 [PMID: 14612886]
 - 61 **Cappuzzo F**, Bartolini S, Ceresoli GL, Tambari S, Spreafico A, Lombardo L, Gregorc V, Toschi L, Calandri C, Villa E, Crinò L. Efficacy and tolerability of gefitinib in pretreated elderly patients with advanced non-small-cell lung cancer (NSCLC). *Br J Cancer* 2004; **90**: 82-86 [PMID: 14710211]
 - 62 **Hotta K**, Ueoka H, Kiura K, Tabata M, Ogino A, Umemura S, Harita S, Gemba K, Yonei T, Bessho A, Maeda T, Tanimoto M. Safety and efficacy of gefitinib treatment in elderly patients with non-small-cell lung cancer: Okayama Lung Cancer Study Group experience. *Acta Oncol* 2005; **44**: 717-722 [PMID: 16227162]
 - 63 **Stinchcombe TE**, Bůžková P, Choksi J, Taylor M, Bakri K, Gillenwater H, Tynan M, Mears A, Jones PE, Socinski MA. A phase I/II trial of weekly docetaxel and gefitinib in elderly patients with stage IIIB/IV non-small cell lung cancer. *Lung Cancer* 2006; **52**: 305-311 [PMID: 16638621]
 - 64 **Bepler G**, Williams CC, Chiappori A, Antonia SJ, Haura EB, Mahani JJ, Galloway T, Simon GR. Docetaxel and gefitinib in the first-line treatment of elderly patients (> 70) with advanced non-small cell lung cancer (ANSLC): Results of a phase II trial. *J Clin Oncol* 2005; **23** (16S): 659s (abstr 7155)
 - 65 **Scagliotti G**, Rossi A, Novello S, De Marinis F, Dogliotti L, Crinò L, Lorusso V, Martoni A, Paccagnella A, Gridelli C. Gefitinib (ZD 1839) combined with gemcitabine or vinorelbine as single-agent in elderly patients with advanced non-small-cell lung cancer (NSCLC). *J Clin Oncol* 2004; **23** (14S): 636s (abstr 7081)
 - 66 **Yoshimura M**, Nakamura S, Imamura F, Ueno K, Yamamoto S, Igarashi T. Severe myelotoxicity in a combination of gefitinib and vinorelbine. *Lung Cancer* 2004; **45**: 121-123 [PMID: 15196742]
 - 67 **Jackman D**, Lucca J, Fidias P, Rabin M, Lynch T, Ostler P, Skarin A, Temel J, Johnson B, Janne PA. Phase II study of the EGFR tyrosine kinase inhibitor erlotinib (Tarceva) in patients > 70 years of age with previously untreated advanced non-small cell lung carcinoma. *Lung Cancer* 2005; **49** (suppl 2): S62 (abstr O-188)
 - 68 **Jackman DM**, Yeap B, Lucca J, Ostler PA, Morse LK, Fidias P, Lynch TJ, Temel J, Johnson BE, Janne PA. Phase II trial of erlotinib in elderly patients (age > 70) with previously untreated advanced non-small cell lung cancer (NSCLC): An analysis quality of life and symptom response. *J Clin Oncol* 2006; **24** (18S): 406S (abstr 7168)
 - 69 **Rosell R**, Carcereny E, Gervais R, Vergnenegre A, Massuti B, Felip E, Palmero R, Garcia-Gomez R, Pallares C, Sanchez JM, Porta R, Cobo M, Garrido P, Longo F, Moran T, Insa A, De Marinis F, Corre R, Bover I, Illiano A, Dansin E, de Castro J, Milella M, Reguart N, Altavilla G, Jimenez U, Provencio M, Moreno MA, Terrasa J, Muñoz-Langa J, Valdivia J, Isla D, Domine M, Molinier O, Mazieres J, Baize N, Garcia-Campelo R, Robinet G, Rodriguez-Abreu D, Lopez-Vivanco G, Gebbia V, Ferrera-Delgado L, Bombaron P, Bernabe R, Bearz A, Artal A, Cortesi E, Rolfo C, Sanchez-Ronco M, Drozdowskyj A, Queralt C, de Aguirre I, Ramirez JL, Sanchez JJ, Molina MA, Taron M, Paz-Ares L. Erlotinib versus standard chemotherapy as first-line treatment for European patients with advanced EGFR mutation-positive non-small-cell lung cancer (EURTAC): a multicentre, open-label, randomised phase 3 trial. *Lancet Oncol* 2012; **13**: 239-246 [PMID: 22285168 DOI: 10.1016/S1470-2045(11)70393-X]
 - 70 **Shepherd FA**, Rodrigues Pereira J, Ciuleanu T, Tan EH, Hirsh V, Thongprasert S, Campos D, Maoleekoonpiroj S, Smylie M, Martins R, van Kooten M, Dediu M, Findlay B, Tu D, Johnston D, Bezzak A, Clark G, Santabárbara P, Seymour L. Erlotinib in previously treated

- non-small-cell lung cancer. *N Engl J Med* 2005; **353**: 123-132 [PMID: 16014882]
- 71 **Chen YM**, Tsai CM, Fan WC, Shih JF, Liu SH, Wu CH, Chou TY, Lee YC, Perng RP, Whang-Peng J. Phase II randomized trial of erlotinib or vinorelbine in chemo-naïve, advanced, non-small cell lung cancer patients aged 70 years or older. *J Thorac Oncol* 2012; **7**: 412-418 [PMID: 22157367 DOI: 10.1097/JTO.0b013e31823a39e8]
- 72 **Stinchcombe TE**, Peterman AH, Lee CB, Moore DT, Beaumont JL, Bradford DS, Bakri K, Taylor M, Crane JM, Schwartz G, Hensing TA, McElroy E, Niell HB, Harper HD, Pal S, Socinski MA. A randomized phase II trial of first-line treatment with gemcitabine, erlotinib, or gemcitabine and erlotinib in elderly patients (age \geq 70 years) with stage IIIB/IV non-small cell lung cancer. *J Thorac Oncol* 2011; **6**: 1569-1577 [PMID: 21716146 DOI: 10.1097/JTO.0b013e3182210430]
- 73 **Weiss GJ**, Langer C, Rosell R, Hanna N, Shepherd F, Einhorn LH, Nguyen B, Paul S, McAndrews P, Bunn PA, Kelly K. Elderly patients benefit from second-line cytotoxic chemotherapy: a subset analysis of a randomized phase III trial of pemetrexed compared with docetaxel in patients with previously treated advanced non-small-cell lung cancer. *J Clin Oncol* 2006; **24**: 4405-4411 [PMID: 16983108]
- 74 **Tibaldi C**, Bernardini I, Chella A, Russo F, Vasile E, Malventi M, Falcone A. Second-line chemotherapy with a modified schedule of docetaxel in elderly patients with advanced-stage non-small-cell lung cancer. *Clin Lung Cancer* 2006; **7**: 401-405 [PMID: 16800966]
- 75 **Wu CH**, Fan WC, Chen YM, Chou KT, Shih JF, Tsai CM, Lee YC, Perng RP. Second-line therapy for elderly patients with non-small cell lung cancer who failed previous chemotherapy is as effective as for younger patients. *J Thorac Oncol* 2010; **5**: 376-379 [PMID: 20104191]
- 76 **Repetto L**, Fratino L, Audisio RA, Venturino A, Gianni W, Vercelli M, Parodi S, Dal Lago D, Gioia F, Monfardini S, Aapro MS, Serraino D, Zagonel V. Comprehensive geriatric assessment adds information to Eastern Cooperative Oncology Group performance status in elderly cancer patients: an Italian Group for Geriatric Oncology Study. *J Clin Oncol* 2002; **20**: 494-502 [PMID: 11786579]
- 77 **Stuck AE**, Siu AL, Wieland GD, Adams J, Rubenstein LZ. Comprehensive geriatric assessment: a meta-analysis of controlled trials. *Lancet* 1993; **342**: 1032-1036 [PMID: 8105269]
- 78 **Puts MT**, Hardt J, Monette J, Girre V, Springall E, Alibhai SM. Use of geriatric assessment for older adults in the oncology setting: a systematic review. *J Natl Cancer Inst* 2012; **104**: 1133-1163 [PMID: 22851269]
- 79 **Rossi A**, Maione P, Gridelli C. Safety profile of platinum-based chemotherapy in the treatment of advanced non-small cell lung cancer in elderly patients. *Expert Opin Drug Saf* 2005; **4**: 1051-1067 [PMID: 16255664]
- 80 **Gridelli C**, Aapro M, Ardizzoni A, Balducci L, De Marinis F, Kelly K, Le Chevalier T, Manegold C, Perrone F, Rosell R, Shepherd F, De Petris L, Di Maio M, Langer C. Treatment of advanced non-small-cell lung cancer in the elderly: results of an international expert panel. *J Clin Oncol* 2005; **23**: 3125-3137 [PMID: 15860872]
- 81 **Gajra A**, Jatoi A. Non-small-cell lung cancer in elderly patients: a discussion of treatment options. *J Clin Oncol* 2014; **32**: 2562-2569 [PMID: 25071101]

P- Reviewer: Fang BL, Tamiya M **S- Editor:** Ji FF **L- Editor:** A
E- Editor: Wang CH

