

Risk Factors of Organ Failure in Cholangitis with Bacteriobilia

Jae Min Lee, M.D.^{1,2}, Sang Hyub Lee, M.D., Ph.D.¹, Kwang Hyun Chung, M.D.¹, Jin Myung Park, M.D.³, Ban Seok Lee, M.D.¹, Woo Hyun Paik, M.D., Ph.D.⁴, Joo Kyung Park, M.D., Ph.D.⁵, Ji Kon Ryu, M.D., Ph.D.¹, Yong-Tae Kim, M.D., Ph.D.¹

¹Departments of Internal Medicine and Liver Research Institute, Seoul National University Hospital, Seoul, Korea

²Department of Internal Medicine, Gyeongsang National University College of Medicine, Gyeongsang National University Hospital, Jinju, Korea


³Department of Internal Medicine, Kangwon National University School of Medicine, Kangwon National University Hospital, Chuncheon, Korea

⁴Department of Internal Medicine, Inje University Ilsan Paik Hospital, Goyang, Korea

⁵Department of Internal Medicine, Seoul National University Gangnam Center, Seoul, Korea

Biostatistics: Not applicable (due to the retrospective study using information contained in medical charts and computerized records).

Signature:



Correspondence to: Sang Hyub Lee, M.D., Ph.D.,

Department of Internal Medicine, Seoul National University College of Medicine,

Seoul National University Hospital,

101 Daehak-ro, Jongno-gu, Seoul, 110-744, Korea

Tel: +82-2-2072-4892/Fax: +82-2-762-9662

E-mail: gidoctor@snuh.org