

Institutional Review Board (IRB) statement

This study (No. 08KY023: Different roles of CE and DBE in obscure small intestinal diseases) was reviewed and approved by the Ethics Committee of Sichuan Province Institution * Sichuan Provincial People's Hospital Institutional Review Board.

Ethics Committee
Application for Approval of Research Protocol

1. Title of the Proposed Project

Different role of CE and DBE in the obscure small intestinal diseases

2. Principal and Co-Investigators and Their Departments

Principal Investigator: Zhang Zhihong

Department of Gastroenterology of Sichuan Provincial People's Hospital

Co-Investigators Qiu Chunhua

Department of Gastroenterology of Sichuan Provincial People's Hospital

Li Yi

Department of Gastroenterology of Sichuan Provincial People's Hospital

3. Outline of the Project

Aim of study: To estimate the role and difference between the capsule endoscopy (CE) and the double-balloon enteroscopy (DBE) in the obscure small bowel diseases.

Background: It is very difficult to diagnose the small intestinal diseases because of the small intestinal specific structure and anatomy location. With the development of capsule endoscopy (CE) and the double-balloon enteroscopy (DBE) in recent years, the prospect has been brought for the diagnosis and treatment of the obscure intestinal diseases. However, until now, because of the expensive cost and some difficult technology, the study with CE followed by DBE simultaneously in one case was still rare.

4. Methods: Patients were recruited as followed:

Inclusion criteria: All patients underwent the gastroscopy and colonoscopy, and part of them were given the radiological small intestinal barium meal, abdominal computed tomography (CT) and magnetic resonance imaging (MRI) scan, mesenteric angiography or others. But the causes of them were still not clear.

Exclusion criteria: The contraindications to CE were excluded, which included the gastrointestinal obstruction, fistula, stricture, cardiac pacemakers, pregnant women, and those who could not accept the capsule retention or capsule removal surgery.

This study was reviewed and approved by the ethics committee of Sichuan Provincial People's Hospital Institutional Review Board

5. The ethics committee of Sichuan Provincial People's Hospital
伦理审查批号:

08 KY023



6. References:

1. [Moshe Nadler](#) and [Rami Eliakim](#). The role of capsule endoscopy in acute gastrointestinal bleeding. *Therap Adv Gastroenterol*. Mar 2014; 7(2): 87–92.
2. Triester SL, Leighton JA, Leontiadis GI, et al. A meta-analysis of the yield of capsule endoscopy compared to other diagnostic modalities in patients with obscure gastrointestinal bleeding. *Am J Gastroenterol*. 2005; 100:2407-18.
3. De Leusse A, Vahedi K, Edery J, et al. Capsule endoscopy or push enteroscopy for first-line exploration of obscure gastrointestinal bleeding? *Gastroenterology*. 2007; 132:855-62.
4. Domagk D, Mensink P, Aktas H, Lenz P, Meister T, et al. Single- vs. double-balloon enteroscopy in small-bowel diagnostics: a randomized multicenter trial. *Endoscopy*. 2011; 43:472-476.
5. Teshima CW, Kuipers EJ, van Zanten SV, Mensink PB. Double balloon enteroscopy and capsule endoscopy for obscure gastrointestinal bleeding: an updated meta-analysis. *J Gastroenterol Hepatol*. 2011; 26:796-801.
6. [Kamalaporn P](#)1, [Cho S](#), [Basset N](#), [Cirocco M](#), et al. Double-balloon enteroscopy following capsule endoscopy in the management of obscure gastrointestinal bleeding: outcome of a combined approach. *Can J Gastroenterol*, 2008; 22(5):491-495.
7. [TsujiKawa T](#)1, [Saitoh Y](#), [Andoh A](#), et al. Novel single-balloon enteroscopy for diagnosis and treatment of the small intestine: preliminary experiences. *Endoscopy*. 2008 ; 40(1):11-5.
8. [Takano N](#)1, [Yamada A](#), [Watabe H](#), et al. Single-balloon versus double-balloon endoscopy for achieving total enteroscopy: a randomized, controlled trial. *Gastrointest Endosc*. 2011; 73(4):734-9.

9. Carey EJ, Leighton JA, Heigh RI, et al. A single-center experience of 260 consecutive patients undergoing capsule endoscopy for obscure gastrointestinal bleeding. *Am J Gastroenterol*. 2007 Jan; 102(1):89-95.
10. Pennazio M, Santucci R, Rondonotti E, et al. Outcome of patients with obscure gastrointestinal bleeding after capsule endoscopy: report of 100 consecutive cases. *Gastroenterology*. 2004 Mar; 126(3):643-53.
11. [Zhang Q](#), [He Q](#), [Liu J](#), et al. Combined use of capsule endoscopy and double-balloon enteroscopy in the diagnosis of obscure gastrointestinal bleeding: meta-analysis and pooled analysis. *Hepatogastroenterology*. 2013; 60(128):1885-91.
12. [Lewis B](#). How to prevent endoscopic capsule retention. *Endoscopy*. 2005 Sep; 37(9):852-6.
13. [Sears DM](#)¹, [Avots-Avotins A](#), [Culp K](#), [Gavin MW](#). Frequency and clinical outcome of capsule retention during capsule endoscopy for GI bleeding of obscure origin. *Gastrointest Endosc*. 2004; 60(5):822-7.
14. Mensink PB, Haringsma J, Kucharzik T, et al. Complications of double balloon enteroscopy: a multicenter survey [J]. *Endoscopy*, 2007, 39 (7) : 613-615.
15. Application of Capsule Endoscopy and Device-assisted Enteroscopy in Detection of Small Bowel Tumors: A Systematic Review. LI Xuan, GE Zhizheng. *Chin J Gastroenterol*, 2013, 18(7): 396-401.
16. [Cazzato IA](#)¹, [Cammarota G](#), [Nista EC](#), et al. Diagnostic and therapeutic impact of double-balloon enteroscopy (DBE) in a series of 100 patients with suspected small bowel diseases. *Dig Liver Dis*. 2007 May;39(5):483-7.
17. [Rahmi G](#)¹, [Samaha E](#), [Vahedi K](#), et al. Multicenter comparison of double-balloon enteroscopy and spiral enteroscopy. *J Gastroenterol Hepatol*. 2013, 28(6):992-8.