



BAISHIDENG PUBLISHING GROUP INC

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242 Fax: +1-925-223-8243

E-mail: bpgoffice@wjgnet.com <http://www.wjgnet.com>

Name of journal: *World Journal of Clinical Cases*

ESPS Manuscript NO: 19693

Manuscript Type: CASE REPORT

Reviewer 1 (Reviewer No. 00214259):

Isolated cardiac sarcoidosis is very rare. Do we know the prevalence of this clinical entity?

We agree that cardiac sarcoidosis is very rare. The estimated prevalence of sarcoidosis varies from ethnicity (4.7-64/100 000). Approximately, 2-7% of patients with sarcoidosis suffer from clinical cardiac manifestations. However, autopsy studies reveal a higher prevalence up to 70-85%. We added this information in the revised manuscript on page 4, lines 2-7.

This case report is support mainly on the endomyocardial biopsy (EMB) results. Do we know the sensibility and specificity of EMB for the diagnostic of cardiac sarcoidosis? Do you suggest that we should perform left ventricular EMB for every case of myocarditis when the etiology of the myocardial injury remained unclear?

Thank you very much for pointing us to this important issue. The sensitivity and specificity of EMB in cardiac sarcoidosis are 20-30, whereas the specificity is unknown. However, CMR could be used to perform targeted EMB in order to improve the diagnostic accuracy. We added this information in the revised manuscript on page 6, lines 7-10. Furthermore, we clarified the role of EMB in current recommendations in the revised manuscript on page 6, lines 10-15.

Reviewer 2 (Reviewer No. 00106145):

Reduced systolic function must be specified. Please indicate ejection fraction, LV volumes and regional wall motion, etc. Preferable, also diastolic filling echo variables along with the

mechanism of mitral regurgitation should be specified. Echo functional status should better be defined also regarding follow-up.

We thank the reviewer for his helpful recommendations, which definitely helped to improve the manuscript.

Thank you very much, we added the missing quantitative values at baseline and follow-up as well as the mechanism of mitral regurgitation, which was functional MR due to the dilated annulus and papillary muscle dysfunction, in the revised manuscript on page 4, lines 21-26.

2) MR images are striking. Echo speckle tracking is not described, but should be added to the discussion as a diagnostic tool.

Thank you very much. Speckle tracking is certainly an important novel feature but was not performed in this case. However, we entered speckle tracking as a diagnostic tool in the revised manuscript on page 5, lines 28-20 and page 6 lines 1-7.

English requires minor revision (e.g.: "Interestingly, additionally laboratory findings...").

Thank you very much, we checked the revised manuscript for English grammar.