

Firstly, I would like to thank each of the reviewers for their hard work in reading, critiquing, and offering their suggestions to help make this a better and more informative paper. Your comments were helpful and I have made efforts to incorporate each of your suggestions. Specifics are listed below.

Reviewer's Code: **00503313**

Revision: Per this reviewer, there were no revisions to be made. However, as they commented on the infrequently used trans-jugular approach for kidney biopsy, this was clarified in a revision. The traditional percutaneous method of obtaining a renal biopsy is equally efficacious. However, if obtaining a liver and kidney biopsy at the same time in a facility with the means, a trans-jugular approach would allow for sampling of both tissues. This addition was highlighted yellow.

Reviewer's Code: **00503339**

Revision: Per this reviewer, a chart indicating the creatinine and bilirubin values over time as well as latest values would be helpful. This was addressed by the inclusion of 2 new tables. Table 1 includes the creatinine and bilirubin values over time, as well as when particular interventions were performed, such as dialysis or plasmapheresis. Table 2 includes the most recent available creatinine and bilirubin values. This addition was highlighted yellow.

Reviewer's Code: **00503241**

Revision: Per this reviewer, (A) the discussion regarding extracorporeal treatment options for removal of bilirubin was incomplete as other potential treatment options were not discussed. (B) The presence of lack of albuminuria was not included in the original manuscript. (C) the trans-jugular approach for renal biopsy at the time of obtaining liver biopsy is not routine. These additions were highlighted in yellow.

- A) Alternative extracorporeal treatment options and their uses were incorporated into the manuscript along with generalized information on improved outcomes in a few studies.
- B) Urinalysis information, including the presence of trace proteinuria was added into the manuscript.

C) Discussion regarding obtaining kidney biopsy via the traditional percutaneous method is acceptable was added, and clarified to state that the trans-jugular approach to obtaining a renal biopsy at the time of obtaining a liver biopsy may be executed in a facility with the means and abilities to do so.