

Name of Journal: *World Journal of Hepatology*

Manuscript NO: 31926

Manuscript Type: Original Article

Basic Study

Angiotensin II or epinephrine hemodynamic and metabolic responses in the liver of L-NAME induced hypertension and spontaneous hypertensive rats

Debora Conte Kimura, Marcia Regina Nagaoka, Durval Rosa Borges, Maria Kouyoumdjian

Abstract

AIM: Hepatic vasoconstriction and glucose release induced by AngII or Epi were studied in rats with pharmacological hypertension and SHR.

METHODS: Isolated liver perfusion was performed following portal vein and vena cava cannulation; AngII or Epi was injected *in bolus* and portal pressure monitored;

Match Overview

Rank	Source	Words	Similarity
1	Internet crawled on 22-Mar-2016 www.clinsci.org	43 words	1%
2	Crossref Christiane Aparecida Badin Tarsitano. "Hepatic morphological alterations, glycogen content and cytochrome P450 act	39 words	1%
3	Internet crawled on 13-May-2016 linknovate.com	34 words	1%
4	Crossref C. B. Herath. "Portal pressure responses and angiotensin peptide production in rat liver are determined by relative ...	33 words	1%
5	Publications "Researchers from University of Maryland Report on Findings in Autacoids (Novel RAAS agonists and ant", Biotech	33 words	1%
6	Crossref H KOBORI. "Young Scholars Award Lecture: Intratubular ... ngiotensinogen in Hypertension and Kidney Diseases", Am	30 words	1%
7	Crossref Waleska C Damas. "Animal models for the study of arterial	26 words	1%



[全部](#) [图片](#) [购物](#) [新闻](#) [视频](#) [更多](#)

[设置](#) [工具](#)

找到约 10,500 条结果 (用时 0.95 秒)

[Abstracts of the 9th International Symposium on SHR and ...](#)
onlinelibrary.wiley.com/doi/10.1111/j.1440-1681.1998.tb02354.../abstract - [翻译此页](#)

ORAL PRESENTATIONS: ROLE OF ENDOGENOUS ANGIOTENSIN II IN THE ... POSTER
PRESENTATIONS: ANALYSIS OF HEMODYNAMIC AND PLASMA RENIN ... L-NAME-INDUCED
HYPERTENSION IN UNINEPHRECTOMIZED RATS. ... RESPONSES TO ANGIOTENSIN H IN
SPONTANEOUSLY HYPERTENSIVE ...

[Angiotensin II pressor response in the L-NAME-induced hypertensive ...](#)
<https://www.ncbi.nlm.nih.gov/pubmed/22128423> - [翻译此页](#)

作者: ES Figueroa-Guillén - 2009 - [被引用次数: 1](#) - [相关文章](#)

Angiotensin II pressor response in the L-NAME-induced hypertensive pithed rat. ... II type 1 (AT1)
receptor role in L-NAME-induced hypertension in pithed rats.

缺少字词: epinephrine hemodynamic metabolic liver spontaneous

[Influence of Fimasartan \(a Novel AT1 Receptor Blocker\) on ... - NCBI](#)
<https://www.ncbi.nlm.nih.gov> > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: HJ Lim - 2013 - [被引用次数: 1](#) - [相关文章](#)

2013年2月14日 - It is the eighth drug marketed as a member of the Ang II receptor ... in the treatment
of hypertension, with two classes of antihypertensive agents targeting the RAAS. ... In
spontaneously hypertensive rats (SHRs), oral administration of the effect of L-NAME on
fimasartan-induced inhibitory responses of CA ...

[Physiopathology of splanchnic vasodilation in portal hypertension](#)
<https://www.ncbi.nlm.nih.gov> > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: M Martell - 2010 - [被引用次数: 61](#) - [相关文章](#)

2010年6月27日 - In liver cirrhosis, the circulatory hemodynamic alterations of portal hypertension ...
and an impaired vasodilatory response of the hepatic vascular bed, are the by eNOS in
mesenteric vessels from portal hypertensive rats[87,88]. ... with cirrhosis, inducing desensitisation of
angiotensin II receptor[111].

找到约 24,400 条结果 (用时 0.39 秒)

[PDF] Effects of COX-1 and COX-2 Inhibitors in L - Jordan Journal of ...

jbs.hu.edu.jo/files/v6n3/Paper%20Number%204m.pdf  翻译此页

作者: IM Maulood - 被引用次数: 1 - 相关文章

Arginine Methyl Ester Induced Hypertensive Rats ... COX-2 group rats compared with the L-NAME group. ... hypertension may be due to vasoconstrictor effect of ... may due to increase in plasma epinephrine and ... important roles in the cellular response to stress ... possible hemodynamic, renal and liver effects of COX-1.

Influence of Fimasartan (a Novel AT1 Receptor Blocker) on ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: HJ Lim - 2013 - 被引用次数: 1 - 相关文章

2013年2月14日 - Generally, the renin-angiotensin II-aldosterone system (RAAS) is an important ... In spontaneously hypertensive rats (SHRs), oral administration of AT1 [37] have suggested that Ang II increase epinephrine release from the adrenal ... the effect of L-NAME on fimasartan-induced inhibitory responses of CA ...

Brain mineralocorticoid receptors in cognition and cardiovascular ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: E Gomez-Sanchez - 2014 - 被引用次数: 12 - 相关文章

11 β -HSD1 inhibitors combat metabolic and cognitive diseases related to The role of brain MR in mineralocorticoid-salt hypertension was confirmed by the lateral ... for MR abrogated the increased SNS activation and hypertension induced by ... the L-NAME, Angiotensin II-salt excess, mineralocorticoid-salt excess, and ...

Tetrahydrobiopterin in Cardiovascular Health and Disease - NCBI

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4038990/> - 翻译此页

作者: JK Bendall - 2014 - 被引用次数: 47 - 相关文章

In the liver, BH4 was shown to be an essential cofactor for the metabolism of ... synthesize NO

[全部](#)[图片](#)[新闻](#)[购物](#)[视频](#)[更多](#)[设置](#)[工具](#)

找到约 24,100 条结果（用时 0.90 秒）

[PDF] [Effects of COX-1 and COX-2 Inhibitors in L - Jordan Journal of ...](#)

jbs.hu.edu.jo/files/v6n3/Paper%20Number%204m.pdf ▾ [翻译此页](#)

作者: IM Maulood - 被引用次数: 1 - [相关文章](#)

Arginine Methyl Ester **Induced Hypertensive Rats** ... COX-2 group **rats** compared with the **L-NAME** group. ... **hypertension** may be due to vasoconstrictor effect of ... may due to increase in plasma **epinephrine** and ... important roles in the cellular **response** to stress ... possible **hemodynamic**, renal and **liver** effects of COX-1.

[Influence of Fimasartan \(a Novel AT1 Receptor Blocker\) on ...](#)

<https://www.ncbi.nlm.nih.gov> > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: HJ Lim - 2013 - 被引用次数: 1 - [相关文章](#)

2013年2月14日 - Generally, the renin-**angiotensin II**-aldosterone system (RAAS) is an important ... In **spontaneously hypertensive rats** (SHRs), oral administration of AT1 [37] have suggested that Ang II increase **epinephrine** release from the adrenal ... the effect of **L-NAME** on **fimasartan-induced** inhibitory **responses** of CA ...

[Brain mineralocorticoid receptors in cognition and cardiovascular ...](#)

<https://www.ncbi.nlm.nih.gov> > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: E Gomez-Sanchez - 2014 - 被引用次数: 12 - [相关文章](#)

The r o le of brain MR in mineralocorticoid-salt **hypertension** was confirmed by the ... for MR abrogated the increased SNS activation and **hypertension induced** by ... the **L-NAME**, **Angiotensin II**-salt excess, mineralocorticoid-salt excess, and of the lower affinity Type II receptors dampened the **responses** to stress; severe or ...

[PDF] [Full text \(PDF file\) - biomed.cas.cz](#)

www.biomed.cas.cz/physiolres/pdf/57/57_37P.pdf ▾ [翻译此页](#)