



32593-Review

Quotes Included
Bibliography Included2%
SIMILAR

Name of Journal: *World Journal of Methodology*

Manuscript NO: 32593

Manuscript Type: Case Report

Laparoscopic-extracorporeal surgery performed with a fixation device for
adnexal masses complicating pregnancy: Report of two cases

Hanako Kasahara, Iwaho Kikuchi, Aya Otsuka, Yoko Tsuzuki, Michio
Nojima, Koyo Yoshida

Abstract

The potential complications associated with an adnexal mass discovered during

Match Overview

1

Crossref 24 words

Tsuguhiro Horikoshi. "Two cases of a fetus with sirenomelia s
equence", *Congenital Anomalies*, 9/2005

2%

[全部](#)
[图片](#)
[新闻](#)
[视频](#)
[购物](#)
[更多](#)
[设置](#)
[工具](#)

找到约 107,000 条结果 (用时 0.67 秒)

Laparoscopic surgery for treating adnexal masses during the first ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: L Minig - 2016 - 被引用次数: 8 - [相关文章](#)

The main indication of **surgery** was cyst torsion in four **cases** (30.7%) and rupture of ... cyst in three patients (23%) and heterotopic pregnancy in **two cases** (15.3%), ... The incidence of **adnexal masses during pregnancy** can range between 1:81 Dorsal lithotomy position for **surgery performed** during the first trimester and a ...

缺少字词: **extracorporeal fixation device**

Global Library of Women's Medicine - Operative Laparoscopy - DOI ...

editorial.glowm.com/ch053/framesets/index.html?p=glowm.cml/...88 ▾ [翻译此页](#)

In the 1900s, one of the first **reports** on operative **laparoscopy** came from **Two** such **devices** include those produced by Gyrus/ACMI and Covidian In patients who have had previous abdominal **surgery**, in children, and in **pregnant** patients, Of the 54,198 laparoscopies, 16,601 were **performed** for **adnexal masses**; ...

Single-port access laparoscopic surgery using a novel laparoscopic ...

[www.tjog-online.com/article/S1028-4559\(11\)00166-5/fulltext](http://www.tjog-online.com/article/S1028-4559(11)00166-5/fulltext) - [翻译此页](#)

作者: T Song - 2011 - 被引用次数: 14 - [相关文章](#)

2011年2月16日 - All procedures were **performed** without complications. ... Octo-Port, a single-port access **laparoscopic device**, showing (A) a ... body **mass** index (BMI, kg/m²), chief complaint, estimated blood loss, operation time, and operative record. ... In the **cases** of SPA **laparoscopic adnexal surgery**, we used several ...

Laparoscopy versus laparotomy in the management of adnexal ...

[www.journals.elsevierhealth.com/article/S0015-0282\(99\)00064-3/fulltext](http://www.journals.elsevierhealth.com/article/S0015-0282(99)00064-3/fulltext) ▾ [翻译此页](#)

作者: D Soriano - 1999 - 被引用次数: 162 - [相关文章](#)

Laparoscopy versus laparotomy in the management of **adnexal masses** ... Numerous **case reports**

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 82,100 条结果 (用时 0.53 秒)

Laparoscopic surgery for treating adnexal masses during the first ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4746970/> ▼ [翻译此页](#)

作者: L Minig - 2016 - 被引用次数: 10 - 相关文章

The main indication of surgery was cyst torsion in four cases (30.7%) and rupture of ... The treatment of complicated adnexal masses by laparoscopic surgery during the first ... [2] The most common causes of ovarian masses during pregnancy include ... [3] Even though surgery has been traditionally performed by laparotomy, ...

缺少字词: extracorporeal fixation device

Laparoscopy versus laparotomy in the management of adnexal ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/10231065> ▼ [翻译此页](#)

作者: D Soriano - 1999 - 被引用次数: 170 - 相关文章

Laparoscopy versus laparotomy in the management of adnexal masses during pregnancy. ... outcome of pregnancy after operative laparoscopy or laparotomy for the management ... SETTING: University tertiary care referral center for endoscopic surgery. ... Laparoscopy was performed during the first trimester in 39 patients.

缺少字词: extracorporeal fixation device report

Surgical Management of Adnexal Masses in Pregnancy - NCBI - NIH

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3939346/> ▼ [翻译此页](#)

作者: SF Ngu - 2014 - 被引用次数: 19 - 相关文章

A retrospective review was performed of pregnant women before 20 weeks of ... Conversion to laparotomy was required in 2 women because of extensive pelvic ... The laparoscopy group had a significantly less blood loss (67.4 ± 55.8 vs 153.6 ... Surgical management of adnexal masses during pregnancy appears to have ...

缺少字词: extracorporeal fixation device